

Appendix I

High Priority Site Subwatershed Analysis

(2008-2010 and 2010-2012)

Table of Content

HIGH PRIORITY SITE SUBWATERSHEDS (2008 -2010)	1
I. DRY CREEK @ WELLSFORD RD	1
MANAGEMENT PLAN CONSTITUENTS.....	1
DESCRIPTION OF DRY CREEK AT WELLSFORD RD SITE SUBWATERSHED	1
SUBWATERSHED MONITORING HISTORY.....	3
EXCEEDANCE HISTORY	9
2007 - 2009 MANAGEMENT PLAN MONITORING RESULTS.....	13
LOAD CALCULATIONS.....	15
SOURCE IDENTIFICATION AND OUTREACH	16
EVALUATION	34
II. DUCK SLOUGH @ HIGHWAY 99	35
MANAGEMENT PLAN CONSTITUENTS.....	35
DESCRIPTION OF DUCK SLOUGH @ HWY 99 SITE SUBWATERSHED	35
SUBWATERSHED MONITORING HISTORY.....	37
EXCEEDANCE HISTORY	43
2007 - 2009 MANAGEMENT PLAN MONITORING RESULTS.....	46
LOAD CALCULATIONS.....	48
SOURCE IDENTIFICATION AND OUTREACH	48
EVALUATION	60
III. PRAIRIE FLOWER DRAIN	61
MANAGEMENT PLAN CONSTITUENTS.....	61
DESCRIPTION OF PRAIRIE FLOWER DRAIN @ CROWS LANDING RD SITE SUBWATERSHED	61
SUBWATERSHED MONITORING HISTORY.....	63
EXCEEDANCE HISTORY	69
2007 - 2009 MANAGEMENT PLAN MONITORING RESULTS.....	74
LOAD CALCULATIONS.....	75
SOURCE IDENTIFICATION AND OUTREACH	76
EVALUATION	82
HIGH PRIORITY SITE SUBWATERSHEDS (2010 -2012)	83
IV. BEAR CREEK @ KIBBY RD	83
MANAGEMENT PLAN CONSTITUENTS.....	83
DESCRIPTION OF BEAR CREEK @ KIBBY RD SITE SUBWATERSHED	83
SUBWATERSHED MONITORING HISTORY.....	85
EXCEEDANCE HISTORY	90
2008 MANAGEMENT PLAN MONITORING RESULTS	93
LOAD CALCULATIONS.....	93
SOURCE IDENTIFICATION AND OUTREACH	94
EVALUATION	102
V. COTTONWOOD CREEK @ RD 20	104
MANAGEMENT PLAN CONSTITUENTS.....	104
DESCRIPTION OF COTTONWOOD CREEK @ Rd 20 SITE SUBWATERSHED.....	104
SUBWATERSHED MONITORING HISTORY.....	106
EXCEEDANCE HISTORY	111

2007 - 2009 MANAGEMENT PLAN MONITORING RESULTS.....	114
LOAD CALCULATIONS.....	116
SOURCE IDENTIFICATION AND OUTREACH	117
EVALUATION	139
VI. DUCK SLOUGH @ GURR RD	141
MANAGEMENT PLAN CONSTITUENTS.....	141
DESCRIPTION OF DUCK SLOUGH @ GURR RD SITE SUBWATERSHED.....	141
SUBWATERSHED MONITORING HISTORY.....	143
EXCEEDANCE HISTORY	148
2007 - 2009 MANAGEMENT PLAN MONITORING RESULTS.....	151
LOAD CALCULATIONS.....	153
SOURCE IDENTIFICATION AND OUTREACH	154
EVALUATION	168
VII. HIGHLINE CANAL @ HWY 99	169
MANAGEMENT PLAN CONSTITUENTS.....	169
DESCRIPTION OF HIGHLINE CANAL @ HWY 99 SITE SUBWATERSHED.....	169
SUBWATERSHED MONITORING HISTORY.....	171
EXCEEDANCE HISTORY	176
2007 - 2009 MANAGEMENT PLAN MONITORING RESULTS.....	179
LOAD CALCULATIONS.....	180
SOURCE IDENTIFICATION AND OUTREACH	181
EVALUATION	196
LOW PRIORITY SUBWATERSHEDS.....	198
MERCED RIVER @ SANTA FE DRIVE	198
DRY CREEK @ OAKDALE RD (UPSTREAM OF MERCED RIVER)	198
MILES CREEK @ REILLY RD.....	199

List of Tables

TABLE I-1. COORDINATES OF THE DRY CREEK SITE SUBWATERSHED SAMPLING LOCATIONS.	1
TABLE I-2. DRY CREEK @ WELLSFORD RD SAMPLING EVENTS PER SEASON AND YEAR.	3
TABLE I-3. NUMBER OF ANALYSES PERFORMED PER ANALYTE IN EACH SAMPLING SEASON AND YEAR FOR THE DRY CREEK @ WELLSFORD RD SAMPLE SITE. ONLY ENVIRONMENTAL SAMPLES WITH A SAMPLE REPLICATE AND LAB REPLICATE NUMBER OF ONE ARE SHOWN.	4
TABLE I-4. DRY CREEK SITE SUBWATERSHED. 2007 MANAGEMENT PLAN ADDITIONAL (A) SAMPLING SCHEDULE FOR CHLORPYRIFOS AND <i>CERIODAPHNIA DUBIA</i> . "X" INDICATES THE SITE, MONTH, AND ANALYTE SAMPLED.	8
TABLE I-5. DRY CREEK SITE SUBWATERSHED. 2008 MANAGEMENT PLAN UPSTREAM (U) SAMPLING SCHEDULE FOR CHLORPYRIFOS AND <i>CERIODAPHNIA DUBIA</i> . "X" INDICATES THE SITE, MONTH, AND ANALYTE SAMPLED.	8
TABLE I-6. DRY CREEK SITE SUBWATERSHED. 2009 MANAGEMENT PLAN MONITORING (MPM) AND UPSTREAM (U) SAMPLING SCHEDULE FOR CHLORPYRIFOS. "X" INDICATES THE SITE, MONTH AND ANALYTE SAMPLED.	8
TABLE I-7. ALL EXCEEDANCES EXPERIENCED IN SAMPLES COLLECTED FROM LOCATIONS WITHIN THE DRY CREEK SITE SUBWATERSHED FROM FEBRUARY 2005 THROUGH DECEMBER 2009 (SORTED BY SEASON AND DATE). IF THE WATER QUALITY TRIGGER LIMIT IS BASED ON HARDNESS THEN THE HARDNESS VALUE IS SHOWN IN PARENTHESIS; OTHERWISE THE WQTL USED TO EVALUATE THE DATA IS LISTED IN THE HEADER AFTER THE ANALYTE. UPSTREAM SITE IS ITALICIZED.	10
TABLE I-8. DRY CREEK SITE SUBWATERSHED. NORMAL MONITORING (NM) AND MANAGEMENT PLAN MONITORING (MPM) RESULTS WHERE 'A' INDICATES ADDITIONAL MPM (2007) AND 'US' INDICATES UPSTREAM MPM (2008) FOR CHLORPYRIFOS AND <i>CERIODAPHNIA</i> FROM THE 2007-2008 IRRIGATION SEASONS, AND NM AND MPM FOR CHLORPYRIFOS FOR THE IRRIGATION SEASONS OF 2009. EXCEEDANCE VALUES ARE IN BOLD.	14
TABLE I-9. DRY CREEK @ WELLSFORD AND DRY CREEK @ WATERFORD. INSTANTANEOUS LOAD CALCULATIONS FOR CHLORPYRIFOS, COPPER AND DIURON WHERE DISCHARGE WAS MEASURED (SORTED BY ANALYTE, SITE AND DATE). UPSTREAM SITE IS ITALICIZED.	15
TABLE I-10. NUMBER OF CHLORPYRIFOS APPLICATIONS, POUNDS AI APPLIED, AND ACRES TREATED BY MONTH FOR JANUARY 2005 THROUGH DECEMBER 2009 IN THE DRY CREEK @ WELLSFORD RD SITE SUBWATERSHED. IF A MONTH IS NOT INCLUDED IN THE TABLE, NO APPLICATIONS WERE MADE.	17
TABLE I-11. TOTAL POUNDS OF AI FOR CHLORPYRIFOS BASED ON PUR DATA FROM 2005-2009 WITHIN THE DRY CREEK @ WELLSFORD RD SUBWATERSHED.	20
TABLE I-12. NUMBER OF COPPER APPLICATIONS, TOTAL POUNDS AI APPLIED AND TOTAL ACRES TREATED BY MONTH FOR JANUARY 2005 THROUGH DECEMBER 2009 IN THE DRY CREEK @ WELLSFORD SITE SUBWATERSHED. IF A MONTH IS NOT INCLUDED IN THE TABLE, NO APPLICATIONS WERE MADE.	23
TABLE I-13. TOTAL POUNDS OF AI FOR COPPER BASED ON PUR DATA FROM 2005-2009 WITHIN THE DRY CREEK @ WELLSFORD RD SUBWATERSHED.	26

TABLE I-14. NUMBER OF DIURON APPLICATIONS, TOTAL POUNDS AI APPLIED AND TOTAL ACRES TREATED BY MONTH FOR JANUARY 2005-DECEMBER 2009 IN THE DRY CREEK @ WELLSFORD RD SITE SUBWATERSHED. IF A MONTH IS NOT INCLUDED IN THE TABLE, NO APPLICATIONS WERE MADE.	29
TABLE I-15. TOTAL POUNDS OF AI FOR DIURON BASED ON PUR DATA FROM 2005-2009 WITHIN THE DRY CREEK @ WELLSFORD RD SUBWATERSHED.	32
TABLE II-1. COORDINATES OF THE DUCK SLOUGH SITE SUBWATERSHED SAMPLING LOCATIONS.	35
TABLE II-2. DUCK SLOUGH @ HWY 99 SAMPLING EVENTS PER SEASON AND YEAR.....	37
TABLE II-3. NUMBER OF ANALYSES PERFORMED PER ANALYTE IN EACH SAMPLING SEASON AND YEAR FOR THE DUCK SLOUGH @ HWY 99 SAMPLE SITE. ONLY ENVIRONMENTAL SAMPLES WITH A SAMPLE REPLICATE AND LAB REPLICATE NUMBER OF ONE ARE SHOWN.	38
TABLE II-4. DUCK SLOUGH SITE SUBWATERSHED. 2007 MANAGEMENT PLAN ADDITIONAL (A) SAMPLING SCHEDULE FOR CHLORPYRIFOS AND COPPER. "X" INDICATES THE SITE, MONTH, AND ANALYTE SAMPLED.....	42
TABLE II-5. DUCK SLOUGH SITE SUBWATERSHED. 2008 MANAGEMENT PLAN UPSTREAM (U) SAMPLING SCHEDULE FOR CHLORPYRIFOS AND COPPER. X INDICATES THE SITE, MONTH AND ANALYTE SAMPLED.	42
TABLE II-6. DUCK SLOUGH SITE SUBWATERSHED. 2009 MANAGEMENT PLAN SAMPLING SCHEDULE FOR CHLORPYRIFOS, COPPER AND <i>SELENASTRUM CAPRICORNUTUM</i> TOXICITY. "X" INDICATES THE SITE, MONTH AND ANALYTE SAMPLED.	42
TABLE II-7. ALL EXCEEDANCES EXPERIENCED IN SAMPLES COLLECTED FROM LOCATIONS WITHIN THE DUCK SLOUGH SITE SUBWATERSHED BETWEEN JULY 2004 AND DECEMBER 2009 (SORTED BY SEASON AND DATE). IF THE WATER QUALITY TRIGGER LIMIT IS BASED ON HARDNESS THEN THE HARDNESS VALUE IS SHOWN IN PARENTHESIS; OTHERWISE THE WQTL USED TO EVALUATE THE DATA IS LISTED IN THE HEADER AFTER THE ANALYTE.	44
TABLE II-8. DUCK SLOUGH SITE SUBWATERSHED. NORMAL MONITORING (NM) AND MANAGEMENT PLAN MONITORING (MPM) RESULTS WHERE 'A' INDICATES ADDITIONAL MPM (2007), AND 'US' INDICATES UPSTREAM MPM FOR CHLORPYRIFOS (2008), COPPER (2007-2009 IRRIGATION SEASONS) AND <i>SELENASTRUM</i> (2009). EXCEEDANCE VALUES ARE IN BOLD.	47
TABLE II-9. DUCK SLOUGH SITE SUBWATERSHED. INSTANTANEOUS LOAD CALCULATIONS FOR CHLORPYRIFOS AND COPPER WHERE DISCHARGE WAS MEASURED (SORTED BY ANALYTE, SITE AND DATE). UPSTREAM SITE IS IN ITALICS.	48
TABLE II-10. NUMBER OF CHLORPYRIFOS APPLICATIONS, TOTAL POUNDS APPLIED, AND TOTAL ACRES TREATED BY MONTH FOR JANUARY 2005 THROUGH DECEMBER 2009 IN THE DUCK SLOUGH @ HWY 99 SITE SUBWATERSHED. IF A MONTH IS NOT INCLUDED IN THE TABLE, NO APPLICATIONS WERE MADE.	49
TABLE II-11. TOTAL POUNDS AI FOR CHLORPYRIFOS BASED ON PUR DATA FROM 2005-2009 WITHIN THE DUCK SLOUGH @ HWY 99 SUBWATERSHED.	52
TABLE II-12. NUMBER OF COPPER APPLICATIONS, TOTAL POUNDS APPLIED AND TOTAL ACRES TREATED BY MONTH FOR JANUARY 2005-DECEMBER 2009 IN THE DUCK SLOUGH @ HWY 99 SITE SUBWATERSHED. IF A MONTH IS NOT INCLUDED IN THE TABLE, NO APPLICATIONS WERE MADE.....	54

TABLE II-13. TOTAL POUNDS AI FOR COPPER BASED ON PUR DATA FROM 2005-2009 WITHIN THE DUCK SLOUGH @ HWY 99 SUBWATERSHED.	57
TABLE III-1. COORDINATES OF THE PRAIRIE FLOWER DRAIN SITE SUBWATERSHED SAMPLING LOCATIONS.	61
TABLE III-2. PRAIRIE FLOWER DRAIN @ CROWS LANDING RD SAMPLING EVENTS PER SEASON AND YEAR.....	63
TABLE III-3. NUMBER OF ANALYSES PERFORMED PER ANALYTE IN EACH SAMPLING SEASON AND YEAR FOR THE PRAIRIE FLOWER DRAIN @ CROWS LANDING RD SAMPLE SITE. ONLY ENVIRONMENTAL SAMPLES WITH A SAMPLE REPLICATE AND LAB REPLICATE NUMBER OF ONE ARE SHOWN.....	64
TABLE III-4. PRAIRIE FLOWER DRAIN @ CROWS LANDING RD. 2007 MANAGEMENT PLAN ADDITIONAL (A) SAMPLING SCHEDULE FOR CHLORPYRIFOS AND <i>PIMEPHALES</i> TOXICITY. "X" INDICATES THE SITE, MONTH AND ANALYTE SAMPLED.	68
TABLE III-5. PRAIRIE FLOWER DRAIN SITE SUBWATERSHED. 2008 MANAGEMENT PLAN UPSTREAM (U) SAMPLING SCHEDULE FOR CHLORPYRIFOS, NITRATE, <i>CERIODAPHNIA</i> TOXICITY, AND <i>PIMEPHALES</i> TOXICITY. "X" INDICATES THE SITE, MONTH AND ANALYTE SAMPLED.	68
TABLE III-6. PRAIRIE FLOWER DRAIN SITE SUBWATERSHED. 2009 MANAGEMENT PLAN SAMPLING SCHEDULE FOR CHLORPYRIFOS AND <i>SELENASTRUM CAPRICORNUTUM</i> TOXICITY. "X" INDICATES THE SITE, MONTH AND ANALYTE SAMPLED.	68
TABLE III-7. ALL EXCEEDANCES EXPERIENCED IN SAMPLES COLLECTED FROM LOCATIONS WITHIN THE PRAIRIE FLOWER DRAIN SITE SUBWATERSHED BETWEEN FEBRUARY 2005 AND DECEMBER 2009 (SORTED BY SEASON AND DATE). IF THE WATER QUALITY TRIGGER LIMIT IS BASED ON HARDNESS THEN THE HARDNESS VALUE IS SHOWN IN PARENTHESIS; OTHERWISE THE WQTL USED TO EVALUATE THE DATA IS LISTED IN THE HEADER AFTER THE ANALYTE.....	70
TABLE III-8. PRAIRIE FLOWER DRAIN SITE SUBWATERSHED. NORMAL MONITORING (NM) AND MANAGEMENT PLAN MONITORING (MPM) RESULTS WHERE 'A' INDICATES ADDITIONAL MPM (2007), 'US' INDICATES UPSTREAM MPM FOR CHLORPYRIFOS (2008-2009), NITRATE, <i>CERIODAPHNIA</i> , AND <i>PIMEPHALES</i> FROM THE 2007-2008 IRRIGATION SEASONS, AND MPM FOR <i>SELENASTRUM</i> (2009). EXCEEDANCE VALUES ARE IN BOLD.....	75
TABLE III-9. PRAIRIE FLOWER DRAIN SITE SUBWATERSHED. INSTANTANEOUS LOAD CALCULATIONS FOR CHLORPYRIFOS WHERE DISCHARGE WAS MEASURED (SORTED BY ANALYTE, SITE AND DATE).	75
TABLE III-10. NUMBER OF CHLORPYRIFOS APPLICATIONS, TOTAL POUNDS APPLIED AND TOTAL ACRES TREATED BY MONTH FOR JANUARY 2005 THROUGH DECEMBER 2009 IN THE PRAIRIE FLOWER DRAIN @ CROWS LANDING ROAD SITE SUBWATERSHED. IF A MONTH IS NOT INCLUDED IN THE TABLE, NO APPLICATIONS WERE MADE.	76
TABLE III-11. TOTAL POUNDS AI FOR CHLORPYRIFOS BASED ON PUR DATA FROM 2005-2009 WITHIN THE PRAIRIE FLOWER DRAIN @ CROWS LANDING RD SUBWATERSHED.	79
TABLE IV-1. COORDINATES OF THE BEAR CREEK SITE SUBWATERSHED SAMPLING LOCATION.	83
TABLE IV-2. BEAR CREEK @ KIBBY RD SAMPLING EVENTS PER SEASON AND YEAR.	85

TABLE IV-3. NUMBER OF ANALYSES PERFORMED PER ANALYTE IN EACH SAMPLING SEASON AND YEAR FOR THE BEAR CREEK @ KIBBY RD SAMPLE SITE. ONLY ENVIRONMENTAL SAMPLES WITH A SAMPLE REPLICATE AND LAB REPLICATE NUMBER OF ONE ARE SHOWN. 86

TABLE IV-4. BEAR CREEK @ KIBBY RD SITE SUBWATERSHED. 2008 MANAGEMENT PLAN ADDITIONAL (A) SAMPLING SCHEDULE FOR CHLORPYRIFOS AND *CERIODAPHNIA*. "X" INDICATES THE SITE, MONTH, AND ANALYTE SAMPLED. 89

TABLE IV-5. ALL EXCEEDANCES EXPERIENCED IN SAMPLES COLLECTED FROM LOCATIONS WITHIN THE BEAR CREEK SITE SUBWATERSHED BETWEEN MARCH 2005 AND OCTOBER 2008 (SORTED BY SEASON AND DATE). IF THE WATER QUALITY TRIGGER LIMIT IS BASED ON HARDNESS THEN THE HARDNESS VALUE IS SHOWN IN PARENTHESIS; OTHERWISE THE WQTL USED TO EVALUATE THE DATA IS LISTED IN THE HEADER AFTER THE ANALYTE. 91

TABLE IV-6. BEAR CREEK @ KIBBY RD SITE SUBWATERSHED. NORMAL MONITORING (NM) AND MANAGEMENT PLAN MONITORING (MPM) RESULTS FOR CHLORPYRIFOS AND *CERIODAPHNIA* FROM THE 2008 IRRIGATION SEASON. EXCEEDANCE VALUES ARE IN BOLD. 93

TABLE IV-7. BEAR CREEK SITE SUBWATERSHED. INSTANTANEOUS LOAD CALCULATIONS FOR CHLORPYRIFOS AND COPPER WHERE DISCHARGE WAS MEASURED (SORTED BY ANALYTE, SITE AND DATE). 94

TABLE IV-8. NUMBER OF CHLORPYRIFOS APPLICATIONS, TOTAL POUNDS APPLIED AND TOTAL ACRES TREATED BY MONTH FOR JANUARY 2005 THROUGH DECEMBER 2009 IN THE BEAR CREEK @ KIBBY RD SITE SUBWATERSHED. IF A MONTH IS NOT INCLUDED IN THE TABLE, NO APPLICATIONS WERE MADE. 95

TABLE IV-9. TOTAL POUNDS AI FOR CHLORPYRIFOS BASED ON PUR DATA FROM 2005-2009 WITHIN THE BEAR CREEK @ KIBBY RD SUBWATERSHED. 97

TABLE IV-10. NUMBER OF COPPER APPLICATIONS, TOTAL POUNDS APPLIED AND TOTAL ACRES TREATED BY MONTH FOR JANUARY 2005 THROUGH DECEMBER 2009 IN THE BEAR CREEK @ KIBBY RD SITE SUBWATERSHED. IF A MONTH IS NOT INCLUDED IN THE TABLE, NO APPLICATIONS WERE MADE. 99

TABLE IV-11. TOTAL POUNDS AI FOR CHOPPER BASED ON PUR DATA FROM 2005-2009 WITHIN THE BEAR CREEK @ KIBBY RD SUBWATERSHED. 101

TABLE V-1. COORDINATES OF THE COTTONWOOD CREEK SITE SUBWATERSHED SAMPLING LOCATIONS. 104

TABLE V-2. COTTONWOOD CREEK @ Rd 20 SAMPLING EVENTS PER SEASON AND YEAR 106

TABLE V-3. NUMBER OF ANALYSES PERFORMED PER ANALYTE IN EACH SAMPLING SEASON AND YEAR FOR THE COTTONWOOD CREEK @ Rd 20 SAMPLE SITE. ONLY ENVIRONMENTAL SAMPLES WITH A SAMPLE REPLICATE AND LAB REPLICATE NUMBER OF ONE ARE SHOWN. 107

TABLE V-4. COTTONWOOD CREEK SITE SUBWATERSHED. 2007 MANAGEMENT PLAN ADDITIONAL (A) SAMPLING SCHEDULE FOR COPPER. "X" INDICATES THE SITE, MONTH, AND ANALYTE SAMPLED. 110

TABLE V-5. COTTONWOOD CREEK SITE SUBWATERSHED. 2008 MANAGEMENT PLAN UPSTREAM (U) SAMPLING SCHEDULE FOR COPPER. "X" INDICATES THE SITE, MONTH AND ANALYTE SAMPLED. 110

TABLE V-6. ALL EXCEEDANCES EXPERIENCED IN SAMPLES COLLECTED FROM LOCATIONS WITHIN THE COTTONWOOD CREEK SITE SUBWATERSHED BETWEEN FEBRUARY 2005 AND DECEMBER 2009 (SORTED BY SEASON AND DATE). IF THE WATER QUALITY TRIGGER LIMIT

IS BASED ON HARDNESS THEN THE HARDNESS VALUE IS SHOWN IN PARENTHESIS; OTHERWISE THE WQTL USED TO EVALUATE THE DATA IS LISTED IN THE HEADER AFTER THE ANALYTE.....112

TABLE V-7. COTTONWOOD CREEK SITE SUBWATERSHED. NORMAL MONITORING (NM) AND MANAGEMENT PLAN MONITORING (MPM) RESULTS WHERE 'A' INDICATES ADDITIONAL MPM (2007) AND 'US' INDICATES UPSTREAM MPM (2008) FOR COPPER. EXCEEDANCE VALUES ARE IN BOLD.115

TABLE V-8. COTTONWOOD CREEK SITE SUBWATERSHED. INSTANTANEOUS LOAD CALCULATIONS FOR CHLORPYRIFOS, COPPER, DIAZINON AND DIURON WHERE DISCHARGE WAS MEASURED (SORTED BY ANALYTE, SITE AND DATE).116

TABLE V-9. NUMBER OF CHLORPYRIFOS APPLICATIONS, TOTAL POUNDS APPLIED AND TOTAL ACRES TREATED BY MONTH FOR JANUARY 2005 THROUGH DECEMBER 2009 IN THE COTTONWOOD CREEK @ Rd 20 SITE SUBWATERSHED. IF A MONTH IS NOT INCLUDED IN THE TABLE, NO APPLICATIONS WERE MADE.118

TABLE V-10. TOTAL POUND AI FOR CHLORPYRIFOS BASED ON PUR DATA FROM 2005-2009 WITHIN THE COTTONWOOD CREEK @ Rd 20 SUBWATERSHED.120

TABLE V-11. NUMBER OF DIAZINON APPLICATIONS, TOTAL POUNDS APPLIED AND TOTAL ACRES TREATED BY MONTH FOR JANUARY 2005 THROUGH DECEMBER 2009 IN THE COTTONWOOD CREEK @ Rd 20 SITE SUBWATERSHED. IF A MONTH IS NOT INCLUDED IN THE TABLE, NO APPLICATIONS WERE MADE.122

TABLE V-12. TOTAL POUND AI FOR DIAZINON BASED ON PUR DATA FROM 2005-2009 WITHIN THE COTTONWOOD CREEK @ Rd 20 SUBWATERSHED.125

TABLE V-13. NUMBER OF COPPER APPLICATIONS, TOTAL POUNDS APPLIED AND TOTAL ACRES TREATED BY MONTH FOR JANUARY 2005 THROUGH DECEMBER 2009 IN THE COTTONWOOD CREEK @ Rd 20 SITE SUBWATERSHED. IF A MONTH IS NOT INCLUDED IN THE TABLE, NO APPLICATIONS WERE MADE.127

TABLE V-14. TOTAL POUND AI FOR COPPER BASED ON PUR DATA FROM 2005-2009 WITHIN THE COTTONWOOD CREEK @ Rd 20 SUBWATERSHED.131

TABLE V-15. NUMBER OF DIURON APPLICATIONS, TOTAL POUNDS APPLIED AND TOTAL ACRES TREATED BY MONTH FOR JANUARY 2005 THROUGH DECEMBER 2009 IN THE COTTONWOOD CREEK @ Rd 20 SITE SUBWATERSHED. IF A MONTH IS NOT INCLUDED IN THE TABLE, NO APPLICATIONS WERE MADE.135

TABLE V-16. TOTAL POUNDS AI FOR DIURON BASED ON PUR DATA FROM 2005-2009 WITHIN THE COTTONWOOD CREEK @ Rd 20 SUBWATERSHED.138

TABLE VI-1. COORDINATES OF THE DUCK SLOUGH SITE SUBWATERSHED SAMPLING LOCATIONS.141

TABLE VI-2. DUCK SLOUGH @ GURR RD SAMPLING EVENTS PER SEASON AND YEAR.143

TABLE VI-3. NUMBER OF ANALYSES PERFORMED PER ANALYTE IN EACH SAMPLING SEASON AND YEAR FOR THE DUCK SLOUGH @ GURR RD SAMPLE SITE. ONLY ENVIRONMENTAL SAMPLES WITH A SAMPLE REPLICATE AND LAB REPLICATE NUMBER OF ONE ARE SHOWN.144

TABLE VI-4. DUCK SLOUGH @ GURR RD. 2007 MANAGEMENT PLAN ADDITIONAL (A) SAMPLING SCHEDULE FOR CHLORPYRIFOS, THIOBENCARB AND COPPER. "X" INDICATES THE SITE, MONTH, AND ANALYTE SAMPLED.	147
TABLE VI-5. DUCK SLOUGH SITE SUBWATERSHED. 2008 MANAGEMENT PLAN UPSTREAM (U) SAMPLING SCHEDULE FOR CHLORPYRIFOS AND COPPER. "X" INDICATES THE SITE, MONTH AND ANALYTE SAMPLED.	147
TABLE VI-6. ALL EXCEEDANCES EXPERIENCED IN SAMPLES COLLECTED FROM LOCATIONS WITHIN THE DUCK SLOUGH SITE SUBWATERSHED BETWEEN FEBRUARY 2005 AND DECEMBER 2009 (SORTED BY SEASON AND DATE). IF THE WATER QUALITY TRIGGER LIMIT IS BASED ON HARDNESS THEN THE HARDNESS VALUE IS SHOWN IN PARENTHESIS; OTHERWISE THE WQTL USED TO EVALUATE THE DATA IS LISTED IN THE HEADER AFTER THE ANALYTE.	149
TABLE VI-7. DUCK SLOUGH @ GURR RD. NORMAL MONITORING (NM) AND MANAGEMENT PLAN MONITORING (MPM) RESULTS WHERE 'A' INDICATES ADDITIONAL MPM (2007) AND 'US' INDICATES UPSTREAM MONITORING FOR CHLORPYRIFOS, COPPER AND THIOBENCARB FROM THE 2007 IRRIGATION SEASON. NM AND MPM (2008) FOR COPPER AND <i>SELENASTRUM</i> . NM FOR COPPER (2009). EXCEEDANCE VALUES ARE IN BOLD.....	152
TABLE VI-8. DUCK SLOUGH SITE SUBWATERSHED. INSTANTANEOUS LOAD CALCULATIONS FOR CHLORPYRIFOS AND COPPER WHERE DISCHARGE WAS MEASURED (SORTED BY ANALYTE, SITE AND DATE).	153
TABLE VI-9. NUMBER OF CHLORPYRIFOS APPLICATIONS, TOTAL POUNDS APPLIED AND TOTAL ACRES TREATED BY MONTH FOR JANUARY 2004 THROUGH DECEMBER 2009 IN THE DUCK SLOUGH @ GURR ROAD SITE SUBWATERSHED. IF A MONTH IS NOT INCLUDED IN THE TABLE, NO APPLICATIONS WERE MADE.	155
TABLE VI-10. TOTAL POUND AI FOR CHLORPYRIFOS BASED ON PUR DATA FROM 2004-2009 WITHIN THE DUCK SLOUGH @ GURR RD SUBWATERSHED.	158
TABLE VI-11. NUMBER OF COPPER APPLICATIONS, TOTAL POUNDS APPLIED AND TOTAL ACRES TREATED BY MONTH FOR JANUARY 2004 THROUGH DECEMBER 2009 IN THE DUCK SLOUGH @ GURR RD SITE SUBWATERSHED. IF A MONTH IS NOT INCLUDED IN THE TABLE, NO APPLICATIONS WERE MADE.	161
TABLE VI-12. TOTAL POUND AI FOR COPPER BASED ON PUR DATA FROM 2004-2009 WITHIN THE DUCK SLOUGH @ GURR RD SUBWATERSHED.	165
TABLE VII-1. COORDINATES OF THE HIGHLINE CANAL SITE SUBWATERSHED SAMPLING LOCATION.....	169
TABLE VII-2. HIGHLINE CANAL @ HWY 99 SAMPLING EVENTS PER SEASON AND YEAR.	171
TABLE VII-3. NUMBER OF ANALYSES PERFORMED PER ANALYTE IN EACH SAMPLING SEASON AND YEAR FOR THE HIGHLINE CANAL @ HWY 99 SAMPLE SITE. ONLY ENVIRONMENTAL SAMPLES WITH A SAMPLE REPLICATE AND LAB REPLICATE NUMBER OF ONE ARE SHOWN.	172
TABLE VII-4. HIGHLINE CANAL @ HWY 99. 2007 MANAGEMENT PLAN ADDITIONAL (A) SAMPLING SCHEDULE FOR <i>CERIODAPHNIA</i> . "X" INDICATES THE SITE, MONTH, AND ANALYTE SAMPLED.	175
TABLE VII-5. HIGHLINE CANAL @ HWY 99. 2008 MANAGEMENT PLAN ADDITIONAL (A) SAMPLING SCHEDULE FOR CHLORPYRIFOS, COPPER, AND <i>CERIODAPHNIA</i> . "X" INDICATES THE SITE, MONTH, AND ANALYTE SAMPLED.....	175

TABLE VII-6. HIGHLINE CANAL @ HWY 99. 2009 MANAGEMENT PLAN SAMPLING SCHEDULE FOR CHLORPYRIFOS AND *SELENASTRUM*. "X" INDICATES THE SITE, MONTH AND ANALYTE SAMPLED.175

TABLE VII-7. ALL EXCEEDANCES EXPERIENCED IN SAMPLES COLLECTED WITHIN THE HIGHLINE CANAL @ HWY 99 SITE SUBWATERSHED BETWEEN MARCH 2005 AND DECEMBER 2009 (SORTED BY SEASON AND DATE). IF THE WATER QUALITY TRIGGER LIMIT IS BASED ON HARDNESS THEN THE HARDNESS VALUE IS SHOWN IN PARENTHESIS; OTHERWISE THE WQTL USED TO EVALUATE THE DATA IS LISTED IN THE HEADER AFTER THE ANALYTE.177

TABLE VII-8. HIGHLINE CANAL @ HWY 99. NORMAL MONITORING (NM) AND MANAGEMENT PLAN MONITORING (MPM) RESULTS FOR *CERIODAPHNIA* FROM THE 2007 IRRIGATION SEASON. NORMAL MONITORING (NM) AND MANAGEMENT PLAN MONITORING (MPM) RESULTS FOR COPPER, CHLORPYRIFOS AND *CERIODAPHNIA* FOR THE 2008 IRRIGATION SEASON AND MANAGEMENT PLAN MONITORING (MPM) RESULTS FOR CHLORPYRIFOS AND *SELENASTRUM* FOR 2009. EXCEEDANCE VALUES ARE IN BOLD.180

TABLE VII-9. HIGHLINE CANAL @ HWY 99. INSTANTANEOUS LOAD CALCULATIONS FOR CHLORPYRIFOS, COPPER AND DIURON WHERE DISCHARGE WAS MEASURED (SORTED BY ANALYTE AND DATE).....180

TABLE VII-10. NUMBER OF CHLORPYRIFOS APPLICATIONS, TOTAL POUNDS APPLIED AND TOTAL ACRES TREATED BY MONTH FOR JANUARY 2005 THROUGH DECEMBER 2009 IN THE HIGHLINE CANAL @ HWY 99 SITE SUBWATERSHED. IF A MONTH IS NOT INCLUDED IN THE TABLE, NO APPLICATIONS WERE MADE.182

TABLE VII-11. TOTAL POUNDS AI FOR CHLORPYRIFOS BASED ON PUR DATA FROM 2005-2009 WITHIN THE HIGHLINE CANAL @ HWY 99 SUBWATERSHED.185

TABLE VII-12. NUMBER OF COPPER APPLICATIONS, TOTAL POUNDS APPLIED AND TOTAL ACRES TREATED BY MONTH FOR JANUARY 2005 THROUGH DECEMBER 2009 IN THE HIGHLINE CANAL @ HWY 99 SITE SUBWATERSHED. IF A MONTH IS NOT INCLUDED IN THE TABLE, NO APPLICATIONS WERE MADE.187

TABLE VII-13. TOTAL POUNDS AI FOR COPPER BASED ON PUR DATA FROM 2005-2009 WITHIN THE HIGHLINE CANAL @ HWY 99 SUBWATERSHED.190

TABLE VII-14. NUMBER OF DIURON APPLICATIONS, TOTAL POUNDS APPLIED AND TOTAL ACRES TREATED BY MONTH FOR JANUARY 2005 THROUGH DECEMBER 2009 IN THE HIGHLINE CANAL @ HWY 99 SITE SUBWATERSHED. IF A MONTH IS NOT INCLUDED IN THE TABLE, NO APPLICATIONS WERE MADE.192

TABLE VII-15. TOTAL POUNDS AI FOR DIURON BASED ON PUR DATA FROM 2005-2009 WITHIN THE HIGHLINE CANAL @ HWY 99 SUBWATERSHED.195

List of Figures

FIGURE I-1. SITE SUBWATERSHED MAP OF LAND USE FOR THE DRY CREEK @ WELLSFORD RD SAMPLE SITE.	2
FIGURE I-2. POUNDS OF CHLORPYRIFOS APPLIED WITHIN THE DRY CREEK @ WELLSFORD RD SITE SUBWATERSHED BY MONTH FOR 2005-2009. ASTERISK (*) DENOTES MONTHS WITH EXCEEDANCES.	19
FIGURE I-3. POUNDS OF COPPER APPLIED WITHIN THE DRY CREEK @ WELLSFORD RD SITE SUBWATERSHED BY MONTH FOR 2005-2009. ASTERISK (*) DENOTES MONTHS WITH EXCEEDANCES.	25
FIGURE I-4. POUNDS OF DIURON APPLIED WITHIN THE DRY CREEK @ WELLSFORD RD SITE SUBWATERSHED BY MONTH FOR 2005-2009. ASTERISK (*) DENOTES MONTHS WITH EXCEEDANCES.	31
FIGURE II-1. SITE SUBWATERSHED MAP OF LAND USE FOR THE DUCK SLOUGH @ HWY 99 SAMPLE SITE.	36
FIGURE II-2. POUNDS OF CHLORPYRIFOS APPLIED WITHIN THE DUCK SLOUGH @ HWY 99 SITE SUBWATERSHED BY MONTH FOR 2005-2009. ASTERISK (*) DENOTES MONTHS WITH EXCEEDANCES.	51
FIGURE II-3. POUNDS OF COPPER APPLIED WITHIN THE DUCK SLOUGH @ HWY 99 SITE SUBWATERSHED BY MONTH FOR 2005-2009. ASTERISK (*) DENOTES MONTHS WITH EXCEEDANCES.	56
FIGURE III-1. SITE SUBWATERSHED MAP OF LAND USE FOR THE PRAIRIE FLOWER DRAIN @ CROWS LANDING RD SAMPLE SITE.	62
FIGURE III-2. POUNDS OF CHLORPYRIFOS ADDED WITHIN THE PRAIRIE FLOWER DRAIN @ CROWS LANDING RD SITE SUBWATERSHED BY MONTH FOR 2005-2009. ASTERISK (*) DENOTES MONTHS WITH EXCEEDANCES. NO USE OF CHLORPYRIFOS WAS REPORTED FOR 2007.	78
FIGURE IV-1. SITE SUBWATERSHED MAP OF LAND USE FOR THE BEAR CREEK @ KIBBY RD SAMPLE SITE.	84
FIGURE IV-2. POUNDS OF CHLORPYRIFOS APPLIED WITHIN THE BEAR CREEK SITE SUBWATERSHED BY MONTH FOR 2005-2009. ASTERISK (*) DENOTES MONTHS WITH EXCEEDANCES.	96
FIGURE IV-3. POUNDS OF COPPER APPLIED WITHIN THE BEAR CREEK @ KIBBY RD SITE SUBWATERSHED BY MONTH FOR 2005-2009. ASTERISK (*) DENOTES MONTHS WITH EXCEEDANCES.	100
FIGURE V-1. SITE SUBWATERSHED MAP OF LAND USE FOR THE COTTONWOOD CREEK @ RD 20 SAMPLE SITE.	105
FIGURE V-2. POUNDS OF CHLORPYRIFOS APPLIED WITHIN THE COTTONWOOD CREEK @ RD 20 SITE SUBWATERSHED BY MONTH FOR 2005-2009. ASTERISK (*) DENOTES MONTHS WITH EXCEEDANCES.	119
FIGURE V-3. POUNDS OF DIAZINON APPLIED WITHIN THE COTTONWOOD CREEK @ RD 20 SITE SUBWATERSHED BY MONTH FOR 2005-2009. ASTERISK (*) DENOTES MONTHS WITH EXCEEDANCES.	124

FIGURE V-4. POUNDS OF COPPER APPLIED WITHIN THE COTTONWOOD CREEK @ RD 20 SITE SUBWATERSHED BY MONTH FOR 2005-2009. ASTERISK (*) DENOTES MONTHS WITH EXCEEDANCES..... 130

FIGURE V-5. POUNDS OF DIURON APPLIED WITHIN THE COTTONWOOD CREEK @ RD 20 SITE SUBWATERSHED BY MONTH FOR 2005-2009. ASTERISK (*) DENOTES MONTHS WITH EXCEEDANCES. 137

FIGURE VI-1. SITE SUBWATERSHED MAP OF LAND USE FOR THE DUCK SLOUGH @ GURR RD SAMPLE SITE..... 142

FIGURE VI-2. POUNDS OF CHLORPYRIFOS APPLIED WITHIN THE DUCK SLOUGH @ GURR RD SITE SUBWATERSHED BY MONTH FOR 2004-2009. ASTERISK (*) DENOTES MONTHS WITH EXCEEDANCES. 157

FIGURE VI-3. POUNDS OF COPPER APPLIED WITHIN THE DUCK SLOUGH @ GURR RD SITE SUBWATERSHED BY MONTH FOR 2004-2009. ASTERISK (*) DENOTES MONTHS WITH EXCEEDANCES..... 164

FIGURE VII-1. SITE SUBWATERSHED MAP OF LAND USE FOR THE HIGHLINE CANAL @ HWY 99 SAMPLE SITE..... 170

FIGURE VII-2. POUNDS OF CHLORPYRIFOS APPLIED WITHIN THE HIGHLINE CANAL @ HWY 99 SITE SUBWATERSHED BY MONTH FOR 2005-2009. ASTERISK (*) DENOTES MONTHS WITH EXCEEDANCES. 184

FIGURE VII-3. POUNDS OF COPPER APPLIED WITHIN THE HIGHLINE CANAL @ HWY 99 SITE SUBWATERSHED BY MONTH FOR 2005-2009. ASTERISK (*) DENOTES MONTHS WITH EXCEEDANCES..... 189

FIGURE VII-4. POUNDS OF DIURON APPLIED WITHIN THE HIGHLINE CANAL @ HWY 99 SITE SUBWATERSHED BY MONTH FOR 2005-2009. ASTERISK (*) DENOTES MONTHS WITH EXCEEDANCES. NO USE OF DIURON WAS REPORTED FOR 2008. 194

High Priority Site Subwatersheds (2008 -2010)

I. DRY CREEK @ WELLSFORD RD

Management Plan Constituents

Priority A/B

- Chlorpyrifos

Priority C

- Copper
- Diuron

Priority D

- *Selenastrum capricornutum* water column toxicity
- *Hyalella azteca* sediment toxicity

Priority E

- Dissolved Oxygen
- *E. coli*
- pH
- Specific Conductance

Description of Dry Creek at Wellsford Rd Site Subwatershed

Dry Creek @ Wellsford Road (23,331 irrigated acres) – This site subwatershed is in the northern part of the Coalition region and drains a combination of field crops, deciduous nuts, and vineyards (Figure I-1). Dry Creek originates to the east of Modesto and drains into the Tuolumne River. This site subwatershed samples Dry Creek at the furthest downstream location that collects agricultural drainage prior to flowing through the urban areas of Modesto. Dairies are located upstream of this site and the town of Waterford may contribute an urban signal. This site subwatershed includes an upstream location (Dry Creek @ Waterford Rd) which was sampled in 2008 and 2009. Table I-1 includes the station name, station code and target latitude/longitudes for sites sampled within this subwatershed.


Table I-1. Coordinates of the Dry Creek site subwatershed sampling locations.

Station Name	Station Code	Target Latitude	Target Longitude
Dry Creek @ Waterford ^u Rd	535XDCAWF	37.65876	-120.77887
Dry Creek @ Wellsford Rd*	535XDCAWR	37.6602	-120.8743

*Original ESJWQC sampling site

^uUpstream sites

Figure I-1. Site subwatershed map of land use for the Dry Creek @ Wellsford Rd sample site.



Subwatershed Monitoring History

Monitoring within the Dry Creek site subwatershed at Dry Creek @ Wellsford Rd was initiated in 2005 and has continued uninterrupted through 2009 (Table I-2). Dry Creek @ Wellsford Rd will remain a Core Monitoring location under the current MRPP through 2010. In 2011 this site will be monitored as an Assessment site. The constituents sampled at this location from 2005-2009 are listed in Table I-3. A summary and discussion of exceedances are provided in the next section (Table I-7).

Management Plan Monitoring for the Coalition was initiated during the 2007 irrigation season and has included both additional sampling at Dry Creek @ Wellsford Road and upstream monitoring at Dry Creek @ Waterford Rd for chlorpyrifos and toxicity to *Ceriodaphnia* (Table I-4, Table I-5, Table I-6).

Dry Creek is considered an impaired water body and is included on the proposed 2008 Central Valley Basin Plan 303d list for chlorpyrifos, diazinon, *E. coli*, and unknown toxicity. Agriculture is listed as the potential source for the impairment caused by chlorpyrifos and diazinon.

Table I-2. Dry Creek @ Wellsford Rd sampling events per season and year.

	2004	2005		2006		2007		2008			2009			
	Irrigation	Storm	Irrigation	Storm	Irrigation	Storm	Irrigation	Storm	Irrigation	Fall	Winter	Storm	Irrigation	Fall
Events Sampled	NA	2	5	2	5	2	6	3	6	2	2	2	6	3
Events Not Sampled	NA	0	0	0	0	0	0	0	0	0	0	0	0	0
Total	NA	2	5	2	5	2	6	3	6	2	2	0	6	3

NA - Not applicable. This site was not sampled during this season/year.

Table I-3. Number of analyses performed per analyte in each sampling season and year for the Dry Creek @ Wellsford Rd sample site. Only environmental samples with a sample replicate and lab replicate number of one are shown.

Method	Analyte	2005		2006		2007		2008			2009			
		Storm	Irrigation	Storm	Irrigation	Storm	Irrigation	Storm	Irrigation	Fall	Storm	Winter	Irrigation	Fall
Field and Physical Parameters														
EPA 110.2	Color	2	5	2	5	2	6	2	6					
EPA 160.1	Dissolved Solids	2	5	2	5	2	6	3	6	2	2	2	6	2
EPA 160.2	Suspended Solids							1		2	2	2	6	2
EPA 180.1	Turbidity	2	5	2	5	2	6	3	6	2	2	2	6	2
EPA 405.1	BOD				1	2	2							
EPA 415.1	Total Organic Carbon	2	5	2	5	2	6	3	6	2	2	2	6	2
SM 9223	E. coli	2	5	2	5	2	6	3	6	2	2	2	6	2
NA	Dissolved Oxygen	2	5	3	5	3	10	7	8	2	2	2	6	2
NA	Specific Conductivity	2	5	3	5	3	10	7	8	2	2	2	6	2
NA	pH	2	5	3	5	3	10	7	8	2	2	2	6	2
Carbamates														
EPA 8321A	Aldicarb				5	2	6	2	6					
EPA 8321A	Carbaryl				5	2	6	2	6					
EPA 8321A	Carbofuran				5	2	6	2	6					
EPA 8321A	Diuron				5	2	6	2	6					
EPA 8321A	Linuron				5	2	6	2	6					
EPA 8321A	Methiocarb				5	2	6	2	6					
EPA 8321A	Methomyl				5	2	6	2	6					
EPA 8321A	Oxamyl				5	2	6	2	6					
Organochlorines														
EPA 8081A	DDD(p,p')				5	2	6	2	6					
EPA 8081A	DDE(p,p')				5	2	6	2	6					
EPA 8081A	DDT(p,p')				5	2	6	2	6					
EPA 8081A	Dicofol				5	2	6	2	6					

Method	Analyte	2005		2006		2007		2008			2009			
		Storm	Irrigation	Storm	Irrigation	Storm	Irrigation	Storm	Irrigation	Fall	Storm	Winter	Irrigation	Fall
EPA 8081A	Dieldrin				5	2	6	2	6					
EPA 8081A	Endrin				5	2	6	2	6					
EPA 8081A	Methoxychlor				5	2	6	2	6					
Organophosphates														
EPA 8141A	Azinphos methyl				5	2	7	2	6					
EPA 8141A	Chlorpyrifos	2	5	2	5	2	8	2	6				2	
EPA 8141A	Diazinon	2	5	2	5	2	7	2	6					
EPA 8141A	Dimethoate	2	5	2	5	2	7	2	6					
EPA 8141A	Disulfoton				5	2	7	2	6					
EPA 8141A	Malathion				5	2	7	2	6					
EPA 8141A	Methamidophos				5	2	6	2	6					
EPA 8141A	Methidathion				5	2	7	2	6					
EPA 8141A	Molinate				5	2	7	2	6					
EPA 8141A	Parathion, Methyl				5	2	7	2	6					
EPA 8141A	Phorate				5	2	7	2	6					
EPA 8141A	Phosmet				5	2	7	2	6					
EPA 8141A	Thiobencarb				5	2	7	2	6					
Pyrethroids														
EPA 8081A	Bifenthrin		1	2	5	2	6							
EPA 8081A	Cyfluthrin, total		1	2	5	2	6							
EPA 8081A	Cyhalothrin, lambda, total	2	5	2	5	2	6							
EPA 8081A	Cypermethrin, total	2	5	2	5	2	6							
EPA 8081A	Esfenvalerate/ Fenvalerate, total	2	5	2	5	2	6							
EPA 8081A	Permethrin, total	2	5	2	5	2	6							
Triazines														
EPA 547M	Glyphosate				5	2	6	2	6					
EPA 549.2M	Paraquat dichloride				5	2	6	2	6					

Method	Analyte	2005		2006		2007		2008			2009			
		Storm	Irrigation	Storm	Irrigation	Storm	Irrigation	Storm	Irrigation	Fall	Storm	Winter	Irrigation	Fall
EPA 619	Atrazine				5	2	6	2	6					
EPA 619	Cyanazine				5	2	6	2	6					
EPA 619	Simazine				5	2	6	2	6					
Metals (Total)														
EPA 200.8	Arsenic				5	2	6	2	6					
EPA 200.8	Boron				5	2	6	2	6					
EPA 200.8	Cadmium				5	2	6	2	6					
EPA 200.8	Copper				5	2	6	2	6					
EPA 200.8	Lead				5	2	6	2	6		1			
EPA 200.8	Nickel				5	2	6	2	6					
EPA 200.8	Selenium				5	2	2	2	6					
EPA 200.8	Zinc				5	2	6	2	6					
Metals (Dissolved)														
EPA 200.8	Cadmium													
EPA 200.8	Copper													
EPA 200.8	Lead										1			
EPA 200.8	Nickel													
EPA 200.8	Zinc													
Nutrients														
SM 2340 C	Hardness as CaCO3				5	2	6	2	6		1			
EPA 300.0	Nitrate as N				5	2	6	2	6					
EPA 350.2	Ammonia as N				5	2	6	3	6	2	2	2	6	2
EPA 351.3	Nitrogen, Total Kjeldahl				5	2	6	3	6	2	1	2	1	
EPA 354.1	Nitrite as N				5	2	6	2	6					
EPA 353.2	Nitrate + Nitrite as N							1		2	2	2	6	2
EPA 365.2	Orthophosphate as P				5	2	6	3	6	2	1	3	1	
EPA 365.2	Phosphate as P				5	2	6	3	6	2	2	2	6	2
Toxicity														

Method	Analyte	2005		2006		2007		2008			2009			
		Storm	Irrigation	Storm	Irrigation	Storm	Irrigation	Storm	Irrigation	Fall	Storm	Winter	Irrigation	Fall
EPA 821-02-012	Ceriodaphnia dubia	2	5	2	6	2	7	2	6					
EPA 821-02-012	Pimephales promelas	2	5	2	5	2	6	2	6					
EPA 821-02-013	Selenastrum capricornutum	2	5	2	5	2	6	4	6					
EPA 600/R-99-064	Hyalella azteca		1	1	1	1	1	2	2					

NA - Not applicable

Table I-4. Dry Creek site subwatershed. 2007 Management Plan additional (A) sampling schedule for chlorpyrifos and *Ceriodaphnia dubia*. "X" indicates the site, month, and analyte sampled.

Sample Site	Date	Type	Chlorpyrifos	<i>Ceriodaphnia dubia</i>
Dry Creek @ Wellsford Rd	31-Jul-07	A	X	
Dry Creek @ Wellsford Rd	28-Aug-07	A	X	
Dry Creek @ Wellsford Rd	25-Sep-07	A		X

Table I-5. Dry Creek site subwatershed. 2008 Management Plan upstream (U) sampling schedule for chlorpyrifos and *Ceriodaphnia dubia*. "X" indicates the site, month, and analyte sampled.

Sample Site	Date	Type	Chlorpyrifos	<i>Ceriodaphnia dubia</i>
Dry Creek @ Waterford Rd	22-Jul-08	U	X	
Dry Creek @ Waterford Rd	19-Aug-08	U	X	
Dry Creek @ Waterford Rd	23-Sep-08	U	X	X

Table I-6. Dry Creek site subwatershed. 2009 Management Plan Monitoring (MPM) and upstream (U) sampling schedule for chlorpyrifos. "X" indicates the site, month and analyte sampled.

Site Name	Date	Type	Chlorpyrifos
Dry Creek @ Waterford Rd	21-Jul-09	U	X
Dry Creek @ Wellsford Rd	21-Jul-09	MPM	X
Dry Creek @ Waterford Rd	18-Aug-09	U	X
Dry Creek @ Wellsford Rd	18-Aug-09	MPM	X

Exceedance History

Ambient water monitoring within the Dry Creek subwatershed was initiated in the 2005 storm sampling season at the site at Wellsford Rd. Dry Creek @ Wellsford Rd samples collected from 2005-2009 have experienced exceedances WQTL for field parameters, *E. coli*, pesticides, and metals (Table I-7). Toxicity in the water column to *Ceriodaphnia dubia*, and *Selenastrum capricornutum*, and sediment toxicity to *Hyalella azteca*, has also occurred (Table I-7).

Exceedances in this subwatershed, including the upstream site, include DO (28), pH (4), SC (1), *E. coli* (29), chlorpyrifos (9), diuron (2), thiobencarb (1), copper (3), and lead (1). Water column toxicity to *Ceriodaphnia dubia* occurred twice, *Selenastrum capricornutum* toxicity occurred five times and sediment toxicity to *Hyalella azteca* occurred twice.

In 2009, exceedances included DO (7), SC (1), *E. coli* (8), and chlorpyrifos (1).

All exceedances are listed in Table I-7 by season and date and are based on WQTLs listed in the introduction of the ESJWQC Management Plan. The priority level (A-E) assigned to each constituent is listed in the bottom row of Table I-7 and is determined using the ESJWQC Management Plan prioritization process flow chart (Figure 4). Specific conductance is now under a management plan following the exceedance during the 2009 sampling season. Additionally, *Ceriodaphnia dubia* water column toxicity is no longer prioritized because the only two toxic samples were collected over three years ago.

Table I-7. All exceedances experienced in samples collected from locations within the Dry Creek site subwatershed from February 2005 through December 2009 (sorted by season and date). If the water quality trigger limit is based on hardness then the hardness value is shown in parenthesis; otherwise the WQTL used to evaluate the data is listed in the header after the analyte. Upstream site is italicized.

Station Name	Season	Sample Date	Oxygen, Dissolved, 7 mg/L	pH, 6.5-8.5 units	Specific Conductivity, 700 µS/cm	<i>E. coli</i> , 235 MPN/100 mL	Copper, Total, 1,300 µg/L	Lead, Total, 15 µg/L	Chlorpyrifos, 0.015 µg/L	Diuron, 2 µg/L	Thiobencarb, 0 µg/L	<i>C. dubia</i> , Survival (%)	<i>H. azteca</i> , Survival (%)	<i>S. capricornutum</i> , Total Cell Count
Dry Creek @ Wellsford Rd	Storm	2/15/2005										80		
Dry Creek @ Wellsford Rd	Storm	3/22/2005		8.96		900								
Dry Creek @ Wellsford Rd	Irrigation	5/11/2005		6.26										
Dry Creek @ Wellsford Rd	Irrigation	6/15/2005	5.9			240								
Dry Creek @ Wellsford Rd	Irrigation	7/13/2005	5.7											
Dry Creek @ Wellsford Rd	Irrigation	8/17/2005		9.18		900			0.024					
Dry Creek @ Wellsford Rd	Irrigation	9/21/2005	6.98			500								
Dry Creek @ Wellsford Rd	Storm	3/1/2006				300								
Dry Creek @ Wellsford Rd	Storm	3/16/2006				1600								
Dry Creek @ Wellsford Rd	Irrigation	5/18/2006				280								
Dry Creek @ Wellsford Rd	Irrigation	6/15/2006	6.08											
Dry Creek @ Wellsford Rd	Irrigation	7/13/2006	6.69						0.026					
Dry Creek @ Wellsford Rd	Irrigation	8/10/2006							0.024					
Dry Creek @ Wellsford Rd	Irrigation	9/14/2006				310				0.1	70			
Dry Creek @ Wellsford Rd	Storm	2/11/2007	6.17			290				37				508000
Dry Creek @ Wellsford Rd	Storm	2/22/2007												220000
Dry Creek @ Wellsford Rd	Storm	2/28/2007				2400	8.4 (7.2)			4				298750
Dry Creek @ Wellsford Rd	Storm	3/7/2007												826000
Dry Creek @ Wellsford Rd	Irrigation	4/17/2007					5.1 (5.0)							

Station Name	Season	Sample Date	Oxygen, Dissolved, 7 mg/L	pH, 6.5-8.5 units	Specific Conductivity, 700 µS/cm	<i>E. coli</i> , 235 MPN/100 mL	Copper, Total, 1,300 µg/L	Lead, Total, 15 µg/L	Chlorpyrifos, 0.015 µg/L	Diuron, 2 µg/L	Thiobencarb, 0 µg/L	<i>C. dubia</i> , Survival (%)	<i>H. azteca</i> , Survival (%)	<i>S. capricornutum</i> , Total Cell Count
Dry Creek @ Wellsford Rd	Irrigation	6/19/2007	5.77											
Dry Creek @ Wellsford Rd	Irrigation	7/17/2007	6.64						0.021					
Dry Creek @ Wellsford Rd	Irrigation	7/31/2007	6.91											
Dry Creek @ Wellsford Rd	Irrigation	8/14/2007	6.58			440								
Dry Creek @ Wellsford Rd	Irrigation	9/11/2007	6.5			420			0.043					
Dry Creek @ Wellsford Rd	Storm	1/24/2008				>2400								
Dry Creek @ Wellsford Rd	Storm	2/25/2008				>2400	11 (6.0)	1.8 (1.7)						494753
Dry Creek @ Wellsford Rd	Sediment	3/4/2008											88	
Dry Creek @ Wellsford Rd	Irrigation	4/22/2008				>2400								
Dry Creek @ Wellsford Rd	Irrigation	5/20/2008	5.67			330								
Dry Creek @ Wellsford Rd	Irrigation	6/17/2008	6.31			>2400								
<i>Dry Creek @ Waterford Rd</i>	<i>Irrigation</i>	<i>7/22/2008</i>	<i>6.08</i>						<i>0.02</i>					
Dry Creek @ Wellsford Rd	Irrigation	7/22/2008	6.67			>2400			0.03					
<i>Dry Creek @ Waterford Rd</i>	<i>Irrigation</i>	<i>8/19/2008</i>	<i>5.93</i>						<i>0.023</i>					
Dry Creek @ Wellsford Rd	Irrigation	8/19/2008	6.85			580								
Dry Creek @ Wellsford Rd	Sediment	8/28/2008	6.64										71	
Dry Creek @ Wellsford Rd	Irrigation	9/23/2008				290								
Dry Creek @ Wellsford Rd	Irrigation	10/2/2008	5.83											
Dry Creek @ Wellsford Rd	Fall	10/21/2008	4.91			550								
Dry Creek @ Wellsford Rd	Fall	12/16/2008	2.77	8.68										
Dry Creek @ Wellsford Rd	Winter	1/20/2009	5.1		707									

Station Name	Season	Sample Date	Oxygen, Dissolved, 7 mg/L	pH, 6.5-8.5 units	Specific Conductivity, 700 µS/cm	<i>E. coli</i> , 235 MPN/100 mL	Copper, Total, 1,300 µg/L	Lead, Total, 15 µg/L	Chlorpyrifos, 0.015 µg/L	Diuron, 2 µg/L	Thiobencarb, 0 µg/L	<i>C. dubia</i> , Survival (%)	<i>H. azteca</i> , Survival (%)	<i>S. capricornutum</i> , Total Cell Count
Dry Creek @ Wellsford Rd	Winter	3/17/2009				250								
Dry Creek @ Wellsford Rd	Irrigation	5/19/2009	6.24			260								
Dry Creek @ Wellsford Rd	Irrigation	6/16/2009				1600								
<i>Dry Creek @ Waterford Rd</i>	<i>Irrigation</i>	<i>7/21/2009</i>	<i>6.89</i>											
Dry Creek @ Wellsford Rd	Irrigation	7/21/2009	5.9			270								
Dry Creek @ Wellsford Rd	Irrigation	8/18/2009				410			0.027					
Dry Creek @ Wellsford Rd	Fall	10/20/2009	4.04			490								
Dry Creek @ Wellsford Rd	Fall	11/17/2009	3.04			730								
Dry Creek @ Wellsford Rd	Storm	12/15/2009	6.65			820								
Constituent Priority			E	E	E	E	C	NP	A/B	C	NP¹	NP	D	D

NP- Not prioritized. Fewer than two exceedances for this constituent at this site within three years and currently no TMDL for constituent

NP¹ - Not Prioritized; thiobencarb is used only by rice and therefore all exceedances are turned over to the Rice Coalition.

2007 - 2009 Management Plan Monitoring Results

Management Plan Monitoring results are included in Table I-8.

2007

Dry Creek was monitored monthly for chlorpyrifos and toxicity to *Ceriodaphnia* during 2007. Additional monitoring also occurred for chlorpyrifos in July and August and for toxicity to *Ceriodaphnia* in September. No toxicity to *Ceriodaphnia* occurred in 2007. Chlorpyrifos exceeded the WQTL of 0.015µg/L during normal monitoring sampling in July and September 2007. Additional samples had no detectable levels of chlorpyrifos (<0.003 µg/L, Table I-8).

2008

In 2008, upstream MPM occurred at Dry Creek @ Waterford Road for chlorpyrifos during July, August, and September and for toxicity to *Ceriodaphnia* during September (Table I-8). Normal monitoring occurred at Dry Creek @ Wellsford Rd for these constituents monthly in 2008. Again, no toxicity to *Ceriodaphnia* occurred in 2008 (Table I-8). Chlorpyrifos exceeded the WQTL during normal monitoring in July (0.03 µg/L) and during upstream monitoring in July and August 2008 (0.02 and 0.023 µg/L, respectively, Table I-8).

2009

Although upstream monitoring is typically scheduled for one season only, the Coalition continued to monitor the Dry Creek upstream site (Dry Creek @ Waterford Rd) during the 2009 irrigation season because chlorpyrifos exceedances were inconsistent at the normal monitoring and upstream locations in 2008 (Table I-8, August 19, 2008). During 2009 monitoring there was a chlorpyrifos detection of 0.013 µg/L at Dry Creek @ Wellsford Rd in samples collected in July and an exceedance of 0.027 µg/L of chlorpyrifos in samples collected in August. Management Plan Monitoring samples collected from Dry Creek @ Waterford Rd during July and August 2009 had no detections of chlorpyrifos.

Management Plan Monitoring for copper, chlorpyrifos, and *Hyalella* toxicity are planned for the 2010 irrigation season. Management Plan Monitoring during February and March 2011 will also occur for copper, diuron and toxicity to *Selenastrum* and *Hyalella*.

Table I-8. Dry Creek site subwatershed. Normal monitoring (NM) and Management Plan Monitoring (MPM) results where 'A' indicates additional MPM (2007) and 'US' indicates upstream MPM (2008) for chlorpyrifos and *Ceriodaphnia* from the 2007-2008 irrigation seasons, and NM and MPM for chlorpyrifos for the irrigation seasons of 2009. Exceedance values are in bold.

Month:		April	May	June	July	August	September
2007 NM (@ Wellsford Rd)	Date	4/17/07	5/15/07	6/19/07	7/17/07	8/14/07	9/11/07
	Chlorpyrifos (µg/L)	<0.003	0.011	<0.003	0.021	<0.003	0.043
	<i>C. dubia</i> toxicity (% Control)	100	100	100	100	100	100
2007 MPM A (@ Wellsford Rd)	Date	NA	NA	NA	7/31/07	8/28/07	9/25/07
	Chlorpyrifos (µg/L)	NA	NA	NA	<0.003	<0.003	NA
	<i>C. dubia</i> toxicity (% Control)	NA	NA	NA	NA	NA	100
2008 NM (@ Wellsford Rd)	Date	4/22/08	5/20/08	6/17/08	7/22/08	8/19/08	9/23/08
	Chlorpyrifos (µg/L)	<0.003	<0.003	<0.003	0.03	<0.003	<0.003
	<i>C. dubia</i> toxicity (% Control)	100	100	100	100	100	100
2008 MPM US (@ Waterford Rd)	Date	NA	NA	NA	7/22/08	8/19/08	9/23/08
	Chlorpyrifos (µg/L)	NA	NA	NA	0.02	0.023	<0.003
	<i>C. dubia</i> toxicity (% Control)	NA	NA	NA	NA	NA	100
2009 MPM (@ Wellsford Rd)	Date	NA	NA	NA	7/21/09	8/18/09	NA
	Chlorpyrifos (µg/L)	NA	NA	NA	0.013	0.027	NA
2009 MPM US (@ Waterford Rd)	Date	NA	NA	NA	7/21/09	8/18/09	NA
	Chlorpyrifos (µg/L)	NA	NA	NA	<0.003	<0.003	NA

NA - Not Applicable; this site was not sampled during this month.

Load Calculations

Loads have been calculated for the chlorpyrifos, copper and diuron detections (Table I-9) based on the following formula:

$$\text{Load} = \text{Discharge (cfs)} \times 28.317\text{L} \times \text{Concentration (milligram/L} \times 1000 \text{ or } \mu\text{g/L)}.$$

The load values calculated and presented for pesticides or other constituents in this report represent instantaneous loads only. These values should not be used to extrapolate loading over any period of time (e.g. weekly, monthly, seasonal or annual). The primary purpose for reporting instantaneous loads is to provide the Regional Water Board with a context for the concentrations of various constituents at the time samples were collected.

Table I-9. Dry Creek @ Wellsford and Dry Creek @ Waterford. Instantaneous load calculations for chlorpyrifos, copper and diuron where discharge was measured (sorted by analyte, site and date). Upstream site is italicized.

Station Name	Analyte Name	Sample Date	Discharge cfs	Concentration $\mu\text{g/L}$	Loading Rate $\mu\text{g/sec}$
Dry Creek @ Wellsford Rd	Chlorpyrifos	17-Aug-05	0	0.024	0
Dry Creek @ Wellsford Rd	Chlorpyrifos	13-Jul-06	123.91	0.026	91.23
Dry Creek @ Wellsford Rd	Chlorpyrifos	15-May-07	51.13	0.011	15.93
Dry Creek @ Wellsford Rd	Chlorpyrifos	17-Jul-07	68.57	0.021	40.78
Dry Creek @ Wellsford Rd	Chlorpyrifos	11-Sep-07	61.02	0.043	74.30
Dry Creek @ Wellsford Rd	Chlorpyrifos	22-Jul-08	63.22	0.03	53.71
Dry Creek @ Wellsford Rd	Chlorpyrifos	21-Jul-09	71.26	0.013	26.23
Dry Creek @ Wellsford Rd	Chlorpyrifos	18-Aug-09	47.07	0.027	35.99
<i>Dry Creek at Waterford</i>	<i>Chlorpyrifos</i>	<i>19-Aug-08</i>	<i>24.24</i>	<i>0.023</i>	<i>15.79</i>
Dry Creek @ Wellsford Rd	Copper	13-Jul-06	123.91	3.1	10877.15
Dry Creek @ Wellsford Rd	Copper	17-Apr-07	30.71	5.1	4435.04
Dry Creek @ Wellsford Rd	Copper	15-May-07	51.13	6.1	8831.87
Dry Creek @ Wellsford Rd	Copper	19-Jun-07	32.21	5.9	5381.33
Dry Creek @ Wellsford Rd	Copper	17-Jul-07	68.57	3.9	7572.62
Dry Creek @ Wellsford Rd	Copper	14-Aug-07	65.83	5.3	9879.77
Dry Creek @ Wellsford Rd	Copper	11-Sep-07	61.02	3.3	5702.08
Dry Creek @ Wellsford Rd	Copper	22-Apr-08	39.8	4.7	5296.98
Dry Creek @ Wellsford Rd	Copper	20-May-08	38.3	3.8	4121.26
Dry Creek @ Wellsford Rd	Copper	17-Jun-08	38.94	3.7	4079.86
Dry Creek @ Wellsford Rd	Copper	22-Jul-08	63.22	3.2	5728.64
Dry Creek @ Wellsford Rd	Copper	19-Aug-08	52.75	5.3	7916.73
Dry Creek @ Wellsford Rd	Copper	23-Sep-08	33.48	2.3	2180.52
Dry Creek @ Wellsford Rd	Diuron	17-Apr-07	30.71	0.29	252.19
Dry Creek @ Wellsford Rd	Diuron	22-Apr-08	39.8	0.2	225.40

Source Identification and Outreach

The Coalition addressed source identification in the 2009 Management Plan Update. The Coalition focuses its efforts on applied pesticides, and PUR data are reviewed to determine the most frequent crop type and timing associated with applications. The Coalition conducted additional and upstream MPM during the months when exceedances occurred in the past to better characterize water quality the subwatershed. Throughout 2009, the Coalition actively engaged in grower outreach and education to address Dry Creek Management Plan's highest priority constituents. This outreach included general surveys used to determine initial management practice utilization, followed by grower meetings and mailings to educate growers about additional management practices and their importance in achieving acceptable water quality, and lastly individual grower meetings to continue to inform growers about management practices as well as to assess expected future implementation of additional management practices.

Priority A/B Constituents

Chlorpyrifos is the only priority A/B constituent under the Dry Creek Management Plan.

Chlorpyrifos

Nine exceedances of the chlorpyrifos WQTL of 0.015 µg/L occurred during the latter months of irrigation seasons in samples collected between 2005 and 2009 (Table I-7), including one exceedance in August of 2009. The Coalition used a combination of monitoring data and evaluation of Pesticide Use Report (PUR) data to identify possible sources.

PUR data are reviewed for the number of monthly chlorpyrifos applications, pounds active ingredient (AI) applied and acres treated (Table I-10, Figure I-2). The greatest amount of chlorpyrifos use occurred in 2005 (9,997 lbs) with the lowest amount of use occurring in 2008 (4,326 lbs, Table I-10, Figure I-2). The greatest chlorpyrifos use has occurred consistently in the month of July; however in 2009 the largest amount of chlorpyrifos was applied in May (Table I-10, Figure I-2). Exceedances often occur in the late summer (Figure I-2). The most recent chlorpyrifos applications preceding the 2009 August chlorpyrifos exceedances occurred in July 2009.

The Coalition also used PUR data to determine which crops receive the most applications of chlorpyrifos. The highest application rates are associated with almond/walnut orchards, corn, oranges, and grapes (Table I-11). The largest amount of chlorpyrifos applied from 2005 through 2009 was to almonds (15,958.278 lbs) followed by walnuts (12,287 lbs, Table I-11). The most common product containing chlorpyrifos used within this subwatershed was Lorsban (Table I-11).

Table I-10. Number of chlorpyrifos applications, pounds AI applied, and acres treated by month for January 2005 through December 2009 in the Dry Creek @ Wellsford Rd site subwatershed. If a month is not included in the table, no applications were made.

Month/Year	Number of Chlorpyrifos Applications	Pounds of AI Applied	Acres Treated
January, 2005	1	152.5	75
February, 2005	3	175.9	94
March, 2005	3	120.9	111
April, 2005	6	464.6	521.5
May, 2005	22	2617.1	2022.41
June, 2005	15	1970.3	1244
July, 2005	50	3923.3	2312
August, 2005	16	569.8	425.75
September, 2005	1	2.0	0.75
February, 2006	1	8.0	83
March, 2006	4	98.1	60.5
May, 2006	27	2881.3	1842.41
June, 2006	14	842.4	538.9
July, 2006	55	3042.9	2625
August, 2006	27	852.8	475
September, 2006	5	123.9	62.5
October, 2006	3	137.4	73.91
December, 2006	1	29.9	15
March, 2007	1	10.2	23
April, 2007	7	139.6	260.4
May, 2007	23	2486.1	1598.9
June, 2007	24	887.9	597
July, 2007	40	2585.9	1611.2
August, 2007	11	414.7	207.75
September, 2007	1	11.6	11
October, 2007	1	117.6	63.3
November, 2007	4	247.4	133.15
January, 2008	1	29.9	14
May, 2008	20	840.6	595
June, 2008	14	858.9	714
July, 2008	41	2012.0	1565.61
August, 2008	10	499.4	350
September, 2008	2	85.6	48
March, 2009	4	381.2	202.98
April, 2009	6	389.9	164.4
May, 2009	19	1355.6	668.5
June, 2009	9	798.2	701

Month/Year	Number of Chlorpyrifos Applications	Pounds of AI Applied	Acres Treated
July, 2009	22	1049.7	611.65
August, 2009	7	294.9	290.75
September, 2009	4	122.5	70.75
November, 2009	3	487.4	259.52
Summaries by Year			
2005 Total	117	9996.5	6806.41
2006 Total	137	8016.6	5776.22
2007 Total	112	6901.1	4505.7
2008 Total	88	4326.4	3286.61
2009 Total	74	4879.6	2969.55
Total	528	34,120.2	23,344.49

Figure I-2. Pounds of chlorpyrifos applied within the Dry Creek @ Wellsford Rd site subwatershed by month for 2005-2009. Asterisk (*) denotes months with exceedances.




Table I-11. Total pounds of AI for chlorpyrifos based on PUR data from 2005-2009 within the Dry Creek @ Wellsford Rd subwatershed.

Chemical Name	Commodity	Product Name	Lbs AI Applied	
CHLORPYRIFOS	ALFALFA	NUFOS 4E	10.168	
	ALFALFA (FORAGE - FODDER) (ALFALFA HAY)	NUFOS 4E	11.225	
	ALMOND		CHLORPYRIFOS 4E AG	382.772
			GOVERN 4E INSECTICIDE	890.849
			LORSBAN 4E INSECTICIDE	596.842
			LORSBAN 4E-HF	1439.236
			LORSBAN ADVANCED	478.890
			LORSBAN-4E	10633.721
			NUFOS 4E	1487.487
	WARHAWK	48.482		
	APPLE	LORSBAN 4E-HF	3.520	
	APRICOT	LORSBAN 4E-HF	8.972	
	CORN (FORAGE - FODDER)		LORSBAN 15G GRANULAR INSECTICIDE	1031.400
			LORSBAN 4E-HF	442.863
			LORSBAN-4E	111.917
	CORN FOR/FOD		GOVERN 4E INSECTICIDE	547.317
			LORSBAN 15G GRANULAR INSECTICIDE	104.544
			LORSBAN 4E-HF	45.604
			LORSBAN ADVANCED	94.745
			LORSBAN-4E	308.153
			NUFOS 15G	618.000
			NUFOS 4E	96.959
	WARHAWK	542.711		
	GRAPES, WINE		LORSBAN 4E-HF	98.137
			LORSBAN ADVANCED	868.574
			LORSBAN-4E	548.892
	N-OUTDR CONTAINER/FLD GRWN PLANTS	LORSBAN 75WG	90.000	
N-OUTDR GRWN TRNSPLNT/PRPGTV MTRL	LORSBAN-4E	244.121		
OP-DEC. TREE		LORSBAN 15G GRANULAR INSECTICIDE	1.125	
		LORSBAN 75WG	46.140	
WALNUT		CHLORPYRIFOS 4E AG	84.435	
		GOVERN 4E INSECTICIDE	40.401	
		LORSBAN 4E INSECTICIDE	240.034	

Chemical Name	Commodity	Product Name	Lbs AI Applied
		LORSBAN 4E-HF	666.949
		LORSBAN ADVANCED	362.454
		LORSBAN-4E	2453.534
		NUFOS 4E	1502.210
		WARHAWK	101.003
	WALNUT (ENGLISH WALNUT, PERSIAN WALNUT)	CHLORPYRIFOS 4E AG	1047.698
		LORSBAN 4E INSECTICIDE	300.195
		LORSBAN 4E-HF	1584.866
		LORSBAN-4E	3579.714
		NUFOS 4E	323.331
ALFALFA – Total Pounds Chlorpyrifos Applied			21.393
ALMOND – Total Pounds Chlorpyrifos Applied			15958.278
APPLE– Total Pounds Chlorpyrifos Applied			3.520
APRICOT – Total Pounds Chlorpyrifos Applied			8.972
CORN – Total Pounds Chlorpyrifos Applied			3944.212
GRAPES – Total Pounds Chlorpyrifos Applied			1515.603
N-OUTDOOR PLANTS – Total Pounds Chlorpyrifos Applied			334.121
OP-DEC. TREE – Total Pounds Chlorpyrifos Applied			47.265
WALNUT – Total Pounds Chlorpyrifos Applied			12286.824
Total pounds chlorpyrifos applied (2005 - 2009)			34,120.189

In the 2009 Management Plan Update, the Coalition determined the most probable source of chlorpyrifos in the Dry Creek waterway was spray drift from adjacent orchards, particularly during the months July and August. Consequently, outreach efforts focus on encouraging management practices aimed at reducing spray drift.

The results from the general surveys administered to all members in the subwatershed are summarized in the ESJWQC General Survey Summary Report submitted January 30, 2009 and were used as a starting point to evaluate management practice implementation within the subwatershed. The general survey assessed practices in use from May 17, 2007 to December 31, 2008. The Coalition noted that management practices specific to spray application and orchard and vineyard water management are not widely implemented across the Coalition region within the subwatershed. For example, out of survey responses, members representing 64 parcels, which accounts for 4,372 acres in the subwatershed, indicated they perform sprayer calibrations. Thus, of the 23,331 irrigated acres in the Dry Creek subwatershed, 19% of acres receive applications from currently calibrated sprayers. Fewer members indicated they allow grass to grow in the center rows of orchards and/or vineyards; grass center rows were present in only 3,678 acres, or 16%, in the subwatershed. And very few growers maintain native buffers along streams and waterways, representing only 92 acres. When assessing the percentages of management practices utilized in the watershed, the Coalition understands that not all

management practices are applicable or even necessary for all irrigated acres. Thus, the Coalition designed outreach to affect an increase in relevant and effective management practices in parcels adjacent to the creek which should result in the greatest improvement in water quality.

Coalition outreach since 2007 has included grower meetings and the mailing/distribution of information. A complete list of Coalition Outreach during 2009 is provided in the Summary of Coalition Outreach Activities section of this report. Prior to the 2009 irrigation season, several group grower meetings were held for members in the Dry Creek subwatershed to make them aware of the current issues and concerns specific to pesticide applications in the upcoming irrigation season and relevant management practices to protect and improve water quality in Dry Creek. The meetings also allowed the Coalition to update members on the strategy and status of the Dry Creek Management Plan.

Additionally, to gain more information about potential sources of discharge into the Dry Creek @ Wellsford Rd site subwatershed, the Stanislaus County Agricultural Commissioner's staff walked the waterway in November 2007 and documented all drains using GPS and digital photos. The Coalition mapped the drains and used this information to further assess possible sources of water quality impairments and to compile a list of growers targeted for outreach. The Coalition notified individuals owning the parcels where drains were found and conducted a joint meeting with the Stanislaus County Agriculture Commissioner in June 2008.

The Coalition then turned its attention to individual contacts. Targeted growers were selected based on their potential to drain or drift to the creek. Of the 16,111 acres in Dry Creek that have the potential to drain directly or indirectly to the creek, 6,658 acres are owned and operated by Coalition members. The Coalition eventually developed a list of 22 targeted members that represented 6,116 acres that have the potential to discharge to the creek. Individual meetings occurred starting in January, 2009 and continued through August, 2009. Coalition representatives met with growers on their property to discuss the Dry Creek Management Plan and offer management practice recommendations specific to the parcel and land owner/operator. This gave the Coalition a chance to better assess currently used management practices as well as planned management practice implementation. Individual contacts focused mainly on chlorpyrifos exceedances however all water quality results were reviewed and discussed including exceedances of copper, diuron, *E. coli*, DO, SC, and pH, and toxicity to *Selenastrum* and *Hyaella*. To date, the Coalition has contacted all 22 targeted members, which represents 6,116 acres or 9% of the total subwatershed acreage. The final targeted member acreage is less than the potential to drain acreage (members) due to parcels/members being dropped from the original targeted list because the acres were 1) not currently farmed, 2) were pasture with no pesticide use, 3) determined not to drain or have the potential to drift to the creek, or 4) there was no reported pesticide use. The Coalition plans to contact targeted members in the spring of 2010 to determine if the owners' planned management practices have been implemented, and will continue to conduct outreach on a grower group level and monitor the creek to assess water quality.

Priority C Constituents

Water samples from Dry Creek @ Wellsford Rd had exceedances for copper and diuron priority C constituents.

Copper

Copper has exceeded the hardness-based WQTL three times in this subwatershed from 2005 through 2009. The Coalition began monitoring for both the total and dissolved copper fractions in October 2008 to better characterize copper contamination in the subwatershed.

PUR data are reviewed for the number of monthly copper applications, pounds active ingredient (AI) applied, and acres treated (Table I-12, Figure I-3). The greatest amount of copper was applied in 2005 (30,342 lbs) and the lowest amount applied in 2008 (8,934 lbs) and 2009 (11,286 lbs, Table I-12, Figure I-3). In 2007, 2008, and 2009 the greatest amount of copper use occurred in April whereas in 2004 the greatest use occurred in January and in 2005 during May (Table I-12, Figure I-3). Exceedances do not always coincide with the greatest use of copper (Figure I-3).

The largest amount of copper applied was associated with almond, peach, and walnut orchards, rice fields, and vineyards/grapes (Table I-13). The largest amount of copper from 2005 through 2009 was applied to almond (38,517 lbs), followed by walnuts (46,224.6 lbs, Table I-13). The most common product containing copper used within this subwatershed was DuPont Kocide and Kocide (Table I-13).

Table I-12. Number of copper applications, total pounds AI applied and total acres treated by month for January 2005 through December 2009 in the Dry Creek @ Wellsford site subwatershed. If a month is not included in the table, no applications were made.

Month/Year	Number of Copper Applications	Pounds of AI Applied	Acres Treated
January, 2005	29	8069.7	1338.41
February, 2005	51	3809.9	1905.25
March, 2005	38	4863.3	1340.5
April, 2005	47	6703.1	2099.3
May, 2005	25	2986.1	721.7
June, 2005	3	161.6	72
July, 2005	1	0.5	10
August, 2005	2	84.3	35.12
September, 2005	1	15.0	10
November, 2005	2	2404.9	605
December, 2005	8	1243.9	343
January, 2006	29	3714.1	916
February, 2006	48	3098.8	1844.5
March, 2006	8	454.7	503

Month/Year	Number of Copper Applications	Pounds of AI Applied	Acres Treated
April, 2006	51	4937.9	1196.88
May, 2006	47	8561.3	1671.75
June, 2006	3	803.6	126
August, 2006	2	49.5	22
November, 2006	4	177.8	57.5
December, 2006	4	353.9	51
January, 2007	20	6954.0	1517.2
February, 2007	46	2299.5	1562
March, 2007	27	3154.5	940
April, 2007	62	7496.8	2287
May, 2007	10	1491.9	224.75
June, 2007	3	938.1	100
July, 2007	4	344.0	86
August, 2007	2	48.0	12
October, 2007	1	32.3	3
November, 2007	5	1714.7	435
December, 2007	5	203.0	59
January, 2008	10	1927.3	501.5
February, 2008	34	1614.1	1134
March, 2008	13	1124.8	674
April, 2008	29	4146.7	1254
May, 2008	3	39.5	25
November, 2008	2	29.5	12
December, 2008	2	53.0	13.5
January, 2009	13	2352.2	577.28
February, 2009	29	851.6	682
March, 2009	1	7.3	18
April, 2009	40	4388.2	1166
May, 2009	17	3274.5	738.5
June, 2009	3	194.2	102.75
November, 2009	4	140.7	21
December, 2009	2	77.3	13.5
Summaries by Year			
2005 Total	207	30342.2	8480.28
2006 Total	196	22151.6	6388.63
2007 Total	185	24676.8	7225.95
2008 Total	93	8934.9	3614
2009 Total	109	11286.2	3319.03
Total	790	97,391.6	29,027.89

Figure I-3. Pounds of copper applied within the Dry Creek @ Wellsford Rd site subwatershed by month for 2005-2009. Asterisk (*) denotes months with exceedances.




Table I-13. Total pounds of AI for copper based on PUR data from 2005-2009 within the Dry Creek @ Wellsford Rd subwatershed.

Chemical Name	Commodity	Product Name	Lbs AI Applied
COPPER	ALMOND	BASIC COPPER 53	5399.800
		BASICOP	628.050
		CHAMP FORMULA 2 FLOWABLE	7109.926
		CHAMPION WETTABLE POWDER	610.957
		CUPROFIX ULTRA 40 DISPERS	127.625
		DUPONT KOCIDE 2000 FUNGICIDE/BACTERICIDE	1910.976
		DUPONT KOCIDE DF FUNGICIDE/BACTERICIDE	27.016
		HYDROX	4864.860
		KOCIDE 101	3328.156
		KOCIDE 2000	2040.694
		KOCIDE DF	1796.257
		NORDOX	49.350
		NORDOX 75 WG	5907.777
		NU-COP 50DF	4602.290
		NUCOP DF	112.500
		NUTRA-SPRAY ZN12-MN2-CU10	0.463
	APRICOT	NU-COP 50DF	6.160
		TRIANGLE BRAND COPPER SULFATE	89.100
	CHERRY	CHAMPION WETTABLE POWDER	107.800
		KOCIDE 101	57.750
	GRAPES, WINE	BRITZ COPPER SULFUR 15-25 DUST	690.800
		CHAMP FORMULA 2 FLOWABLE	979.388
		DUPONT KOCIDE 2000 FUNGICIDE/BACTERICIDE	42.997
		KOCIDE 2000	82.228
		KOCIDE DF	58.391
		NU-COP 3L	35.817
	NECTARINE	CHAMPION WETTABLE POWDER	492.800
		KOCIDE 101	154.000
	N-OUTDR CONTAINER/FLD GRWN PLANTS	KOCIDE 101	140.140
		KOCIDE DF	92.100
	N-OUTDR GRWN TRNSPLNT/PRPGTV MTRL	CHAMP FORMULA 2 FLOWABLE	386.446
	OLIVE	CUPROFIX ULTRA 40 DISPERS	2.666
	OP-DEC. TREE	DUPONT KOCIDE 2000 FUNGICIDE/BACTERICIDE	51.648
KOCIDE DF		29.472	
PEACH	BASIC COPPER 53	73.500	

Chemical Name	Commodity	Product Name	Lbs AI Applied
		CHAMPION WETTABLE POWDER	554.400
		DUPONT KOCIDE 2000 FUNGICIDE/BACTERICIDE	7.263
		HYDROX	161.700
		KOCIDE 101	16.170
		KOCIDE 2000	33.894
		KOCIDE DF	299.632
		NORDOX 75 WG	149.929
		NU-COP 50 WP	164.780
	PEACH PROCESSNG	DUPONT KOCIDE 2000 FUNGICIDE/BACTERICIDE	10.491
		KOCIDE DF	34.384
		NORDOX 75 WG	74.168
	PISTACHIO	CUPROFIX ULTRA 40 DISPERSS	582.736
	PLUM	CHAMPION WETTABLE POWDER	138.600
		NORDOX 75 WG	11.410
	PLUM (INCLUDES WILD PLUMS FOR HUMAN CONSUMPTION)	CHAMPION WETTABLE POWDER	46.200
	RICE	BLUE VIKING COPPER SULFATE STAR SHINE CRYSTAL	910.800
		CHIPCO COPPER SULFATE CRYSTALS	2346.300
		KOCIDE COPPER SULFATE STAR SHINE CRYSTALS	782.100
	RICE (ALL OR UNSPEC)	CHIPCO COPPER SULFATE CRYSTALS	2752.200
	WALNUT	CHAMP FLOWABLE	55.726
		CHAMP FORMULA 2 FLOWABLE	1042.754
		CHAMPION WETTABLE POWDER	1663.200
		CUPROFIX ULTRA 40 DISPERSS	255.960
		DUPONT GX-569 FUNGICIDE/BACTERICIDE	528.306
		DUPONT KOCIDE 101 FUNGICIDE/BACTERICIDE	123.200
		DUPONT KOCIDE 2000 FUNGICIDE/BACTERICIDE	5116.929
		DUPONT KOCIDE 3000 FUNGICIDE/BACTERICIDE	395.538
		DUPONT KOCIDE DF FUNGICIDE/BACTERICIDE	92.100
		HYDROX	677.600
		KENTAN DF	208.420
		KOCIDE 101	458.920
		KOCIDE 2000	3727.802
		KOCIDE DF	1274.050

Chemical Name	Commodity	Product Name	Lbs AI Applied
		NORDOX 75 WG	1254.305
		NU-COP 50 WP	281.820
		NU-COP 50DF	4392.080
	WALNUT (ENGLISH WALNUT, PERSIAN WALNUT)	CHAMP FLOWABLE	23.713
		CHAMP FORMULA 2 FLOWABLE	1604.720
		CHAMPION WETTABLE POWDER	308.000
		DUPONT KOCIDE 2000 FUNGICIDE/BACTERICIDE	309.888
		HYDROX	229.460
		KOCIDE 101	4965.214
		KOCIDE 2000	4136.822
		KOCIDE DF	11514.624
		NORDOX	450.072
		NORDOX 75 WG	46.145
		NU-COP 50DF	1087.240
		ALMOND - Total Pounds Copper Applied	38516.694
		APRICOT – Total Pounds Copper Applied	95.260
	CHERRY – Total Pounds Copper Applied	165.550	
	GRAPES – Total Pounds Copper Applied	1889.621	
	NECTARINE – Total Pounds Copper Applied	646.800	
	N-OUTDOOR PLANTS – Total Pounds Copper Applied	618.686	
	OLIVE – Total Pounds Copper Applied	2.666	
	OP-DEC. TREE – Total Pounds Copper Applied	81.120	
	PEACH – Total Pounds Copper Applied	1580.311	
	PISTACHIO – Total Pounds Copper Applied	582.736	
	PLUM – Total Pounds Copper Applied	196.210	
	RICE – Total Pounds Copper Applied	6791.400	
	WALNUT – Total Pounds Copper Applied	46224.608	
	Total pounds copper applied (2005 - 2009)	97,391.663	

In the 2009 Management Plan Update, the Coalition determined future outreach should focus on providing information to specific growers located adjacent to Dry Creek to encourage the retention of water and sediment on the fields, especially during the dormant season. This coincides with the Coalition’s management plan strategy outlined above under the chlorpyrifos outreach section. Orchard and vineyard operators will be advised to consider management practices to prevent copper from entering the waterway with runoff when applicable. Storm runoff management relevant to copper applications will also be discussed to prevent the winter exceedances.

Diuron

Diuron is a soluble herbicide applied in the Dry Creek watershed primarily in the months of October – May. The largest amount of diuron used was in 2006 (1,063 lbs) and the least amount used was in 2008 (160 lbs, Table I-14). The month with the largest amount AI applied varied from year to year with large applications in January and November 2006, January 2007, April 2008, and November 2009 (Table I-14, Figure I-4). The highest application rates and largest amount of diuron applied from 2005 through 2009 are associated with walnuts and grapes (including wine and raisin, Table I-15).

There are only two applications associated with the two storm 2007 exceedances within one month of sampling. A third application occurred approximately five weeks prior to the first exceedance and a fourth application occurred approximately seven weeks prior to the first exceedance. Diuron did not exceed the WQTL (2 µg/L) during 2008 monitoring. Dry Creek was sampled as a Core site during 2009, and diuron was not monitored as a part of 2009 MPM.

Table I-14. Number of diuron applications, total pounds AI applied and total acres treated by month for January 2005-December 2009 in the Dry Creek @ Wellsford Rd site subwatershed. If a month is not included in the table, no applications were made.

Month/Year	Number of Diuron Applications	Pounds of AI Applied	Acres Treated
January, 2005	2	40.5	42
February, 2005	1	9.8	6
March, 2005	1	40.2	25.76
April, 2005	3	72.4	70
June, 2005	1	34.2	17.5
November, 2005	3	36.1	47
January, 2006	2	348.1	338.5
February, 2006	3	93.0	38
May, 2006	2	41.8	33
June, 2006	1	233.9	320
November, 2006	4	336.3	430
December, 2006	1	10.1	7
January, 2007	2	220.4	156
February, 2007	1	29.3	40
November, 2007	1	215.6	275
December, 2007	6	128.1	409
February, 2008	2	9.0	11.3
March, 2008	1	7.7	24
April, 2008	1	58.5	40
July, 2008	4	39.6	39
August, 2008	2	28.4	26
October, 2008	1	17.0	42.5

Month/Year	Number of Diuron Applications	Pounds of AI Applied	Acres Treated
January, 2009	3	16.0	40
February, 2009	1	4.0	5
November, 2009	5	237.8	795
December, 2009	1	53.8	185
Summaries by Year			
2005 Total	11	233.1	208.26
2006 Total	13	1063.2	1166.5
2007 Total	10	593.4	880
2008 Total	11	160.2	182.8
2009 Total	10	311.6	1025
Total	55	2,361.5	3,462.6

Figure I-4. Pounds of diuron applied within the Dry Creek @ Wellsford Rd site subwatershed by month for 2005-2009. Asterisk (*) denotes months with exceedances.




Table I-15. Total pounds of AI for diuron based on PUR data from 2005-2009 within the Dry Creek @ Wellsford Rd subwatershed.

Chemical Name	Commodity	Product Name	Lbs AI Applied
DIURON	ALFALFA (FORAGE - FODDER) (ALFALFA HAY)	DIREX 4L	10.136
	GRAPES, WINE	DIREX 4L	371.521
		DIURON 4L	215.550
		DUPONT DIREX 4L HERBICIDE	412.721
	PASTURES (ALL OR UNSPEC)	DIREX 4L	19.006
	WALNUT	DIREX 4L	376.219
		DUPONT DIREX 4L HERBICIDE	55.377
	WALNUT (ENGLISH WALNUT, PERSIAN WALNUT)	DIREX 4L	900.881
ALFALFA – Total Pounds Diuron Applied			10.136
GRAPES – Total Pounds Diuron Applied			999.793
PASTURES – Total Pounds Diuron Applied			19.006
WALNUT – Total Pounds Diuron Applied			1332.477
Total pounds diuron applied (2005 - 2009)			2,361.411

Priority D Constituents

Water column toxicity to *Selenastrum capricornutum* and sediment toxicity to *Hyalella azteca* are listed as priority D constituents under the Dry Creek @ Wellsford Rd Site Subwatershed Management Plan.

***Selenastrum capricornutum* toxicity**

Selenastrum toxicity was not monitored during 2009 sampling at Dry Creek. *Selenastrum* toxicity occurred once during the 2008 storm season and was associated with an exceedance of copper. The two are closely tied as copper is used as an algaecide as well as a fungicide.

The Coalition's strategy for eliminating *Selenastrum* toxicity involves focusing on copper and diuron with individual contacts as well as conducting outreach regarding the retention of storm water runoff rather than discharging to drainage canals and creeks. If these two constituents can be prevented from entering storm water during the winter rainy season, the Coalition believes that *Selenastrum* toxicity can also be reduced or eliminated. If the Coalition finds that diuron and copper exceedances are eliminated and *Selenastrum* toxicity persists, then additional herbicides will be targeted for outreach and management.

***Hyalella azteca* toxicity**

As a result of 2008 monitoring, *Hyalella azteca* was listed as a Priority D constituent in the Dry Creek site subwatershed. Both the 2008 storm and irrigation seasons experienced a single toxicity each. Sediment monitoring was not conducted at Dry Creek during 2009.

The Coalition has begun to conduct management plan monitoring for sediment toxicity. As of 2009, the Coalition now tests for total organic carbon and grain size in all sediments and pyrethroids and chlorpyrifos in sediment that exhibits toxicity. Management Plan Monitoring is scheduled for this site in 2010 and 2011.

Priority E Constituents

The following priority E constituents are listed under the Dry Creek @ Wellsford Rd Site Subwatershed Management Plan: DO, *E. coli*, pH, SC.

Several DO and *E. coli* exceedances occurred throughout 2009 sampling. There were no pH exceedances. These constituents will remain low priority but will be discussed during individual contacts and annual grower meetings.

Evaluation

This subwatershed is one of the first three priority site subwatersheds within the ESJWQC and therefore the Coalition has focused its resources on identifying the sources of agricultural discharge within this subwatershed that could result in water quality impairments, extending outreach to individual Coalition members, and setting evaluation goals.

The Coalition's strategy for the Dry Creek subwatershed has been to target growers adjacent to the waterway that have the potential to discharge or drift. Outreach includes grower notification, management practice outreach and education, tracking of management practices implementation, and providing information to growers on studies of management practice efficacy. The Coalition believes that because land use in this subwatershed is predominately orchards and vineyards, outreach aimed toward orchard and vineyard relevant management practice implementation will target the largest audience and result in the greatest improvement of water quality.

Individual surveys to document current management practice implementation and assess future implementations have been completed and returned representing 100% of targeted growers. The Coalition is now in the process of assessing these results and planning the final stages of outreach.

Both the acres treated and pounds of chlorpyrifos applied have decreased overall since 2005, which may be a result of many factors including educating growers about alternative products. Usage has decreased in July and August, when the largest number of exceedances have occurred. Despite this, chlorpyrifos exceeded the WQTL in Dry Creek in August 2009 although the exceedance was traced to a non member dairy. Management Plan Monitoring in 2010 will further indicate if a change in management practices as a result of individual contacts has resulted in long-term improvements to downstream water quality.

II. DUCK SLOUGH @ HIGHWAY 99

Management Plan Constituents

Priority A/B

- Chlorpyrifos

Priority C

- Copper

Priority D

- *Selenastrum capricornutum* water column toxicity

Priority E

- Dissolved Oxygen
- *E. coli*
- pH
- Lead

Description of Duck Slough @ Hwy 99 Site Subwatershed

Duck Slough @ Hwy 99 (10,695 irrigated acres) – This site subwatershed is located upstream of the Duck Slough @ Gurr Road site and was selected to determine relative contribution of water quality impairments from the upstream portion of the Duck Slough subwatershed. Duck Slough originates in the Sierra foothills and flows west eventually joining with Deadman Creek in the western portion of the Coalition region. The monitoring site is located just east of Highway 99, south of Planada and Merced. Irrigated agriculture in this site subwatershed is primarily deciduous nuts, with truck crops and irrigated pasture the next most common land uses (Figure II-1). This site subwatershed includes an upstream location (Duck Slough @ Whealan Rd) which was sampled in 2008. Table II-1 includes the station name, station code and target latitude/longitudes for sites sampled within this subwatershed.


Table II-1. Coordinates of the Duck Slough site subwatershed sampling locations.

Station Name	Station Code	Target Latitude	Target Longitude
Duck Slough @ Hwy 99*	535XDSAHN	37.2501	-120.4100
Duck Slough @ Whealan Rd ^U	535XDSAWH	37.26149	-120.34325

*Original ESJWQC sampling site

^U Upstream sites

Figure II-1. Site subwatershed map of land use for the Duck Slough @ Hwy 99 sample site.



Subwatershed Monitoring History

Monitoring at Duck Slough began in the storm season of 2005 at Duck Slough @ Highway 99 and continued throughout the storm and irrigation seasons from 2006 through 2008 (Table II-2). Starting in October 2008, Duck Slough @ Hwy 99 was removed from the monitoring schedule except for MPM events which occurred during the irrigation season of 2009 (Table II-2). As of October 2008, Duck Slough @ Hwy 99 became an Assessment site and will be monitored again for normal monitoring in 2013 and 2014. The constituents sampled at this location from 2005-2009 are listed in Table II-3.

Management Plan Monitoring occurred at Duck Slough @ Hwy 99 in 2007, 2008 and 2009 for copper and chlorpyrifos and for *S. capricornutum* toxicity in 2009 only (Table II-4, Table II-5, and Table II-6). Duck Slough was monitored twice a month in 2007, at additional upstream locations in 2008 and monitored in 2009 during months of past exceedances.

Duck Slough is listed as an impaired waterbody on the proposed 2008 Central Valley Basin Plan 303d list for chlorpyrifos, *E. coli*, sediment toxicity and unknown toxicity. Agriculture is listed as the potential source for the impairment caused by chlorpyrifos; all other impairments have unknown sources.

Table II-2. Duck Slough @ Hwy 99 sampling events per season and year.

	2004	2005		2006		2007		2008			2009			
	Irrigation	Storm	Irrigation	Storm	Irrigation	Storm	Irrigation	Storm	Irrigation	Fall	Winter	Storm	Irrigation	Fall
Events Sampled	NA	2	5	2	5	2	6	2	6	NA	NA	NA	6	NA
Events Not Sampled	NA	0	0	0	0	0	0	0	0	NA	NA	NA	0	NA
Total	NA	2	5	2	5	2	6	2	6	NA	NA	NA	6	NA

NA indicates that this site was not sampled during this season/year.

Table II-3. Number of analyses performed per analyte in each sampling season and year for the Duck Slough @ Hwy 99 sample site. Only environmental samples with a sample replicate and lab replicate number of one are shown.

Method	Analyte	2005		2006		2007		2008			2009			
		Storm	Irrigation	Storm	Irrigation	Storm	Irrigation	Storm	Irrigation	Fall*	Winter*	Storm*	Irrigation	Fall*
Field and Physical Parameters														
EPA 110.2	Color	2	5	2	5	2	6	2	6					
EPA 160.1	Dissolved Solids	2	5	2	5	2	6	2	6					
EPA 180.1	Turbidity	2	5	2	5	2	6	2	6					
EPA 405.1	BOD				1	2	2							
EPA 415.1	Total Organic Carbon	2	5	2	5	2	6	2	6					
SM 9223 B	E. coli	2	5	2	5	2	6	2	6					
NA	Dissolved Oxygen	2	6	3	6	3	10	3	9				6	
NA	Specific Conductivity	2	6	3	6	3	10	3	9				6	
NA	pH	2	6	3	6	3	10	3	9				6	
Carbamates														
EPA 8321A	Aldicarb				5	2	6	2	6					
EPA 8321A	Carbaryl				5	2	6	2	6					
EPA 8321A	Carbofuran				5	2	6	2	6					
EPA 8321A	Diuron				5	2	6	2	6					
EPA 8321A	Linuron				5	2	6	2	6					
EPA 8321A	Methiocarb				5	2	6	2	6					
EPA 8321A	Methomyl				5	2	6	2	6					
EPA 8321A	Oxamyl				5	2	6	2	6					
Organochlorines														
EPA 8081A	DDD(p,p')				5	2	6	2	6					
EPA 8081A	DDE(p,p')				5	2	6	2	6					
EPA 8081A	DDT(p,p')				5	2	6	2	6					
EPA 8081A	Dicofol				5	2	6	2	6					
EPA 8081A	Dieldrin				5	2	6	2	6					

Method	Analyte	2005		2006		2007		2008			2009			
		Storm	Irrigation	Storm	Irrigation	Storm	Irrigation	Storm	Irrigation	Fall*	Winter*	Storm*	Irrigation	Fall*
EPA 8081A	Endrin				5	2	6	2	6					
EPA 8081A	Methoxychlor				5	2	6	2	6					
Organophosphates														
EPA 8141A	Azinphos methyl				5	2	7	2	6					
EPA 8141A	Chlorpyrifos	2	5	2	5	2	7	2	6				3	
EPA 8141A	Diazinon	2	5	2	5	2	7	2	6					
EPA 8141A	Dimethoate				5	2	7	2	6					
EPA 8141A	Disulfoton				5	2	7	2	6					
EPA 8141A	Malathion				5	2	7	2	6					
EPA 8141A	Methamidophos				5	2	6	2	6					
EPA 8141A	Methidathion				5	2	7	2	6					
EPA 8141A	Molinate				5	2	7	2	6					
EPA 8141A	Parathion, Methyl				5	2	7	2	6					
EPA 8141A	Phorate				5	2	7	2	6					
EPA 8141A	Phosmet				5	2	7	2	6					
EPA 8141A	Thiobencarb				5	2	7	2	6					
Pyrethroids														
EPA 8081A	Bifenthrin			2	5	2	6							
EPA 8081A	Cyfluthrin, total			2	5	2	6							
EPA 8081A	Cyhalothrin, lambda, total	2	5	2	5	2	6							
EPA 8081A	Cypermethrin, total	2	5	2	5	2	6							
EPA 8081A	Esfenvalerate/ Fenvalerate, total	2	5	2	5	2	6							
EPA 8081A	Permethrin, total	2	5	2	5	2	6							
Triazines														
EPA 547M	Glyphosate				5	2	6	2	6					
EPA 549.2M	Paraquat dichloride				5	2	6	2	6					
EPA 619	Atrazine				5	2	6	2	6					

Method	Analyte	2005		2006		2007		2008			2009			
		Storm	Irrigation	Storm	Irrigation	Storm	Irrigation	Storm	Irrigation	Fall*	Winter*	Storm*	Irrigation	Fall*
EPA 619	Cyanazine				5	2	6	2	6					
EPA 619	Simazine				5	2	6	2	6					
Metals (Total)														
EPA 200.8	Arsenic				5	2	6	2	6					
EPA 200.8	Boron				5	2	6	2	6					
EPA 200.8	Cadmium				5	2	6	2	6					
EPA 200.8	Copper				5	2	8	2	6				4	
EPA 200.8	Lead				5	2	6	2	6					
EPA 200.8	Nickel				5	2	6	2	6					
EPA 200.8	Selenium				5	2	2	2	6					
EPA 200.8	Zinc				5	2	6	2	6					
Metals (Dissolved)														
EPA 200.8	Cadmium													
EPA 200.8	Copper												4	
EPA 200.8	Lead													
EPA 200.8	Nickel													
EPA 200.8	Zinc													
Nutrients														
SM 2340 C	Hardness as CaCO3				5	2	8	2	6				4	
EPA 300.0	Nitrate as N				5	2	6	2	6					
EPA 350.2	Ammonia as N				5	2	6	2	6					
EPA 351.3	Nitrogen, Total Kjeldahl				5	2	6	2	6					
EPA 354.1	Nitrite as N				5	2	6	2	6					
EPA 353.2	Nitrate + Nitrite as N													
EPA 365.2	Orthophosphate as P				5	2	6	2	6					
EPA 365.2	Phosphate as P				5	2	6	2	6					
Toxicity														
EPA 821/R-02-012	Ceriodaphnia dubia	2	5	2	6	2	7	2	6					

Method	Analyte	2005		2006		2007		2008			2009			
		Storm	Irrigation	Storm	Irrigation	Storm	Irrigation	Storm	Irrigation	Fall*	Winter*	Storm*	Irrigation	Fall*
EPA 821/R-02-012	Pimephales promelas	2	5	2	5	2	7	2	6					
EPA 821/R-02-013	Selenastrum capricornutum	2	7	2	5	2	7	2	7				1	
EPA 600/R-99-064	Hyalella azteca			1	1	1	1	1	2					

*Site not sampled during the 2008 fall and 2009 winter, storm and fall seasons.

Table II-4. Duck Slough site subwatershed. 2007 Management Plan additional (A) sampling schedule for chlorpyrifos and copper. "X" indicates the site, month, and analyte sampled.

Sample site	Date	Type	Chlorpyrifos	Copper
Duck Slough @ Hwy 99	24-Jul-07	A	x	
Duck Slough @ Hwy 99	21-Aug-07	A		x
Duck Slough @ Hwy 99	18-Sep-07	A		x

Table II-5. Duck Slough site subwatershed. 2008 Management Plan upstream (U) sampling schedule for chlorpyrifos and copper. X indicates the site, month and analyte sampled.

Station Name	Date	Type	Chlorpyrifos	Copper
Duck Slough @ Whealan Rd	29-Apr-08	U		X
Duck Slough @ Whealan Rd	27-May-08	U	X	
Duck Slough @ Whealan Rd	24-Jun-08	U		X
Duck Slough @ Whealan Rd	29-Jul-08	U	X	X
Duck Slough @ Whealan Rd	26-Aug-08	U		X
Duck Slough @ Whealan Rd	30-Sep-08	U		X

Table II-6. Duck Slough site subwatershed. 2009 Management Plan sampling schedule for chlorpyrifos, copper and *Selenastrum capricornutum* toxicity. "X" indicates the site, month and analyte sampled.

Site Name	Sample Date	<i>S. capricornutum</i>	Copper	Chlorpyrifos
Duck Slough @ Hwy 99	21-Apr-09	X		
Duck Slough @ Hwy 99	19-May-09			X
Duck Slough @ Hwy 99	16-Jun-09		X	
Duck Slough @ Hwy 99	21-Jul-09		X	X
Duck Slough @ Hwy 99	18-Aug-09		X	
Duck Slough @ Hwy 99	22-Sep-09		X	X

Exceedance History

Sampling at Duck Slough @ Hwy 99 resulted in exceedances of field and physical parameters, *E. coli*, metals, and chlorpyrifos. Toxicity in the water column to *Ceriodaphnia dubia* and *Selenastrum capricornutum* and in the sediment to *Hyalella azteca* also occurred at this monitoring site (Table II-7). During ambient water monitoring at Duck Slough @ Hwy 99 (including upstream sampling at Duck Slough @ Whealan Rd), exceedances of numerous field, physical, inorganic and organic parameters occurred including DO (2), pH (3), and *E. coli* (12, Table II-7). Exceedances of the chlorpyrifos (4), copper (14), and lead (11) WQTL have occurred. Samples taken from Duck Slough have also resulted in water column toxicity to *Ceriodaphnia* once and to *Selenastrum* three times. Sediment toxicity to *Hyalella* occurred twice, both during the 2008 irrigation season (Table II-7).

In 2009 there was one exceedance of the DO WQTL that occurred in June at Duck Slough @ Hwy 99 during the MPM irrigation season.

All exceedances are listed in Table II-7 by season and date and are based on WQTLs listed in the introduction of the ESJWQC Management Plan. The priority level (A-E) assigned to each constituent is listed in the bottom row of Table II-7 and is determined using the ESJWQC Management Plan prioritization process flow chart (Figure 4). Dissolved oxygen is now under a management plan following the exceedance during the 2009 sampling season. *Selenastrum capricornutum* and *Hyalella azteca* toxicity are listed as priority level D since a review of past exceedances.

Table II-7. All exceedances experienced in samples collected from locations within the Duck Slough site subwatershed between July 2004 and December 2009 (sorted by season and date). If the water quality trigger limit is based on hardness then the hardness value is shown in parenthesis; otherwise the WQTL used to evaluate the data is listed in the header after the analyte.

Station Name	Season	Sample Date	Oxygen, Dissolved, 7 mg/L	pH, 6.5-8.5 units	<i>E. coli</i> , 235 MPN/100 mL	Copper, Total, 1,300 µg/L	Lead, Total, 15 µg/L	Chlorpyrifos, 0.015 µg/L	<i>C. dubia</i> , Survival (%)	<i>H. azteca</i> , Survival (%)	<i>S. capricornutum</i> , Total Cell Count
Duck Slough @ Hwy 99	Storm	3/21/2005			1600						
Duck Slough @ Hwy 99	Irrigation	5/10/2005			1600						
Duck Slough @ Hwy 99	Irrigation	7/12/2005						0.026			1320000
Duck Slough @ Hwy 99	Storm	3/15/2006			900						
Duck Slough @ Hwy 99	Irrigation	5/17/2006		8.57	280		5.2 (3.02)	0.27	0		
Duck Slough @ Hwy 99	Irrigation	6/14/2006			260						
Duck Slough @ Hwy 99	Irrigation	8/8/2006				3.4 (2.4)	2.3 (0.41)				
Duck Slough @ Hwy 99	Irrigation	9/13/2006	6.72		340	19 (5.0)	24 (1.25)				
Duck Slough @ Hwy 99	Storm	2/12/2007			2400	31 (10.1)	15 (3.59)				
Duck Slough @ Hwy 99	Storm	2/28/2007			2400						
Duck Slough @ Hwy 99	Irrigation	4/24/2007				4.1 (3.7)	1.5 (0.81)				
Duck Slough @ Hwy 99	Irrigation	6/26/2007				3.0 (2.4)	0.68 (0.41)				
Duck Slough @ Hwy 99	Irrigation	7/24/2007				3.5 (3.0)	0.64 (0.57)				
Duck Slough @ Hwy 99	Irrigation	7/31/2007		8.8				0.042			
Duck Slough @ Hwy 99	Irrigation	8/21/2007				5.5 (3.3)	1.1 (0.69)				
Duck Slough @ Hwy 99	Irrigation	8/28/2007				3.1 (2.4)					
Duck Slough @ Hwy 99	Irrigation	9/18/2007			610	6.9 (2.8)	1.8 (0.52)				
Duck Slough @ Hwy 99	Storm	1/25/2008			>2400						
Duck Slough @ Hwy 99	Storm	2/25/2008			>2400	9.9 (8.0)					
Duck Slough @ Hwy 99	Sediment	3/4/2008		8.65							

Station Name	Season	Sample Date	Oxygen, Dissolved, 7 mg/L	pH, 6.5-8.5 units	<i>E. coli</i> , 235 MPN/100 mL	Copper, Total, 1,300 µg/L	Lead, Total, 15 µg/L	Chlorpyrifos, 0.015 µg/L	<i>C. dubia</i> , Survival (%)	<i>H. azteca</i> , Survival (%)	<i>S. capricornutum</i> , Total Cell Count
Duck Slough @ Hwy 99	Irrigation	4/29/2008			280						937637
Duck Slough @ Hwy 99	Irrigation	5/7/2008									18219
<i>Duck Slough @ Whealan Rd</i>	<i>Irrigation</i>	<i>6/24/2008</i>				73 (5.0)					
Duck Slough @ Hwy 99	Irrigation	7/29/2008				2.7 (2.6)	0.69 (0.5)				
Duck Slough @ Hwy 99	Irrigation	8/26/2008					0.72 (0.69)				
<i>Duck Slough @ Whealan Rd</i>	<i>Irrigation</i>	<i>8/26/2008</i>				3.4 (1.9)					
Duck Slough @ Hwy 99	Sediment	8/28/2008								84	
Duck Slough @ Hwy 99	Irrigation	9/30/2008						0.034			
<i>Duck Slough @ Whealan Rd</i>	<i>Irrigation</i>	<i>9/30/2008</i>				3.7 (1.3)					
Duck Slough @ Hwy 99	Sediment	10/2/2008								87	
Duck Slough @ Hwy 99	Irrigation	6/16/2009	6.78								
Constituent Priority			E	E	E	C	E	A/B	NP	NP¹	D

NP- Not prioritized. Fewer than two exceedances for this constituent at this site within three years and currently no TMDL for constituent
NP¹ – Not prioritized; both toxic samples were from the same sampling event (sample and resample to test for persistence).

2007 - 2009 Management Plan Monitoring Results

Management Plan monitoring results are included in Table II-8 for chlorpyrifos, copper and *S. capricornutum* for the years from 2007 through 2009.

2007

In 2007, Duck Slough @ Hwy 99 was monitored monthly for chlorpyrifos and copper from April through September. Additional monitoring was conducted in July for chlorpyrifos and in August for copper. Chlorpyrifos was detected in samples collected during normal monitoring below the WQTL (0.011 µg/L), samples collected one week later on July 31, 2007 exceeded the WQTL (0.42 µg/L, Table II-8).

Copper exceedances (based on total copper) occurred in samples collected in 2007 during all months from February – September (except for May) including additional samples collected on August 28, 2007 (Table II-8).

2008

In accordance with the 2008 upstream MPM schedule, Duck Slough @ Whealan Road was monitored for chlorpyrifos during May and July 2008 and for copper during April, June, July, August, and September 2008 (previous section, Table II-5). Chlorpyrifos was detected at both Hwy 99 and Whealan Rd during July monitoring and exceeded the chlorpyrifos WQTL at Duck Slough @ Highway 99 during September monitoring.

Copper exceedances occurred once during normal monitoring in July and three times at the upstream site during June, August, and September (based on total copper, Table II-8).

2009

Duck Slough @ Hwy 99 was monitored for chlorpyrifos during May, July, and September 2009 during MPM; normal monitoring did not occur in this subwatershed during 2009. There were no exceedances or detections of chlorpyrifos in samples collected from Duck Slough during 2009.

There were also no exceedances of the copper WQTL in 2009 MPM samples analyzed for total and dissolved copper collected from June through September 2009. This was the first year that both total and dissolved copper were analyzed. In previous years, the WQTL was based on total copper which used hardness to determine the concentration considered toxic. In 2009 the determination was based on the dissolved fraction only.

Toxicity to *S. capricornutum* was also part of 2009 MPM; the sample collected in April 2009 was not toxic.

Management Plan Monitoring for copper, chlorpyrifos, and *Selenastrum* toxicity are planned for the 2010 irrigation season. Management Plan Monitoring during February 2011 will also occur for copper.

Table II-8. Duck Slough site subwatershed. Normal monitoring (NM) and Management Plan Monitoring (MPM) results where 'A' indicates additional MPM (2007), and 'US' indicates upstream MPM for chlorpyrifos (2008), copper (2007-2009 irrigation seasons) and *Selenastrum* (2009). Exceedance values are in bold.

Month:		April	May	June	July	August	September
2007 NM (@ Hwy 99)	Date	4/24/07	5/29/07	6/26/07	7/24/07	8/21/07	9/18/07
	Chlorpyrifos (µg/L)	<0.003	<0.003	<0.003	0.011	<0.003	<0.003
	Copper (µg/L)	4.1	<0.06	3	3.5	5.5	6.9
2007 MPM A (@ Hwy 99)	Date	NA	NA	NA	7/31/07	8/28/07	9/25/07
	Chlorpyrifos (µg/L)	NA	NA*	NA	0.42	NA	NA
	Copper (µg/L)	NA	NA	NA	NA	3.1	2.5
2008 NM (@ Hwy 99)	Date	4/29/08	5/27/08	6/24/08	7/29/08	8/26/08	9/30/08
	Chlorpyrifos (µg/L)	<0.003	<0.003	<0.0026	0.0067	<0.003	0.034
	Copper (µg/L)	3.3	2.4	2.9	2.7	3	3.8
2008 MPM US (@ Whealan Rd)	Date	4/29/08	5/27/08	6/24/08	7/29/08	8/26/08	9/30/08
	Chlorpyrifos (µg/L)	NA	<0.003	NA	0.0081	NA	NA
	Copper (µg/L)	3.5	NA	73	3	3.4	3.7
2009 MPM (@ Hwy 99)	Date	4/21/09	5/19/09	6/16/09	7/21/09	8/18/09	9/22/09
	Chlorpyrifos (µg/L)	NA	<0.003	NA	<0.003	NA	<0.003
	Copper, total (µg/L)	NA	NA	2.3	2.5	2.4	2.7
	Copper, dissolved (µg/L)	NA	NA	0.61	0.7	0.61	0.25
	<i>S. capricornutum</i> toxicity (% Control)	222	NA	NA	NA	NA	NA

*Additional monitoring was initiated in June 2007; therefore no sample was collected for scheduled May additional monitoring.
NA - Not applicable. This site was not sampled during this month.

Load Calculations

Loads were calculated for the chlorpyrifos and copper detections (Table II-9) based on the following formula:

$$\text{Load} = \text{Discharge (cfs)} \times 28.317\text{L/ft}^3 \times \text{Concentration (milligram/L} \times 1000 \text{ or } \mu\text{g/L)}.$$

The load values presented for constituents in this report represent instantaneous loads only. These values should not be used to extrapolate loading over any period of time (e.g. weekly, monthly, seasonal or annual). The primary purpose for reporting instantaneous loads is to provide a normalization of the concentrations by flow for various constituents at the time the samples were collected.

Because very few discharge measurements have been taken at the Duck Slough @ Hwy 99 site since it is usually too deep to wade, only exceedances and/or detections that occurred when discharge was taken during an event are shown in Table II-9.

Table II-9. Duck Slough site subwatershed. Instantaneous load calculations for chlorpyrifos and copper where discharge was measured (sorted by analyte, site and date). Upstream site is in italics.

Station Name	Analyte Name	Sample Date	Discharge cfs	Concentration $\mu\text{g/L}$	Loading Rate $\mu\text{g/sec}$
<i>Duck Slough @ Whealan Rd</i>	<i>Chlorpyrifos</i>	<i>29-Jul-08</i>	<i>18.73</i>	<i>0.0081</i>	<i>4.30</i>
Duck Slough @ Hwy 99	Copper	12-Jul-06	58.3	2.7	4457.38
Duck Slough @ Hwy 99	Copper	18-Sep-07	0	6.9	0
<i>Duck Slough @ Whealan Rd</i>	<i>Copper</i>	<i>29-Apr-08</i>	<i>5.73</i>	<i>3.5</i>	<i>567.90</i>
<i>Duck Slough @ Whealan Rd</i>	<i>Copper</i>	<i>24-Jun-08</i>	<i>22.15</i>	<i>73</i>	<i>45787.17</i>
<i>Duck Slough @ Whealan Rd</i>	<i>Copper</i>	<i>29-Jul-08</i>	<i>18.73</i>	<i>3</i>	<i>1591.13</i>
<i>Duck Slough @ Whealan Rd</i>	<i>Copper</i>	<i>26-Aug-08</i>	<i>21.6</i>	<i>3.4</i>	<i>2079.60</i>
<i>Duck Slough @ Whealan Rd</i>	<i>Copper</i>	<i>30-Sep-08</i>	<i>7.56</i>	<i>3.7</i>	<i>792.08</i>

Source Identification and Outreach

Priority A/B Constituents

The only priority A/B constituent under the Duck Slough Management Plan is chlorpyrifos.

Chlorpyrifos

Chlorpyrifos exceeded the WQTL of 0.015 $\mu\text{g/L}$ four times in this subwatershed from 2005 through 2008 (May, twice in July and September); there were no exceedances during 2009 (Table II-7).

PUR data are reviewed for the number of monthly chlorpyrifos applications, pounds active ingredient (AI) applied, and acres treated (Table II-10, Figure II-2). The greatest amount of chlorpyrifos applied was in 2005 (4,790 lbs) with the lowest amount applied in 2007 (662 lbs, Table II-10, Figure II-2). In 2005, the greatest chlorpyrifos use occurred in July whereas in 2006, 2008 and 2009 the greatest use occurred in May and in 2007 during March (Table II-10, Figure II-2). Not all exceedances occurred during months of greatest use (Figure II-2).

The Coalition also used PUR data to assess which crops receive the most applications of chlorpyrifos. The highest application rates are associated with almonds and alfalfa (Table II-11). The largest amount of chlorpyrifos applied from 2005 through 2009 was to almonds (5,851 lbs) followed by corn (1,593 lbs, Table II-11). The most common product used within this subwatershed containing chlorpyrifos was Lorsban (Table II-11).

Table II-10. Number of chlorpyrifos applications, total pounds applied, and total acres treated by month for January 2005 through December 2009 in the Duck Slough @ Hwy 99 site subwatershed. If a month is not included in the table, no applications were made.

Month/Year	Number of Chlorpyrifos Applications	Pounds of AI Applied	Acres Treated
January, 2005	4	76.9	38.1
March, 2005	14	408.7	695
June, 2005	2	50.0	30
July, 2005	17	4226.8	812
September, 2005	1	28.0	14
January, 2006	6	119.7	59
March, 2006	3	56.1	151
May, 2006	13	706.5	433
June, 2006	1	137.6	69
July, 2006	5	570.2	508
August, 2006	5	199.9	136
September, 2006	2	145.6	80
January, 2007	2	29.9	15
March, 2007	15	370.0	433.81
April, 2007	3	28.8	57.42
June, 2007	1	19.9	10
July, 2007	1	10.2	10
August, 2007	4	204.0	261.33
March, 2008	15	290.4	347.53
May, 2008	8	674.8	442
June, 2008	1	10.2	8
July, 2008	6	261.8	522.9
August, 2008	7	264.0	426.5
March, 2009	1	57.5	136
May, 2009	4	825.9	632

Month/Year	Number of Chlorpyrifos Applications	Pounds of AI Applied	Acres Treated
July, 2009	2	70.1	140
September, 2009	2	25.9	103.4
Summaries by Year			
2005 Total	38	4790.4	1589.1
2006 Total	35	1935.5	1436
2007 Total	26	662.8	787.56
2008 Total	37	1501.2	1746.93
2009 Total	9	979.4	1011.4
Total	145	9,869.3	6,570.99

Figure II-2. Pounds of chlorpyrifos applied within the Duck Slough @ Hwy 99 site subwatershed by month for 2005-2009. Asterisk (*) denotes months with exceedances.




Table II-11. Total pounds AI for chlorpyrifos based on PUR data from 2005-2009 within the Duck Slough @ Hwy 99 subwatershed.

Chemical Name	Commodity	Product Name	Lbs AI Applied
CHLORPYRIFOS	ALFALFA	COBALT	57.529
		GOVERN 4E INSECTICIDE	184.604
		LOCK-ON INSECTICIDE	397.255
		LORSBAN 4E-HF	320.676
		LORSBAN-4E	42.404
		NUFOS 4E	51.855
	ALFALFA (FORAGE - FODDER) (ALFALFA HAY)	LORSBAN 4E INSECTICIDE	63.921
		LORSBAN 4E-HF	337.449
		LORSBAN 4E-HF INSECTICIDE	63.435
	ALMOND	CHLORPYRIFOS 4E AG	84.746
		GOVERN 4E INSECTICIDE	498.956
		LORSBAN 4E INSECTICIDE	175.710
		LORSBAN 4E-HF	1072.692
		LORSBAN-4E	3619.113
		NUFOS 4E	20.335
		WHIRLWIND	379.940
	CORN (FORAGE - FODDER)	CHLORPYRIFOS 4E AG	10.089
	CORN FOR/FOD	CHLORPYRIFOS 4E AG	121.754
		LORSBAN 15G GRANULAR INSECTICIDE	1451.281
		NUFOS 4E	10.168
	PEACH	CHLORPYRIFOS 4E AG	76.877
		LORSBAN 4E-HF	121.625
WALNUT	GOVERN 4E INSECTICIDE	15.150	
	LORSBAN-4E	406.509	
WALNUT (ENGLISH WALNUT, PERSIAN WALNUT)	LORSBAN 4E-HF	71.779	
	LORSBAN-4E	213.393	
ALFALFA - Total Pounds Chlorpyrifos Applied			1519.127
ALMOND – Total Pounds Chlorpyrifos Applied			5851.493
CORN – Total Pounds Chlorpyrifos Applied			1593.291
PEACH – Total Pounds Chlorpyrifos Applied			198.502
WALNUT – Total Pounds Chlorpyrifos Applied			706.831
Total pounds chlorpyrifos applied (2005 - 2009)			9,869.245

Previous analyses conducted by the Coalition in the 2009 Management Plan Update suggested that management of chlorpyrifos in this watershed should focus on providing information on a targeted basis to growers to encourage the proper management of irrigation tailwater. Also, TRS' with applications associated with past chlorpyrifos exceedances are close to and/or

adjacent to Duck Slough suggesting that drift could play a major role in generating the exceedances and should be one of the subjects of focused outreach. The analysis concluded that outreach should target specific growers applying chlorpyrifos and/or members farming parcels adjacent to the creek. Growers are encouraged to review their operation to determine if irrigation return flows are managed properly. Owners operating orchards are instructed to evaluate their applications to minimize spray drift. Growers are targeted for outreach prior to the irrigation season and heavy application periods. Although approximately 38% of the member acreage utilize grass orchard or vineyard row centers and approximately 31% of the member acreage utilizes recirculation and/or tailwater return systems, the Coalition anticipates that targeted outreach will increase the use of these practices and other irrigation water management practices (see the ESJWQC General Survey Summary Report submitted on January 30, 2009 for details on the management practices within this subwatershed).

Coalition outreach since 2007 has included grower meetings and the mailing/distribution of information. A complete list of Coalition Outreach during 2009 is provided in the Summary of Coalition Outreach Activities section of the Management Plan Update Report. The Coalition focused outreach to growers applying chlorpyrifos and those owning parcels with the potential to drain or drift to the creek. Prior to the 2009 irrigation season, several group grower meetings were held for members in the Duck Slough subwatershed to make them aware of the current issues and concerns specific to pesticide applications, and relevant management practices to protect and improve water quality in Duck Slough. The meetings also allowed the Coalition to update members on the strategy and status of the Duck Slough Management Plan.

Beginning in May 2009 and continuing through August 2009, the Coalition conducted individual meetings with targeted members. Targeted members were determined by considering a grower's chlorpyrifos use and if a grower's parcels had the potential to drain or drift to the creek. Of the 10,695 irrigated acres in the Duck Slough @ Hwy 99 subwatershed, 5,767 acres have the potential to drain or drift to the waterway and 4,440 of those acres are owned and operated by Coalition members. The Coalition eventually developed a list of 24 targeted members that represented 4,016 acres that have the potential to discharge to the creek. Coalition representatives then met with growers on their property to discuss the Duck Slough Management Plan and offer management practice recommendations specific to the land owner. Current management practices were assessed by Coalition representatives and additional management practices were discussed and encouraged. Individual contacts focused mainly on chlorpyrifos exceedances however all water quality results were reviewed and discussed including copper, *E. coli*, DO, lead, and pH, and toxicity to *Selenastrum*. To date, the Coalition has contacted all its targeted members, representing 4,016 acres or 23% of the acreage in the subwatershed. The Coalition re-contacted previously visited members via a group grower meeting held on February 19, 2010. Using the Coalition's Interactive Survey Devices, growers were polled to determine what additional management practices were implemented in 2009 and if additional practices are planned for 2010. The Coalition will continue to monitor the creek to assess water quality in 2010.

Priority C Constituents

Copper

Copper has exceeded the hardness based WQTL numerous times in this subwatershed from 2005 through 2009 based on results from total copper analysis; however there were no exceedances during 2009 (Table II-7).

PUR data are reviewed for the number of monthly copper applications, pounds active ingredient (AI) applied, and acres treated (Table II-12, Figure II-3). The amount of copper applied within the subwatershed has generally decreased. The greatest amount of copper applied was in 2005 (5,753.6 lbs) and the lowest amount applied was in 2009 (2,789.8 lbs). The greatest use occurred in January for 2005 and 2006; in 2008 and 2009 the greatest use occurred in December (Table II-12, Figure II-3) indicating winter season use is the most common. Not all exceedances occurred during months of the greatest use, and interestingly several exceedances occurred in later irrigation season months that had few or no reported applications (Figure II-3).

The largest applications were associated with almonds, nectarines, peaches, tomatoes, and walnuts (Table II-13). The largest amount of copper applied from 2005 through 2009 was to walnuts (4,948.881 lbs) followed by almonds (4,620.083 lbs) (Table II-13). The most common products containing copper used within this subwatershed were Kocide and Dupont Kocide (Table II-13).

Table II-12. Number of copper applications, total pounds applied and total acres treated by month for January 2005-December 2009 in the Duck Slough @ Hwy 99 site subwatershed. If a month is not included in the table, no applications were made.

Month/Year	Number of Copper Applications	Pounds of AI Applied	Acres Treated
January, 2005	12	1568.7	376.1
February, 2005	5	514.8	159.7
March, 2005	3	369.9	81
April, 2005	26	1363.2	483.25
May, 2005	10	497.1	214.5
August, 2005	3	139.8	127.8
September, 2005	2	82.4	74.7
October, 2005	1	25.8	21
November, 2005	1	69.1	45
December, 2005	5	1122.8	153
January, 2006	12	1872.3	321
February, 2006	9	814.0	251
March, 2006	4	288.0	360
April, 2006	12	1038.3	263.5
May, 2006	9	935.4	367

Month/Year	Number of Copper Applications	Pounds of AI Applied	Acres Treated
August, 2006	1	113.0	103
December, 2006	2	539.0	70
January, 2007	9	1502.7	223
February, 2007	8	533.7	280
March, 2007	3	309.2	115
April, 2007	10	422.9	147.5
May, 2007	1	8.3	33
December, 2007	5	247.0	88
January, 2008	6	826.5	136
February, 2008	2	650.3	42
April, 2008	1	27.7	15
May, 2008	5	318.9	165.5
June, 2008	1	17.9	25.84
November, 2008	1	166.0	60
December, 2008	3	901.0	100
January, 2009	2	847.7	102
February, 2009	1	35.6	58
April, 2009	12	596.1	294.5
May, 2009	6	303.2	170.5
August, 2009	1	69.3	30
November, 2009	1	77.0	10
December, 2009	4	861.0	73
Summaries by Year			
2005 Total	68	5753.6	1736.05
2006 Total	49	5600.0	1735.5
2007 Total	36	3023.9	886.5
2008 Total	19	2908.1	544.34
2009 Total	27	2789.8	738
Total	199	20,075.4	5,640.39

Figure II-3. Pounds of copper applied within the Duck Slough @ Hwy 99 site subwatershed by month for 2005-2009. Asterisk (*) denotes months with exceedances.




Table II-13. Total pounds AI for copper based on PUR data from 2005-2009 within the Duck Slough @ Hwy 99 subwatershed.

Chemical Name	Commodity	Product Name	Lbs AI Applied
COPPER	ALFALFA (FORAGE - FODDER) (ALFALFA HAY)	KOCIDE 2000	63.753
	ALMOND	BASIC COPPER 53	220.500
		COOKE BLUESTONE	8.250
		CUPROFIX ULTRA 40 DISPERS	602.217
		DUPONT KOCIDE 2000 FUNGICIDE/BACTERICIDE	117.822
		DUPONT KOCIDE 3000 FUNGICIDE/BACTERICIDE	59.008
		DUPONT KOCIDE DF FUNGICIDE/BACTERICIDE	35.612
		KOCIDE 101	887.040
		KOCIDE 2000	228.650
		KOCIDE DF	853.460
		NORDOX 75 WG	1607.524
	APRICOT	BASIC COPPER 53	58.800
		KOCIDE 2000	118.360
		KOCIDE DF	122.800
	CHERRY	NU-COP 50DF	154.000
	NECTARINE	BASIC COPPER 53	98.000
		DUPONT GX-569 FUNGICIDE/BACTERICIDE	27.660
		DUPONT KOCIDE 2000 FUNGICIDE/BACTERICIDE	42.368
		DUPONT KOCIDE 3000 FUNGICIDE/BACTERICIDE	23.050
		KOCIDE 101	1155.000
		KOCIDE 2000	201.750
	N-OUTDOOR PLANT	BASIC COPPER 53	1519.000
		DUPONT KOCIDE 3000 FUNGICIDE/BACTERICIDE	331.920
	N-OUTDR CONTAINER/FLD GRWN PLANTS	DUPONT GX-569 FUNGICIDE/BACTERICIDE	53.430
		DUPONT KOCIDE 3000 FUNGICIDE/BACTERICIDE	165.960
		KOCIDE 101	473.550
	PEACH	BASIC COPPER 53	2528.400
		DUPONT KOCIDE 2000 FUNGICIDE/BACTERICIDE	172.160
		KOCIDE 101	154.000
		KOCIDE DF	761.974
	PEPPERS (FRUITING VEGETABLE), (BELL, CHILI, ETC.)	KOCIDE 101	121.275
		KOCIDE DF	27.016

Chemical Name	Commodity	Product Name	Lbs AI Applied
	PISTACHIO (PISTACHE NUT)	CUPROFIX DISPERSS DRY FLOWABLE FUNGICIDE/BACTERICIDE FOR TURF AND ORNAMENTAL USE	8.303
	PLUM	DUPONT KOCIDE 2000 FUNGICIDE/BACTERICIDE	80.700
	PLUM (INCLUDES WILD PLUMS FOR HUMAN CONSUMPTION)	KOCIDE 101	154.000
	PRUNE	NORDOX 75 WG	519.173
	TOMATO	DUPONT KOCIDE 2000 FUNGICIDE/BACTERICIDE	19.368
		DUPONT KOCIDE 3000 FUNGICIDE/BACTERICIDE	43.449
		KOCIDE 101	422.153
		KOCIDE 2000	669.390
		KOCIDE DF	78.899
	TOMATO PROCESSING	CHAMPION WETTABLE POWDER	69.300
		DUPONT KOCIDE 3000 FUNGICIDE/BACTERICIDE	67.541
	WALNUT	DUPONT GX-569 FUNGICIDE/BACTERICIDE	212.982
		DUPONT KOCIDE 2000 FUNGICIDE/BACTERICIDE	648.828
		DUPONT KOCIDE 3000 FUNGICIDE/BACTERICIDE	355.431
		KOCIDE 101	309.232
		KOCIDE 2000	350.238
		DUPONT KOCIDE 2000 FUNGICIDE/BACTERICIDE	167.453
	WALNUT (ENGLISH WALNUT, PERSIAN WALNUT)	KOCIDE 101	1188.880
		KOCIDE 2000	1168.267
		KOP-HYDROXIDE 50W	396.550
		NORDOX 75 WG	151.020
		ALFALFA – Total Pounds Copper Applied	63.753
		ALMOND – Total Pounds Copper Applied	4,620.083
		APRICOT – Total Pounds Copper Applied	299.960
		CHERRY – Total Pounds Copper Applied	154.000
		NECTARINE – Total Pounds Copper Applied	1,547.828
		N-OUTDOOR PLANTS – Total Pounds Copper Applied	2543.86
		PEACH – Total Pounds Copper Applied	3,616.534
		PEPPERS – Total Pounds Copper Applied	148.291
		PISTACHIO – Total Pounds Copper Applied	8.303
		PLUM (INCLUES PRUNES) – Total Pounds Copper Applied	753.873
		TOMATO – Total Pounds Copper Applied	1370.1

Chemical Name	Commodity	Product Name	Lbs AI Applied
		WALNUT – Total Pounds Copper Applied	4,948.881
		Total pounds copper applied (2005 - 2009)	20,075.465

Previous analyses conducted by the Coalition and reported in the 2009 Management Plan Update suggested that a small number of applications in the watershed are responsible for past copper exceedances, most likely through irrigation runoff as most copper applications are specified as ground. The Coalition has thus focused its outreach to target specific growers with the potential to discharge to the creek, which coincides with the outreach strategy for chlorpyrifos. Growers should be encouraged to review their operation to determine if irrigation return flows are managed properly, particularly in the irrigation months when copper exceedances are common. Orchard operators are advised to consider storm runoff relevant to copper applications to prevent the winter exceedances of copper. No copper exceedances occurred in 2009 during the months when this site was sampled. Individual contacts that occurred within this subwatershed are described under the chlorpyrifos outreach section above and included discussions of copper exceedances and the above management practices.

Priority D Constituents

Water column toxicity to *Selenastrum capricornutum* and sediment toxicity to *Hyalella azteca* are listed as priority D Constituents under the Duck Slough @ Hwy 99 Site Subwatershed Management Plan.

***Selenastrum capricornutum* toxicity**

Toxicity to *Selenastrum* occurred in July 2005 and in April 2008 and was persistent in the resample collected in May 2008. Although there were not any copper exceedances associated with toxicity, the Coalition elected to list water column toxicity to *Selenastrum* and *Hyalella azteca* sediment toxicity as a priority D constituent because of the high frequency of copper exceedances at this site. There was no toxicity associated with the MPM sample collected in April 2009.

The Coalition’s strategy for eliminating *Selenastrum* toxicity is to focus on copper with individual contacts as well as conduct outreach regarding the retention of storm and irrigation water runoff rather than discharging to drainage canals and creeks. If copper can be prevented from entering waterways, the Coalition believes that *Selenastrum* toxicity can also be reduced or eliminated. If the Coalition finds that copper exceedances are eliminated but *Selenastrum* toxicity persists, then additional herbicides will be targeted for outreach and management.

Priority E Constituents

Priority E constituents include pH, DO, *E. coli*, and lead. Although these constituents will remain low priority, the Coalition will continue to collect these data and these constituents have been discussed during individual contacts and annual grower meetings.

Evaluation

Duck Slough @ Hwy 99 is one of the first three priority site subwatersheds within the ESJWQC and is in its second year of focused outreach (2008-2010). The Coalition's strategy for the Duck Slough subwatershed has been to target growers along or adjacent to the waterway that have the potential to discharge. Focus has been on both aerial application and tailwater retention management practices. Outreach includes grower notification, management practice outreach and education, tracking of management practices implementation, and providing information on special studies of management practice efficacy.

Individual surveys to document current management practices and assess planned implementations have been completed by 100% of targeted growers. The Coalition is now in the process of assessing these results and planning the final stages of outreach including re-contacting growers to identify newly implemented practices.

The amount of chlorpyrifos applied as well as the acreage treated with chlorpyrifos products has decreased since 2005. Results of Management Plan Monitoring conducted in 2009 included no exceedances of chlorpyrifos or copper and no toxicity to *S. capricornutum* indicating improved water quality. Management Plan Monitoring in 2010 will indicate if a change in management practices as a result of individual contacts has resulted in long-term improvements to downstream water quality.

III. PRAIRIE FLOWER DRAIN

Management Plan Constituents

Priority A/B

- Chlorpyrifos

Priority D

- Nitrate / Nitrate + Nitrite
- *Ceriodaphnia dubia* water column toxicity
- *Hyalella azteca* sediment toxicity

Priority E

- Ammonia
- Dissolved Oxygen
- pH
- Specific Conductivity
- Total Dissolved Solids
- *E. coli*
- *Selenastrum capricornutum* water column toxicity

Description of Prairie Flower Drain @ Crows Landing Rd Site Subwatershed

Prairie Flower Drain @ Crows Landing Rd (3,611 irrigated acres) – Relative to other drains in the western portion of the Coalition region, Prairie Flower Drain is longer and appears to drain mostly irrigated agriculture. Dairies and feedlots are ubiquitous in this part of the Coalition region and this drain may receive runoff from several dairies immediately upstream. Upstream agriculture is field crops (Figure III-1). Many of the drains in the eastern part of the Coalition region, including Prairie Flower Drain, were constructed to intercept shallow ground water and lower the water table sufficiently to allow agriculture. Consequently, agriculture in the watershed tends to be dominated by shallow rooted field crops. This site subwatershed includes an upstream location (Prairie Flower Drain @ Morgan Rd) which was sampled in 2008. Table III-1 includes the station name, station code and target latitude/longitudes for sites sampled within this subwatershed.


Table III-1. Coordinates of the Prairie Flower Drain site subwatershed sampling locations.

Station Name	Station Code	Target Latitude	Target Longitude
Prairie Flower Drain @ Crows Landing Road*	535XPFDCL	37.4422	-121.00236
Prairie Flower Drain @ Morgan Rd ^U	535XPFDMR	37.437875	-120.97566

* Original ESJWQC sampling site

^U Upstream sites

Figure III-1. Site subwatershed map of land use for the Prairie Flower Drain @ Crows Landing Rd sample site.



Subwatershed Monitoring History

Ambient monitoring began at Prairie Flower Drain @ Crows Landing Rd during the storm season of 2005 and continued uninterrupted through 2009 (Table III-2). Specific information on the analysis conducted across each of the monitoring seasons is provided below (Table III-3).

Prairie Flower Drain @ Crows Landing Rd will remain a Core Monitoring location under the new MRPP through 2010. In 2011 this site will be monitored as an Assessment site.

Management Plan Monitoring for the Coalition was initiated during the 2007 irrigation season. Additional monitoring for chlorpyrifos and toxicity to *Pimephales promelas* occurred in 2007 (Table III-4). Upstream monitoring at Prairie Flower Drain @Morgan Rd occurred for chlorpyrifos, nitrate, and toxicity to *Ceriodaphnia* and *Pimephales* in 2008 (Table III-5). Management Plan Monitoring for 2009 occurred at Prairie Flower Drain @ Crows Landing Rd for chlorpyrifos and toxicity to *Selenastrum* (Table III-6).

Prairie Flower Drain is not considered impaired under the proposed Basin Plan and is not listed on the 303d list. However the section of the San Joaquin River which receives water from Prairie Flower Drain is listed as impaired for BHC-alpha, boron, chlorpyrifos, DDE, DDT, SC, Group A pesticides, mercury, water temperature, and unknown toxicity.

Table III-2. Prairie Flower Drain @ Crows Landing Rd sampling events per season and year

	2004	2005		2006		2007		2008			2009			
	Irrigation	Storm	Irrigation	Storm	Irrigation	Storm	Irrigation	Storm	Irrigation	Fall	Winter	Storm	Irrigation	Fall
Events Sampled	NA	2	5	2	5	2	6	3	6	2	2	2	6	3
Events Not Sampled	NA	0	0	0	0	0	0	0	0	0	0	0	0	0
Total	NA	2	5	2	5	2	6	3	6	2	2	2	6	3

NA indicates that this site was not sampled during this season/year.

Table III-3. Number of analyses performed per analyte in each sampling season and year for the Prairie Flower Drain @ Crows Landing Rd sample site. Only environmental samples with a sample replicate and lab replicate number of one are shown.

Method	Analyte	2005		2006		2007		2008			2009			
		Storm	Irrigation	Storm	Irrigation	Storm	Irrigation	Storm	Irrigation	Fall	Winter	Storm	Irrigation	Fall
Field and Physical Parameters														
EPA 110.2	Color	2	5	2	5	3	6	2	6					
EPA 160.1	Dissolved Solids	2	5	2	5	2	6	3	6	2	2	2	6	2
EPA 160.2	Suspended Solids							1		2	2	2	6	2
EPA 180.1	Turbidity	2	5	2	5	2	6	3	6	2	2	2	6	2
EPA 405.1	BOD				1	2	2							
EPA 415.1	Total Organic Carbon	2	5	2	5	2	6	3	6	2	2	2	6	2
SM 9223 B	E. coli	2	5	2	5	2	6	3	6	2	2	2	6	2
NA	Dissolved Oxygen	2	5	4	6	3	13	5	10	2	2	3	6	2
NA	Specific Conductivity	2	5	4	6	3	13	5	10	2	2	3	6	2
NA	pH	2	5	4	6	3	13	5	10	2	2	3	6	2
Carbamates														
EPA 8321A	Aldicarb				5	2	6	2	6					
EPA 8321A	Carbaryl				5	2	6	2	6					
EPA 8321A	Carbofuran				5	2	6	2	6					
EPA 8321A	Diuron				5	2	6	2	6					
EPA 8321A	Linuron				5	2	6	2	6					
EPA 8321A	Methiocarb				5	2	6	2	6					
EPA 8321A	Methomyl				5	2	6	2	6					
EPA 8321A	Oxamyl				5	2	6	2	6					
Organochlorines														
EPA 8081A	DDD(p,p')				5	2	6	2	6					
EPA 8081A	DDE(p,p')				5	2	6	2	6					
EPA 8081A	DDT(p,p')				5	2	6	2	6					
EPA 8081A	Dicofol				5	2	6	2	6					
EPA 8081A	Dieldrin				5	2	6	2	6					

Method	Analyte	2005		2006		2007		2008			2009			
		Storm	Irrigation	Storm	Irrigation	Storm	Irrigation	Storm	Irrigation	Fall	Winter	Storm	Irrigation	Fall
EPA 8081A	Endrin				5	2	6	2	6					
EPA 8081A	Methoxychlor				5	2	6	2	6					
Organophosphates														
EPA 8141A	Azinphos methyl				5	2	6	2	6					
EPA 8141A	Chlorpyrifos	2	5	2	5	2	8	2	6				1	
EPA 8141A	Diazinon	2	5	2	5	2	6	2	6					
EPA 8141A	Dimethoate				5	2	6	2	6					
EPA 8141A	Disulfoton				5	2	6	2	6					
EPA 8141A	Malathion				5	2	6	2	6					
EPA 8141A	Methamidophos				5	2	6	2	6					
EPA 8141A	Methodathion				5	2	6	2	6					
EPA 8141A	Molinate				5	2	6	2	6					
EPA 8141A	Parathion, Methyl				5	2	6	2	6					
EPA 8141A	Phorate				5	2	6	2	6					
EPA 8141A	Phosmet				5	2	6	2	6					
EPA 8141A	Thiobencarb				5	2	6	2	6					
Pyrethroids														
EPA 8081A	Bifenthrin		1	2	5	2	6							
EPA 8081A	Cyfluthrin, total		1	2	5	2	6							
EPA 8081A	Cyhalothrin, lambda, total	2	5	2	5	2	6							
EPA 8081A	Cypermethrin, total	2	5	2	5	2	6							
EPA 8081A	Esfenvalerate/ Fenvalerate, total	2	5	2	5	2	6							
EPA 8081A	Permethrin, total	2	5	2	5	2	6							
Triazines														
EPA 547M	Glyphosate				5	2	6	2	6					
EPA 549.2M	Paraquat dichloride				5	2	6	2	6					
EPA 619	Atrazine				5	2	6	2	6					
EPA 619	Cyanazine				5	2	6	2	6					

Method	Analyte	2005		2006		2007		2008			2009			
		Storm	Irrigation	Storm	Irrigation	Storm	Irrigation	Storm	Irrigation	Fall	Winter	Storm	Irrigation	Fall
EPA 619	Simazine				5	2	6	2	6					
Metals (Total)														
EPA 200.8	Arsenic				5	2	6	2	6					
EPA 200.8	Boron				5	2	6	2	6					
EPA 200.8	Cadmium				5	2	6	2	6					
EPA 200.8	Copper				5	2	6	2	6					
EPA 200.8	Lead				5	2	6	2	6					
EPA 200.8	Nickel				5	2	6	2	6					
EPA 200.8	Selenium				5	2	2	2	6					
EPA 200.8	Zinc				5	2	6	2	6					
Metals (Dissolved)														
EPA 200.8	Cadmium													
EPA 200.8	Copper													
EPA 200.8	Lead													
EPA 200.8	Nickel													
EPA 200.8	Zinc													
Nutrients														
SM 2340 C	Hardness as CaCO3				5	2	6	2	6					
EPA 300.0	Nitrate as N				5	2	6	2	6					
EPA 350.2	Ammonia as N				5	2	6	3	6	2	2	2	6	2
EPA 351.3	Nitrogen, Total Kjeldahl				5	2	6	3	6	2	2	1	1	
EPA 354.1	Nitrite as N				5	2	6	2	6					
EPA 353.2	Nitrate + Nitrite as N							1		2	2	2	6	2
EPA 365.2	Orthophosphate as P				5	2	6	3	6	2	2	1	1	
EPA 365.2	Phosphate as P				5	2	6	3	6	2	2	2	6	2
Toxicity														
EPA 821/R-02-012	Ceriodaphnia dubia	2	5	3	5	2	8	2	6					
EPA 821/R-02-012	Pimephales promelas	2	5	2	6	2	8	2	6					

Method	Analyte	2005		2006		2007		2008			2009			
		Storm	Irrigation	Storm	Irrigation	Storm	Irrigation	Storm	Irrigation	Fall	Winter	Storm	Irrigation	Fall
EPA 821/R-02-013	Selenastrum capricornutum	2	5	2	5	2	8	4	8				2	
EPA 600/R-99-064	Hyalella azteca		5	1	1	1	3	1	2					

Table III-4. Prairie Flower Drain @ Crows Landing Rd. 2007 Management Plan additional (A) sampling schedule for chlorpyrifos and *Pimephales* toxicity. "X" indicates the site, month and analyte sampled.

Sample Site	Date	Type	Chlorpyrifos	<i>Pimephales promelas</i>
Prairie Flower @ Crows Landing Rd	31-Jul-07	A		X
Prairie Flower @ Crows Landing Rd	28-Aug-07	A	X	
Prairie Flower @ Crows Landing Rd	25-Sep-07	A	X	

Table III-5. Prairie Flower Drain site subwatershed. 2008 Management Plan upstream (U) sampling schedule for chlorpyrifos, nitrate, *Ceriodaphnia* toxicity, and *Pimephales* toxicity. "X" indicates the site, month and analyte sampled.

Sample Site	Date	Type	Chlorpyrifos	Nitrate	<i>Ceriodaphnia dubia</i>	<i>Pimephales promelas</i>
Prairie Flower @ Morgan Rd	22-Apr-08	U		X		
Prairie Flower @ Morgan Rd	20-May-08	U		X		
Prairie Flower @ Morgan Rd	17-Jun-08	U		X		
Prairie Flower @ Morgan Rd	22-Jul-08	U		X		X
Prairie Flower @ Morgan Rd	19-Aug-08	U	X	X		
Prairie Flower @ Morgan Rd	23-Sep-08	U	X	X	X	

Table III-6. Prairie Flower Drain site subwatershed. 2009 Management Plan sampling schedule for chlorpyrifos and *Selenastrum capricornutum* toxicity. "X" indicates the site, month and analyte sampled.

Site Name	Sample Date	<i>Selenastrum capricornutum</i>	Chlorpyrifos
Prairie Flower Drain @ Crows Landing Rd	21-Apr-09	X	
Prairie Flower Drain @ Crows Landing Rd	19-May-09	X	
Prairie Flower Drain @ Crows Landing Rd	18-Aug-09		X

Exceedance History

During Coalition monitoring, exceedances of WQTLs for field and physical parameters, *E. coli*, nutrients, metals, pesticides, and water column and sediment toxicity occurred in the Prairie Flower Drain site subwatershed (Table III-7). Samples collected over the entire five years of monitoring at the Prairie Flower Drain sample site resulted in exceedances including DO (19), pH (6), SC (66), TDS (45), *E. coli* (31), nitrate (23), nitrite (1), nitrate + nitrite (15), ammonia (6), chlorpyrifos (4), and one each of arsenic, dimethoate, and malathion. Toxicity has occurred twice to *Ceriodaphnia dubia*, eight times to *Selenastrum capricornutum* and twice to *Pimephales promelas*. Sediment toxicity to *Hyalella azteca* has occurred six times.

During 2009, exceedances detected at the normal monitoring and upstream MPM sites included specific conductivity (12), TDS (12), *E. coli* (6), ammonia (4), and nitrate + nitrite (12). One water sample tested toxic *Selenastrum capricornutum*.

All exceedances are listed in Table III-7 by season and date and are based on water quality trigger limits (WQTL) listed in the introduction to the ESJWQC Management Plan. The priority level (A-E) assigned to each constituent is listed in the bottom row of Table III-7 and is determined using the ESJWQC Management Plan prioritization process flow chart (Figure 4).

Table III-7. All exceedances experienced in samples collected from locations within the Prairie Flower Drain site subwatershed between February 2005 and December 2009 (sorted by season and date). If the water quality trigger limit is based on hardness then the hardness value is shown in parenthesis; otherwise the WQTL used to evaluate the data is listed in the header after the analyte.

Station Name	Season	Sample Date	Oxygen, Dissolved, 7 mg/L	pH, 6.5-8.5 units	Specific Conductivity, 700 µS/cm	Dissolved Solids, 450 mg/L	<i>E. coli</i> , 235 MPN/100 mL	Ammonia, 1.5 mg/L	Nitrate as N, 10 mg/L	Nitrite as N, 1 mg/L	Nitrate + Nitrite as N, 10 mg/L	Arsenic, 10 µg/L	Chlorpyrifos, 0.015 µg/L	Dimethoate, µ1 g/L	Malathion, 0 µg/L	<i>C. dubia</i> , Survival (%)	<i>P. promelas</i> , Survival (%)	<i>S. capricornutum</i> , Total Cell Count	<i>H. azteca</i> , Survival (%)
Prairie Flower Drain @ Crows Landing Rd	Storm	2/15/2005			2561	1600													
Prairie Flower Drain @ Crows Landing Rd	Storm	3/22/2005	6.5		2568	1600	1600												
Prairie Flower Drain @ Crows Landing Rd	Irrigation	5/11/2005			3168	1600	500												
Prairie Flower Drain @ Crows Landing Rd	Irrigation	6/15/2005			1705	1300	300												
Prairie Flower Drain @ Crows Landing Rd	Irrigation	7/13/2005	3.2		1723	1100	1600												
Prairie Flower Drain @ Crows Landing Rd	Irrigation	8/17/2005			1779	990	1600					0.029							
Prairie Flower Drain @ Crows Landing Rd	Irrigation	9/21/2005	5.22		791	460	500					0.018							83.8
Prairie Flower Drain @ Crows Landing Rd	Storm	3/1/2006			2419	1600	900												
Prairie Flower Drain @ Crows Landing Rd	Storm	3/16/2006		8.77	2728	1600	300									75			
Prairie Flower Drain @ Crows Landing Rd	Storm	3/24/2006			2782														
Prairie Flower Drain @ Crows Landing Rd	Storm	5/2/2006			2724														88.75
Prairie Flower Drain @ Crows Landing Rd	Irrigation	5/18/2006			2958	1700	550												
Prairie Flower Drain @ Crows Landing Rd	Irrigation	6/15/2006			2660	1700	1300		21										
Prairie Flower Drain @ Crows Landing Rd	Irrigation	7/13/2006	5.45	8.85	1560	720	790	18									8		
Prairie Flower Drain @ Crows Landing Rd	Irrigation	7/20/2006	6.41		1950												70		
Prairie Flower Drain @ Crows Landing Rd	Irrigation	8/10/2006			2302	1800	820		17	1.1									
Prairie Flower Drain @ Crows Landing Rd	Irrigation	9/14/2006	6.01		1276	760	2400		11										
Prairie Flower Drain @ Crows Landing Rd	Storm	2/11/2007		6.12	2659	1600	2400		24										
Prairie Flower Drain @ Crows Landing Rd	Storm	3/1/2007		8.57	2592	1500			42										

Station Name	Season	Sample Date	Oxygen, Dissolved, 7 mg/L	pH, 6.5-8.5 units	Specific Conductivity, 700 µS/cm	Dissolved Solids, 450 mg/L	<i>E. coli</i> , 235 MPN/100 mL	Ammonia, 1.5 mg/L	Nitrate as N, 10 mg/L	Nitrite as N, 1 mg/L	Nitrate + Nitrite as N, 10 mg/L	Arsenic, 10 µg/L	Chlorpyrifos, 0.015 µg/L	Dimethoate, µ1 g/L	Malathion, 0 µg/L	<i>C. dubia</i> , Survival (%)	<i>P. promelas</i> , Survival (%)	<i>S. capricornutum</i> , Total Cell Count	<i>H. azteca</i> , Survival (%)
Prairie Flower Drain @ Crows Landing Rd	Storm	3/7/2007			4798														
Prairie Flower Drain @ Crows Landing Rd	Irrigation	4/17/2007			2127	1700			25										
Prairie Flower Drain @ Crows Landing Rd	Irrigation	5/15/2007	5.59		2473	1500	920		32									1071154	
Prairie Flower Drain @ Crows Landing Rd	Irrigation	5/23/2007			2390														
Prairie Flower Drain @ Crows Landing Rd	Irrigation	6/19/2007		8.54	2304	1500			41			12							
Prairie Flower Drain @ Crows Landing Rd	Irrigation	7/17/2007	4.3		1067	730			13										
Prairie Flower Drain @ Crows Landing Rd	Irrigation	8/14/2007			1126	700	260		16										
Prairie Flower Drain @ Crows Landing Rd	Irrigation	8/16/2007			2562														58
Prairie Flower Drain @ Crows Landing Rd	Irrigation	8/28/2007	3.64		1015								0.094						
Prairie Flower Drain @ Crows Landing Rd	Irrigation	9/11/2007	7.86		1097	540	2400									0			16
Prairie Flower Drain @ Crows Landing Rd	Irrigation	9/18/2007			2262														
Prairie Flower Drain @ Crows Landing Rd	Irrigation	9/25/2007			2489														
Prairie Flower Drain @ Crows Landing Rd	Storm	1/24/2008			2371	1500	1100		23									797608	
Prairie Flower Drain @ Crows Landing Rd	Storm	1/30/2008			2944														
Prairie Flower Drain @ Crows Landing Rd	Storm	2/26/2008			2722	1600			28									442649	
Prairie Flower Drain @ Crows Landing Rd	Sediment	3/4/2008			2639														
Prairie Flower Drain @ Crows Landing Rd	Irrigation	4/22/2008			2548	1700	370		23									403571	
<i>Prairie Flower Drain @ Morgan Rd</i>	<i>Irrigation</i>	<i>4/22/2008</i>	<i>3.29</i>		<i>2574</i>				<i>35</i>										
Prairie Flower Drain @ Crows Landing Rd	Irrigation	4/29/2008	5.44		1739													517549	
Prairie Flower Drain @ Crows Landing Rd	Irrigation	5/20/2008			2526	1600	610		26									771556	
<i>Prairie Flower Drain @ Morgan Rd</i>	<i>Irrigation</i>	<i>5/20/2008</i>	<i>1.17</i>		<i>2026</i>				<i>22</i>										
Prairie Flower Drain @ Crows Landing Rd	Irrigation	5/27/2008			2273													215568	

Station Name	Season	Sample Date	Oxygen, Dissolved, 7 mg/L	pH, 6.5-8.5 units	Specific Conductivity, 700 µS/cm	Dissolved Solids, 450 mg/L	E. coli, 235 MPN/100 mL	Ammonia, 1.5 mg/L	Nitrate as N, 10 mg/L	Nitrite as N, 1 mg/L	Nitrate + Nitrite as N, 10 mg/L	Arsenic, 10 µg/L	Chlorpyrifos, 0.015 µg/L	Dimethoate, µ1 g/L	Malathion, 0 µg/L	C. dubia, Survival (%)	P. promelas, Survival (%)	S. capricornutum, Total Cell Count	H. azteca, Survival (%)
Prairie Flower Drain @ Crows Landing Rd	Irrigation	6/17/2008			2049	1200	1300	2.1	19										
Prairie Flower Drain @ Morgan Rd	Irrigation	6/17/2008			2893				30										
Prairie Flower Drain @ Crows Landing Rd	Irrigation	7/22/2008	2.51		1012	620	250		11				2.7						
Prairie Flower Drain @ Morgan Rd	Irrigation	7/22/2008	2.76		1417														
Prairie Flower Drain @ Crows Landing Rd	Irrigation	8/19/2008	4.93		956	610	440		13			0.024		0.12					
Prairie Flower Drain @ Morgan Rd	Irrigation	8/19/2008	3.63		1300				20										
Prairie Flower Drain @ Crows Landing Rd	Sediment	8/28/2008			1114														87
Prairie Flower Drain @ Crows Landing Rd	Irrigation	9/23/2008			2525	1800			33										
Prairie Flower Drain @ Morgan Rd	Irrigation	9/23/2008	3.3		2675				29										
Prairie Flower Drain @ Crows Landing Rd	Sediment	10/2/2008			2449														83
Prairie Flower Drain @ Crows Landing Rd	Fall	10/21/2008			1742	1100	370				27								
Prairie Flower Drain @ Crows Landing Rd	Fall	11/11/2008			2151	1500					39								
Prairie Flower Drain @ Crows Landing Rd	Fall	12/16/2008			2298	2900	1300				40								
Prairie Flower Drain @ Crows Landing Rd	Winter	1/20/2009			2414	1500					43								
Prairie Flower Drain @ Crows Landing Rd	Storm	2/7/2009			2255	1300					31								
Prairie Flower Drain @ Crows Landing Rd	Winter	3/17/2009		8.74	2394	1400					34								
Prairie Flower Drain @ Crows Landing Rd	Irrigation	4/21/2009			2223	1400	410				24								
Prairie Flower Drain @ Crows Landing Rd	Irrigation	5/19/2009	4.78		2066	1200	>2400	3.2			20							266798	
Prairie Flower Drain @ Crows Landing Rd	Irrigation	6/16/2009			2417	1400		1.3			22								
Prairie Flower Drain @ Crows Landing Rd	Irrigation	7/21/2009			1366	820		1.8			14								
Prairie Flower Drain @ Crows Landing Rd	Irrigation	8/18/2009			1984	1200					22								
Prairie Flower Drain @ Crows Landing Rd	Irrigation	9/22/2009			2171	1400	1300				35								

Station Name	Season	Sample Date	Oxygen, Dissolved, 7 mg/L	pH, 6.5-8.5 units	Specific Conductivity, 700 µS/cm	Dissolved Solids, 450 mg/L	<i>E. coli</i> , 235 MPN/100 mL	Ammonia, 1.5 mg/L	Nitrate as N, 10 mg/L	Nitrite as N, 1 mg/L	Nitrate + Nitrite as N, 10 mg/L	Arsenic, 10 µg/L	Chlorpyrifos, 0.015 µg/L	Dimethoate, µ1 g/L	Malathion, 0 µg/L	<i>C. dubia</i> , Survival (%)	<i>P. promelas</i> , Survival (%)	<i>S. capricornutum</i> , Total Cell Count	<i>H. azteca</i> , Survival (%)	
Prairie Flower Drain @ Crows Landing Rd	Fall	10/20/2009			2459	1400	1300				25									
Prairie Flower Drain @ Crows Landing Rd	Fall	11/17/2009			2415	1500	>2400	8.8			36									
Prairie Flower Drain @ Crows Landing Rd	Storm	12/15/2009			2695	1600	2000				36									
Constituent Priority			E	E	E	E	E	E	D	NP	D	NP	A/B	NP	NP	D	NP¹	E	D	

NP- Not prioritized. Fewer than two exceedances for this constituent at this site within three years and currently no TMDL for constituent
NP¹ – Not prioritized; both toxic samples were from the same sampling event (sample and resample to test for persistence).

2007 - 2009 Management Plan Monitoring Results

Management Plan Monitoring results are included in Table III-8 for chlorpyrifos, nitrate, and toxicity to *Ceriodaphnia dubia* and *Pimephales promelas* from 2007 through 2009. The 2009 MPM results also include *Selenastrum capricornutum*.

2007

In 2007, MPM was implemented at the Prairie Flower Drain monitoring site for chlorpyrifos and *Pimephales promelas* toxicity. The site was monitored monthly for chlorpyrifos and *Pimephales promelas* toxicity. Additional MPM occurred for *Pimephales* toxicity in July and for chlorpyrifos in August and September of 2007 (Table III-8). In August, the normal monitoring sample did not contain chlorpyrifos above the detection limit however the additional sample (collected two weeks later) contained 0.094µg/L of chlorpyrifos. No toxicity to *Pimephales* occurred in the samples collected in 2007.

2008

Upstream MPM for chlorpyrifos, nitrate, and toxicity to *Ceriodaphnia dubia* and *Pimephales promelas* occurred during the 2008 irrigation season at Prairie Flower Drain @ Morgan Road. Table III-8 provides monitoring results for chlorpyrifos, nitrate, *Ceriodaphnia*, and *Pimephales* toxicity from all sampling events during the 2008 irrigation season. Chlorpyrifos exceeded the WQTL once on August 19, 2008 during normal monitoring (0.024 µg/L). With the exception of July upstream MPM, nitrate exceedances were experienced during every sampling event in 2008 at both monitoring locations. No samples were toxic to *Ceriodaphnia* or *Pimephales* during the 2008 irrigation season.

2009

Management Plan Monitoring at Prairie Flower Drain @ Crows Landing Rd for chlorpyrifos resulted in no detections in 2009. Monitoring for *Selenastrum* toxicity in April and May resulted in toxic samples in May. A TIE was initiated on May 27, 2009 and it was concluded that cationic chemical(s) and ammonia were reported as the cause of toxicity to *Selenastrum*.

Management Plan Monitoring for chlorpyrifos, *Ceriodaphnia* toxicity, *Selenastrum* toxicity and *Hyalella* toxicity are planned for the 2010 irrigation season. Management Plan Monitoring during February and March 2011 will also occur for *Ceriodaphnia* and *Selenastrum* toxicity.

Table III-8. Prairie Flower Drain site subwatershed. Normal monitoring (NM) and Management Plan Monitoring (MPM) results where 'A' indicates additional MPM (2007), 'US' indicates upstream MPM for chlorpyrifos (2008-2009), nitrate, *Ceriodaphnia*, and *Pimephales* from the 2007-2008 irrigation seasons, and MPM for *Selenastrum* (2009). Exceedance values are in bold.

Month:		April	May	June	July	August	September
2007 NM (@ Crows Landing Rd)	Date	4/17/07	5/15/07	6/19/07	7/17/07	8/14/07	9/25/07
	Chlorpyrifos (µg/L)	<0.003	<0.003	<0.003	0.009	<0.003	<0.003
	Nitrate (mg/L)	25	41	32	13	16	9.8
	<i>C. dubia</i> toxicity (% Control)	100	90	95	100	100	0
	<i>P. promelas</i> toxicity (% Control)	97.5	100	100	97.5	100	100
2007 MPM A (@ Crows Landing Rd)	Date	NA	NA	NA	7/31/07	8/28/07	9/25/07
	Chlorpyrifos (µg/L)	NA	NA	NA	NA	0.094	<0.003
	<i>P. promelas</i> toxicity (% Control)	NA	NA	NA	100	NA	NA
2008 NM (@ Crows Landing Rd)	Date	4/22/08	5/20/08	6/17/08	7/22/08	8/19/08	9/23/08
	Chlorpyrifos (µg/L)	<0.003	<0.003	<0.003	<0.003	0.024	<0.003
	Nitrate (mg/L)	23	26	19	11	13	33
	<i>C. dubia</i> toxicity (% Control)	100	95	100	95	100	95
	<i>P. promelas</i> toxicity (% Control)	100	100	100	100	100	100
2008 MPM US (@ Morgan Rd)	Date	4/22/08	5/20/08	6/17/08	7/22/08	8/19/08	9/23/08
	Chlorpyrifos (µg/L)	NA	NA	NA	NA	<0.003	<0.003
	Nitrate (mg/L)	35	22	30	0.053	20	29
	<i>C. dubia</i> toxicity (% Control)	NA	NA	NA	NA	NA	100
	<i>P. promelas</i> toxicity (% Control)	NA	NA	NA	100	NA	NA
2009 MPM (@ Crows Landing Rd)	Date	4/21/09	5/19/09	NA	NA	8/18/09	NA
	Chlorpyrifos (µg/L)	<0.003	<0.003	NA	NA	<0.003	NA
	<i>S. capricornutum</i> toxicity (% Control)	248	30	NA	NA	NA	NA

NA - Not applicable. This site was not sampled during this month.

Load Calculations

Loads have been calculated for all chlorpyrifos detections in the site subwatershed (Table III-9) based on the following formula:

$$\text{Load} = \text{Discharge (cfs)} \times 28.317\text{L/ft}^3 \times \text{Concentration (milligram/L} \times 1000 \text{ or } \mu\text{g/L)}.$$

The load values for constituents in this report represent instantaneous loads only. These values should not be used to extrapolate loading over any period of time (e.g. weekly, monthly, seasonal or annual). The primary purpose for reporting instantaneous loads is to provide a normalization of the concentrations by flow for various constituents at the time the samples were collected.

Table III-9. Prairie Flower Drain site subwatershed. Instantaneous load calculations for chlorpyrifos where discharge was measured (sorted by analyte, site and date).

Station Name	Analyte Name	Sample Date	Discharge cfs	Concentration µg/L	Loading Rate µg/sec
Prairie Flower Drain @ Crows Landing Rd	Chlorpyrifos	17-Aug-05	0	0.029	0
Prairie Flower Drain @ Crows Landing Rd	Chlorpyrifos	21-Sep-05	0	0.018	0

Station Name	Analyte Name	Sample Date	Discharge cfs	Concentration µg/L	Loading Rate µg/sec
Prairie Flower Drain @ Crows Landing Rd*	Chlorpyrifos	21-Sep-05	0	0.018	0
Prairie Flower Drain @ Crows Landing Rd	Chlorpyrifos	13-Jul-06	0	0.014	0
Prairie Flower Drain @ Crows Landing Rd	Chlorpyrifos	17-Jul-07	2.32	0.009	0.591259
Prairie Flower Drain @ Crows Landing Rd	Chlorpyrifos	19-Aug-08	4.67	0.024	3.173769

*Field Duplicate

Source Identification and Outreach

Priority A/B Constituents

Chlorpyrifos is the only priority A/B constituent under the Prairie Flower Management Plan.

Chlorpyrifos

The WQTL for chlorpyrifos was exceeded four times at this site, once during the month of September (2005) and three times during the month of August (2005, 2007, 2008, Table III-7). There were no exceedances during 2009 MPM.

The number of chlorpyrifos applications, amount of chlorpyrifos applied, and acres treated decreased dramatically in 2009 when compared to 2005 and 2008 levels, in particular for the month of July during which most applications occur (Table III-10 and Figure III-2). The greatest use of chlorpyrifos occurred in 2008 (740 lbs) with the lowest use was in 2009 (239 lbs, Table III-10). Exceedances do not always occur in months with the highest applications (Figure III-2).

Only two crops receive applications of chlorpyrifos in the subwatershed, alfalfa and corn, with corn receiving the most pounds of active ingredient applied (1,600 lbs) and alfalfa receiving (475 lbs, Table III-11).

Table III-10. Number of chlorpyrifos applications, total pounds applied and total acres treated by month for January 2005 through December 2009 in the Prairie Flower Drain @ Crows Landing Road site subwatershed. If a month is not included in the table, no applications were made.

Month/Year*	Number of Chlorpyrifos Applications	Pounds of AI Applied	Acres Treated
March, 2005	1	61.0	120
July, 2005	5	256.6	312
August, 2005	8	415.7	485
March, 2006	1	19.9	40
July, 2006	4	194.2	191
August, 2006	3	147.4	207
May, 2008	2	212.1	140
July, 2008	6	437.8	561
August, 2008	4	90.5	89
March, 2009	1	76.3	150

Month/Year*	Number of Chlorpyrifos Applications	Pounds of AI Applied	Acres Treated
July, 2009	2	142.3	140
August, 2009	1	20.3	20
Summaries by Year			
2005 Total	14	733.3	917
2006 Total	8	361.5	438
2008 Total	12	740.4	790
2009 Total	4	238.9	310
Total	38	2,074.2	2,455

*No applications of chlorpyrifos reported for 2007.

Figure III-2. Pounds of chlorpyrifos added within the Prairie Flower Drain @ Crows Landing Rd site subwatershed by month for 2005-2009. Asterisk (*) denotes months with exceedances. No use of chlorpyrifos was reported for 2007.




Table III-11. Total pounds AI for chlorpyrifos based on PUR data from 2005-2009 within the Prairie Flower Drain @ Crows Landing Rd subwatershed.

Chemical Name	Commodity	Product Name	Lbs AI Applied
CHLORPYRIFOS*	ALFALFA	LOCK-ON INSECTICIDE	128.689
		NUFOS 4E	76.257
	ALFALFA (FORAGE - FODDER) (ALFALFA HAY)	LOCK-ON INSECTICIDE	188.778
		LORSBAN 4E-HF	19.939
		NUFOS 4E	61.006
	CORN FOR/FOD	GOVERN 4E INSECTICIDE	212.107
		NUFOS 4E	562.271
		CORN (FORAGE - FODDER)	CHLORPYRIFOS 4E AG
	LORSBAN-4E		46.014
	NUFOS 4E		685.300
ALFALFA – Total Pounds Chlorpyrifos Applied			474.669
CORN – Total Pounds Chlorpyrifos Applied			1599.508
Total pounds chlorpyrifos applied (2005 - 2009)			2,074.177

*No applications of chlorpyrifos reported for 2007.

In the 2009 Management Plan Update, the Coalition determined the most probable sources of chlorpyrifos in the Prairie Flower Drain were runoff from row crops recently receiving chlorpyrifos applications. Unfortunately, of the areas associated with exceedances in 2008, Coalition members own only a small portion of the acreage (2009 Management Plan Update). A large amount of the acreage is owned and farmed by dairies and it is assumed that the land not covered by the Coalition is part of the dairy program. This pattern of ownership suggests that it may be difficult to improve water quality in this site subwatershed. It is difficult to receive cooperation from nonmembers as well as track implementation of management practices on nonmember land.

The results from the general surveys administered to all members in the subwatershed are summarized in the ESJWQC General Survey Summary Report submitted January 30, 2009. The results were used as a starting point to evaluate management practice implementation within the subwatershed. The survey assessed practices used in the period from May 17, 2007 to December 2009. The Coalition noted that irrigation water is actively managed by members in this subwatershed. Growers accounting for a combined 48% of acreage indicated either use of recirculation tailwater return systems or restriction of water flow. However, growers accounting for 35% of the acreage represented in the response indicated that they used no practices to reduce runoff. Using this information, the Coalition designed outreach to affect an increase in relevant and effective management practice implementation in member parcels.

Coalition outreach since 2007 has included grower meetings and the mailing/distribution of information. A complete list of Coalition Outreach during 2009 is provided in the Summary of Coalition Outreach Activities section of this report. Prior to the 2009 irrigation season, several group grower meetings were held for members in the Prairie Flower Drain subwatershed to make them aware of the current issues and concerns specific to pesticide applications in the upcoming irrigation season and relevant management practices to protect and improve water quality in the waterway. The meetings also allowed the Coalition to update members on the strategy and status of the Prairie Flower Drain Management Plan.

The Coalition then turned its attention to individual contacts. Targeted growers were selected based on their crop type, pesticide usage, and potential to drain to the creek. Of the 3,106 acres in Prairie Flower Drain subwatershed that have the potential to drain directly to the creek, 900 acres are owned and operated by Coalition members. The Coalition produced a final list of 11 target members representing 865 acres in the subwatershed. Individual meetings occurred starting in June, 2009 and continued through September, 2009. Coalition representatives met with growers on their property to discuss the Prairie Flower Drain Management Plan and offer management practice recommendations specific to the land owner. This also gave the Coalition a chance to better assess currently implemented management practices as well as future planned management practice implementation. Individual contacts focused mainly on chlorpyrifos exceedances however all water quality results were reviewed and discussed including nitrate, ammonia, *E. coli*, DO, SC, TDS, and pH exceedances and toxicity to *Ceriodaphnia*, *Selenastrum*, and *Hyaella*. To date, the Coalition has contacted all its targeted members, which encompasses 865 acres or 28% of the total subwatershed acreage. The Coalition plans to contact target members in the spring of 2010 to determine if owner's planned future management practices have been implemented, and will continue to conduct outreach on a grower group level and monitor the creek to assess water quality.

Priority D Constituents

Prairie Flower Drain is listed for nitrate / nitrate + nitrite, sediment toxicity to *Hyaella azteca* and water toxicity *Ceriodaphnia dubia* as priority D constituents. The Coalition replaced separate analysis of nitrate and nitrite with the combined analysis of nitrate + nitrite starting in October 2008. Exceedances of the nitrate WQTL, either of separately measured nitrate or of combined nitrate + nitrite analysis, have occurred during every sample event.

Nitrate + Nitrite / nitrate

There were 12 additional exceedances of the nitrate WQTL in the Prairie Flower Drain site subwatershed in 2009. Sources of nitrate are discussed in the introduction of the ESJWQC Management Plan submitted in September, 2006 and include fertilizer

applications and dairy manure waste. Prairie Flower Drain subwatershed contains approximately 443 acres of confined animal operations most of which are dairies. In addition, nitrate contamination of ground water is common in the region, and Prairie Flower Drain intercepts shallow ground water. It is not clear if the nitrates are originating with recent applications of dairy waste that may be reaching surface waters, or from past contamination of ground water which is being intercepted by the drain. The Coalition does not have the capabilities to track fertilizer applications and therefore cannot determine the exact source of the nitrate exceedances. In 2008, exceedances occurred during every sampling event and during upstream sampling at Morgan Rd in all months except July. Exceedances prior to 2008 occurred in most of the months of the year including February, March, May, June, July, August, and September. It is unlikely that applications of manure would occur in every month, indicating that ground water and/or dairy discharge are most likely both sources.

***Ceriodaphnia* toxicity**

Toxicity to *Ceriodaphnia* was not monitored during 2009. *Ceriodaphnia* toxicity occurred once during March 2006 and once during September 2007. The 2007 toxicity was associated with an exceedance of chlorpyrifos in August 2007. Due to this correlation, the Coalition's strategy for eliminating *Ceriodaphnia* toxicity will involve focusing on chlorpyrifos. If chlorpyrifos can be prevented from entering waterways throughout the irrigation season, the Coalition believes that *Ceriodaphnia* toxicity can also be reduced or eliminated.

***Hyalella* toxicity**

Sediment toxicity to *Hyalella* has occurred during September 2005 and May 2006. It occurred most recently on August 28, 2008 and also in the resample collected on October 2, 2008. Both samples had survival greater than 80% compared to the control. Toxicity to *Hyalella* was not monitored during 2009.

The Coalition conducted management plan sampling for sediment toxicities. As of 2009, the Coalition now tests for total organic carbon and grain size in all sediments and pyrethroids and chlorpyrifos in sediment that exhibits toxicity. Management Plan Monitoring is scheduled for this site in 2010.

Priority E Constituents

The following priority E constituents are listed under the Prairie Flower Drain management plan: ammonia, DO, pH, SC, dissolved solids, *E. coli*, and water column toxicity to *Selenastrum capricornutum*. Exceedances of ammonia, DO, pH, SC, TDS, and *E. coli* continued in 2009. Toxicity to *Selenastrum* has occurred several times at the site, usually in the winter or early irrigation months, but is never associated with an herbicide or metal exceedance. Toxicity to *Selenastrum* occurred once during May of 2009 MPM,

and was again not associated with an herbicide or metal exceedance. These constituents will remain low priority but will be discussed during individual contacts and annual grower meetings.

Evaluation

The Prairie Flower Drain subwatershed is one of the first three priority site subwatersheds within the ESJWQC and therefore the Coalition has focused its resources on identifying the sources of agricultural discharge within this subwatershed that could result in water quality impairments, extending outreach to individual Coalition members, and setting evaluation goals.

The Coalition's strategy for the Prairie Flower Drain subwatershed has been to target growers who farm row crops and have the potential to discharge. Outreach includes grower notification, management practice education, tracking of management practices implementation, and providing information on special studies of management practice efficacy. The Coalition believes that because land use in this subwatershed is predominately row crops, outreach should continue to focus on management practices that restrict runoff. Growers who farm alfalfa and corn should be targeted.

Individual surveys to document current management practices and assess planned implementations have been completed representing 100% of targeted growers. The Coalition is now in the process of evaluating these results and planning the final stages of outreach.

Acres treated with and pounds applied of chlorpyrifos decreased overall since 2005. A decrease was seen between 2008 usage and 2009 usage, which may further demonstrate that recent Coalition outreach has had an impact. Additionally, usage decreased in July and August, when the most exceedances have occurred. Also encouraging was the absence of chlorpyrifos exceedances and detections during 2009 Management Plan sampling. Management Plan Monitoring in 2010 will indicate if a change in management practices as a result of individual contacts results in long-term improvements to water quality.

IV. BEAR CREEK @ KIBBY RD

Management Plan Constituents

Priority A/B

- Chlorpyrifos

Priority C

- Copper

Priority D

- *Ceriodaphnia dubia* water column toxicity

Priority E

- Dissolved Oxygen
- pH
- *E. coli*



Description of Bear Creek @ Kibby Rd Site Subwatershed

Bear Creek @ Kibby Road (7, 298 irrigated acres) – This site subwatershed drains an eastern portion of the Coalition region in Merced County. Bear Creek originates in the foothills of the Sierras with Burn’s Creek as one of the major tributaries. Bear Creek drains to the east just north of the towns of Planada, through Merced and eventually to the San Joaquin River. The primary irrigated agriculture in the site subwatershed includes deciduous nuts, field crops, truck crops, and irrigated pasture (Figure IV-1). Table IV-1 includes the station name, station code and target latitude/longitudes for the site sampled within this subwatershed.

Table IV-1. Coordinates of the Bear Creek site subwatershed sampling location.

Station Name	Station Code	Target Latitude	Target Longitude
Bear Creek @ Kibby Rd	535XBCAKR	37.3128	-120.4138

Figure IV-1. Site subwatershed map of land use for the Bear Creek @ Kibby Rd sample site.



Bear Creek @ Kibby Rd

Date Prepared: 02/04/10
 ESJWQC

Subwatershed Monitoring History

Ambient water monitoring conducted at Bear Creek @ Kibby Road was initiated during the storm season of 2005 continuing through fall season of 2008 (Table IV-2). Starting in October 2008, Bear Creek @ Kibby Rd was removed from the monitoring schedule and became an Assessment site under the current MRPP. It is scheduled for Assessment monitoring in 2025-2026. The constituents sampled at this location from 2005-2008 are listed in Table IV-3.

Management Plan Monitoring for the Coalition at Bear Creek @ Kibby Rd occurred in May and July of 2008 and included additional sampling for chlorpyrifos and *Ceriodaphnia dubia* toxicity (Table IV-4). The Coalition will conduct MPM for chlorpyrifos and *Ceriodaphnia dubia* toxicity again starting May 2010. There were no samples collected from this site during 2009. Included in this section is a summary and of exceedances experienced at this site (Table IV-5).

Bear Creek is scheduled as a high priority site starting in 2010. Bear Creek is listed as an impaired waterbody for *E. coli* and unknown toxicity in the proposed 2008 Central Valley Basin Plan 303d list.

Table IV-2. Bear Creek @ Kibby Rd sampling events per season and year.

	2004	2005		2006		2007		2008			2009			
	Irrigation	Storm	Irrigation	Storm	Irrigation	Storm	Irrigation	Storm	Irrigation	Fall	Winter	Storm	Irrigation	Fall
Events Sampled	NA	1	5	2	5	2	6	NA	6	NA	NA	NA	NA	NA
Events Not Sampled	NA	1	0	0	0	0	0	NA	0	NA	NA	NA	NA	NA
Total	NA	2	5	2	5	2	6	NA	6	NA	NA	NA	NA	NA

NA indicates that this site was not sampled during this season/year.

Table IV-3. Number of analyses performed per analyte in each sampling season and year for the Bear Creek @ Kibby Rd sample site. Only environmental samples with a sample replicate and lab replicate number of one are shown.

Method	Analyte	2005		2006		2007		2008		
		Storm	Irrigation	Storm	Irrigation	Storm	Irrigation	Storm	Irrigation	Fall*
Field and Physical Parameters										
EPA 110.2	Color	1	5	2	5	2	6	2	6	
EPA 160.1	Dissolved Solids	1	5	2	5	2	6	2	6	
EPA 180.1	Turbidity	1	5	2	5	2	6	2	6	
EPA 405.1	BOD				1	2	2			
EPA 415.1	Total Organic Carbon	1	5	2	5	2	6	2	6	
SM 9223	E. coli	1	5	2	5	2	6	2	6	
NA	Dissolved Oxygen	1	6	3	6	3	7	3	11	
NA	Specific Conductivity	1	6	3	6	3	7	3	11	
NA	pH	1	6	3	6	3	7	3	11	
Carbamates										
EPA 8321A	Aldicarb				5	2	6	2	6	
EPA 8321A	Carbaryl				5	2	6	2	6	
EPA 8321A	Carbofuran				5	2	6	2	6	
EPA 8321A	Diuron				5	2	6	2	6	
EPA 8321A	Linuron				5	2	6	2	6	
EPA 8321A	Methiocarb				5	2	6	2	6	
EPA 8321A	Methomyl				5	2	6	2	6	
EPA 8321A	Oxamyl				5	2	6	2	6	
Organochlorines										
EPA 8081A	DDD (p,p')				5	2	6	2	6	
EPA 8081A	DDE (p,p')				5	2	6	2	6	
EPA 8081A	DDT (p,p')				5	2	6	2	6	
EPA 8081A	Dicofol				5	2	6	2	6	
EPA 8081A	Dieldrin				5	2	6	2	6	
EPA 8081A	Endrin				5	2	6	2	6	

Method	Analyte	2005		2006		2007		2008		
		Storm	Irrigation	Storm	Irrigation	Storm	Irrigation	Storm	Irrigation	Fall*
EPA 8081A	Methoxychlor				5	2	6	2	6	
Organophosphates										
EPA 8141A	Azinphos methyl				5	2	6	2	6	
EPA 8141A	Chlorpyrifos	1	5	2	5	2	6	2	8	
EPA 8141A	Diazinon				5	2	6	2	6	
EPA 8141A	Dimethoate				5	2	6	2	6	
EPA 8141A	Disulfoton				5	2	6	2	6	
EPA 8141A	Malathion				5	2	6	2	6	
EPA 8141A	Methamidophos				5	2	6	2	6	
EPA 8141A	Methidathion				5	2	6	2	6	
EPA 8141A	Molinate				5	2	6	2	6	
EPA 8141A	Parathion, Methyl				5	2	6	2	6	
EPA 8141A	Phorate				5	2	6	2	6	
EPA 8141A	Phosmet				5	2	6	2	6	
EPA 8141A	Thiobencarb				5	2	6	2	6	
Pyrethroids										
EPA 8081A	Bifenthrin		1	2	5	2	6			
EPA 8081A	Cyfluthrin, total		1	2	5	2	6			
EPA 8081A	Cyhalothrin, lambda, total	1	5	2	5	2	6			
EPA 8081A	Cypermethrin, total	1	5	2	5	2	6			
EPA 8081A	Esfenvalerate/ Fenvalerate, total	1	5	2	5	2	6			
EPA 8081A	Permethrin, total	1	5	2	5	2	6			
Triazines										
EPA 547M	Glyphosate				5	2	6	2	6	
EPA 549.2M	Paraquat dichloride				5	2	6	2	6	
EPA 619	Atrazine				5	2	6	2	6	
EPA 619	Cyanazine				5	2	6	2	6	

Method	Analyte	2005		2006		2007		2008		
		Storm	Irrigation	Storm	Irrigation	Storm	Irrigation	Storm	Irrigation	Fall*
EPA 619	Simazine				5	2	6	2	6	
Metals (Total)										
EPA 200.8	Arsenic				5	2	6	2	6	
EPA 200.8	Boron				5	2	6	2	6	
EPA 200.8	Cadmium				5	2	6	2	6	
EPA 200.8	Copper				5	2	6	2	6	
EPA 200.8	Lead				5	2	6	2	6	
EPA 200.8	Nickel				5	2	6	2	6	
EPA 200.8	Selenium				5	2	2	2	6	
EPA 200.8	Zinc				5	2	6	2	6	
Nutrients										
SM 2340 C	Hardness as CaCO3				5	2	6	2	6	
EPA 300.0	Nitrate as N				5	2	6	2	6	
EPA 350.2	Ammonia as N				5	2	6	2	6	
EPA 351.3	Nitrogen, Total Kjeldahl				5	2	6	2	6	
EPA 354.1	Nitrite as N				5	2	6	2	6	
EPA 353.2	Nitrate + Nitrite as N									
EPA 365.2	Orthophosphate as P				5	2	6	2	6	
EPA 365.2	Phosphate as P				5	2	6	2	6	
Toxicity										
EPA 821/R-02-012	Ceriodaphnia dubia	1	6	2	6	2	8	2	8	
EPA 821/R-02-012	Pimephales promelas	1	6	2	5	2	7	2	6	
EPA 821/R-02-013	Selenastrum capricornutum	1	6	2	5	2	7		7	
EPA 600/R-99-064	Hyaella azteca		5	1	1	1	1	1	2	

Table IV-4. Bear Creek @ Kibby Rd site subwatershed. 2008 Management Plan additional (A) sampling schedule for chlorpyrifos and *Ceriodaphnia*. "X" indicates the site, month, and analyte sampled.

Sample Site	Month	Type	Chlorpyrifos	<i>Ceriodaphnia dubia</i>
Bear Creek @ Kibby Rd	7-May-08	A	X	X
Bear Creek @ Kibby Rd	8-Jul-08	A	X	X

Exceedance History

During Coalition monitoring, exceedances of WQTLs for field and physical parameters, *E. coli*, nutrients, metals, pesticides, and water column and sediment toxicity occurred within the Bear Creek site subwatershed (Table IV-5). During ambient water monitoring several exceedances occurred, including DO 92), pH (3), *E. coli* (7), arsenic (1), copper (4), chlorpyrifos (2), and DDT (1). Water column toxicity occurred three times to *Ceriodaphnia dubia* and twice to *Selenastrum capricornutum* including a resample to initial toxicity. Sediment toxicity to *Hyalella azteca* has occurred twice.

All exceedances are listed in Table IV-5 by season and date and are based on WQTL listed in the introduction to the ESJWQC Management Plan. Using the ESJWQC Management Plan prioritization process flow chart (Figure 4), priorities were assigned to management plan constituents and are listed in the bottom row of Table IV-5. The highest priority constituents in the Bear Creek subwatershed are chlorpyrifos (A), copper (C), and toxicity to *Ceriodaphnia* and *Hyalella* (D).

Table IV-5. All exceedances experienced in samples collected from locations within the Bear Creek site subwatershed between March 2005 and October 2008 (sorted by season and date). If the water quality trigger limit is based on hardness then the hardness value is shown in parenthesis; otherwise the WQTL used to evaluate the data is listed in the header after the analyte.

Station Name	Season	Sample Date	Oxygen, Dissolved, 7 mg/L	pH, 6.5-8.5 units	<i>E. coli</i> , 235 MPN/100 mL	Arsenic, 10 µg/L	Copper, Total, 1,300 µg/L	Chlorpyrifos, 0.015 µg/L	DDT (p,p'), 0.00059 µg/L	<i>C. dubia</i> , Survival (%)	<i>H. azteca</i> , Survival (%)	<i>S. capricornutum</i> , Total Cell Count
Bear Creek @ Kibby Rd	Storm	3/21/2005	4.4		1600							
Bear Creek @ Kibby Rd	Irrigation	5/10/2005			280					5		
Bear Creek @ Kibby Rd	Storm	3/15/2006			1600							
Bear Creek @ Kibby Rd	Irrigation	5/17/2006						0.52		0		
Bear Creek @ Kibby Rd	Irrigation	6/13/2006	6.99	8.69								
Bear Creek @ Kibby Rd	Storm	2/12/2007			2400		12 (9.3)		0.0091			
Bear Creek @ Kibby Rd	Storm	3/1/2007			1300							
Bear Creek @ Kibby Rd	Irrigation	7/24/2007						0.049		0		
Bear Creek @ Kibby Rd	Irrigation	8/21/2007		8.69								
Bear Creek @ Kibby Rd	Storm	1/24/2008			2400		8.6 (7.7)					
Bear Creek @ Kibby Rd	Storm	2/25/2008			>2400		7.2 (6.4)					
Bear Creek @ Kibby Rd	Sediment	3/4/2008		8.72								
Bear Creek @ Kibby Rd	Irrigation	4/29/2008										42100
Bear Creek @ Kibby Rd	Irrigation	5/7/2008										735734
Bear Creek @ Kibby Rd	Irrigation	6/24/2008				17						

Station Name	Season	Sample Date	Oxygen, Dissolved, 7 mg/L	pH, 6.5-8.5 units	<i>E. coli</i> , 235 MPN/100 mL	Arsenic, 10 µg/L	Copper, Total, 1,300 µg/L	Chlorpyrifos, 0.015 µg/L	DDT (p,p'), 0.00059 µg/L	<i>C. dubia</i> , Survival (%)	<i>H. azteca</i> , Survival (%)	<i>S. capricornutum</i> , Total Cell Count
Bear Creek @ Kibby Rd	Irrigation	7/29/2008										
Bear Creek @ Kibby Rd	Irrigation	8/26/2008					7.1 (2.4)					
Bear Creek @ Kibby Rd	Sediment	8/28/2008									90	
Bear Creek @ Kibby Rd	Sediment	10/2/2008									88	
Constituent Priority			E	E	E	NP	C	A/B	NP	D	NP¹	NP¹

NP- Not prioritized. Fewer than two exceedances for this constituent at this site within three years and currently no TMDL for constituent
NP¹ – Not prioritized; both toxic samples were from the same sampling event (sample and resample to test for persistence).

2008 Management Plan Monitoring Results

Management Plan Monitoring results are included in Table IV-6 for chlorpyrifos and toxicity to *Ceriodaphnia* for 2008; MPM did not occur for either during 2009. Toxicity to *Ceriodaphnia* occurred in May of 2005 and 2006, and in July of 2007. The 2006 and 2007 toxicities both coincided with an exceedance of chlorpyrifos (Table IV-5).

2008

In 2008, Bear Creek @ Kibby Rd was monitored monthly for chlorpyrifos and *Ceriodaphnia dubia* toxicity from April through September plus additional monitoring conducted in May and July (Table IV-6). Both normal monitoring and MPM for chlorpyrifos indicated no detections for any of the irrigation months in 2008, and there were no toxic samples to *Ceriodaphnia*. Copper concentrations exceeded the WQTL based on hardness three times during 2008 normal monitoring (January, February, and August) and was added to the Bear Creek Management Plan.

2009

Bear Creek @ Kibby Rd was not scheduled for 2009 MPM.

Management Plan Monitoring for chlorpyrifos, copper, and *Ceriodaphnia dubia* toxicity will be conducted during the 2010 irrigation season. Copper is also scheduled for MPM during January and February 2011.

Table IV-6. Bear Creek @ Kibby Rd site subwatershed. Normal monitoring (NM) and Management Plan Monitoring (MPM) results for chlorpyrifos and *Ceriodaphnia* from the 2008 irrigation season. Exceedance values are in bold.

Month:		April	May	June	July	August	September
2008 NM (@ Kibby Rd)	Date	4/22/08	5/20/08	6/17/08	7/22/08	8/19/08	9/23/08
	Chlorpyrifos (µg/L)	<0.003	<0.003	<0.003	<0.003	<0.003	<0.003
	<i>C. dubia</i> toxicity (% Control)	100	100	100	100	100	100
2008 MPM (@ Kibby Rd)	Date	NA	5/07/08	NA	7/08/08	NA	NA
	Chlorpyrifos (µg/L)	NA	<0.003	NA	<0.003	NA	NA
	<i>C. dubia</i> toxicity (% Control)	NA	100	NA	95	NA	NA

NA - Not applicable. This site was not sampled during this month.

Load Calculations

Loads have been calculated for all chlorpyrifos detections in the site subwatershed (Table IV-7) based on the following formula:

$$\text{Load} = \text{Discharge (cfs)} \times 28.317\text{L/ft}^3 \times \text{Concentration (milligram/L} \times 1000 \text{ or } \mu\text{g/L)}.$$

The load values for constituents in this report represent instantaneous loads only. These values should not be used to extrapolate loading over any period of time (e.g. weekly, monthly, seasonal or annual). The primary purpose for reporting instantaneous loads is to provide a normalization of the concentrations by flow for various constituents at the time the samples were collected.

Table IV-7. Bear Creek site subwatershed. Instantaneous load calculations for chlorpyrifos and copper where discharge was measured (sorted by analyte, site and date).

Station Name	Analyte Name	Sample Date	Discharge cfs	Concentration µg/L	Loading Rate µg/sec
Bear Creek @ Kibby Rd*	Copper	17-Apr-07	29.13	2	1649.748
Bear Creek @ Kibby Rd	Copper	17-Apr-07	29.13	2.1	1732.236
Bear Creek @ Kibby Rd	Copper	21-Aug-07	56.1	1.6	2541.734
Bear Creek @ Kibby Rd	Copper	18-Sep-07	51.02	1.7	2456.047
Bear Creek @ Kibby Rd	Copper	29-Apr-08	113.34	1.1	3530.394
Bear Creek @ Kibby Rd	Copper	27-May-08	147.77	1.4	5858.164
Bear Creek @ Kibby Rd	Copper	26-Aug-08	94.9	7.1	19079.71
Bear Creek @ Kibby Rd	Copper	30-Sep-08	33.1	1.3	1218.481

*Field Duplicate

Source Identification and Outreach

Priority A/B Constituents

Chlorpyrifos is the only priority A/B constituent listed under the Bear Creek Management Plan.

Chlorpyrifos

Chlorpyrifos has exceeded the WQTL of 0.015µg/L two times in this subwatershed from 2006 through 2008 (May, 2006 and July, 2007). Chlorpyrifos was omitted from the monitoring schedule in October 2008 when the Coalition began monitoring under a new MRPP. Starting in May 2010, chlorpyrifos will be monitored as a part of MPM. The Coalition uses a combination of monitoring data and evaluation of Pesticide Use Report (PUR) data to identify possible sources.

PUR data are reviewed for the number of monthly chlorpyrifos applications, pounds active ingredient (AI) applied, and acres treated (Table IV-8, Figure IV-2). The amount of chlorpyrifos applied within the subwatershed decreased in 2009 significantly (328 lbs), however the amount applied in 2008 was the greatest amount of chlorpyrifos use (1,817 lb) (Table IV-8, Figure IV-2). The months with the greatest amount of chlorpyrifos use from 2005 through 2009 were May, June, and July (Table IV-8, Figure IV-2). Exceedances occurred during months of the greatest amount of use (Figure IV-2).

The Coalition also uses PUR data to assess which crops receive the most applications of chlorpyrifos. The highest application rates and largest amount of chlorpyrifos applied are associated with almond and walnut orchards (Table IV-9). The most common product containing chlorpyrifos used within this subwatershed was Lorsban (Table IV-9).

Table IV-8. Number of chlorpyrifos applications, total pounds applied and total acres treated by month for January 2005 through December 2009 in the Bear Creek @ Kibby Rd site subwatershed. If a month is not included in the table, no applications were made.

Month/Year	Number of Chlorpyrifos Applications	Pounds of AI Applied	Acres Treated
March, 2005	1	39.6	99
May, 2005	5	347.0	178
June, 2005	2	49.9	25
July, 2005	2	70.0	35
August, 2005	1	55.8	30
September, 2005	1	18.8	10
May, 2006	11	980.6	520
June, 2006	4	154.9	78
July, 2006	3	403.6	200
August, 2006	1	111.4	61
May, 2007	9	363.3	196
June, 2007	1	29.9	15
July, 2007	5	518.7	264
February, 2008	3	492.5	265
May, 2008	2	40.0	20
June, 2008	18	729.1	485
July, 2008	6	458.4	352
August, 2008	2	97.4	106
May, 2009	3	134.8	165
July, 2009	2	193.4	103
Summaries by Year			
2005 Total	12	581.0	377
2006 Total	19	1650.5	859
2007 Total	15	912.0	475
2008 Total	31	1817.4	1228
2009 Total	5	328.2	268
Total	82	5,289.1	3,207

Figure IV-2. Pounds of chlorpyrifos applied within the Bear Creek site subwatershed by month for 2005-2009. Asterisk (*) denotes months with exceedances.




Table IV-9. Total pounds AI for chlorpyrifos based on PUR data from 2005-2009 within the Bear Creek @ Kibby Rd subwatershed.

Chemical Name	Commodity	Product Name	Lbs AI Applied
CHLORPYRIFOS	ALFALFA (FORAGE - FODDER) (ALFALFA HAY)	CHLORPYRIFOS 4E AG	39.629
	ALMOND	GOVERN 4E INSECTICIDE	997.913
		LORSBAN 4E INSECTICIDE	80.113
		LORSBAN 4E-HF	517.405
		LORSBAN-4E	843.047
	CORN FOR/FOD	NUFOS 4E	623.114
		SAURUS	134.764
	PEACH	LORSBAN-4E	157.962
	WALNUT	CHLORPYRIFOS 4E AG	11.258
		LORSBAN 4E-HF	149.539
		LORSBAN ADVANCED	193.434
		LORSBAN-4E	666.940
	WALNUT (ENGLISH WALNUT, PERSIAN WALNUT)	CHLORPYRIFOS 4E AG	46.908
		LORSBAN 4E INSECTICIDE	40.006
		LORSBAN 4E-HF	356.900
		LORSBAN-4E	430.168
ALFALFA – Total Pounds Chlorpyrifos Applied			39.629
ALMOND – Total Pounds Chlorpyrifos Applied			2438.477
CORN – Total Pounds Chlorpyrifos Applied			757.878
PEACH – Total Pounds Chlorpyrifos Applied			157.962
WALNUT – Total Pounds Chlorpyrifos Applied			1895.153
Total pounds chlorpyrifos applied (2005 - 2009)			5,289.100

Based on results from general management practice surveys, approximately 61% of the member acreage indicates no runoff occurs and 32% of the member acreage uses micro and/or drip irrigation. Exceedances of the WQTL occurred in months with the highest application rates, thus the Coalition suspects a combination of spray drift and irrigation water runoff are the cause of chlorpyrifos in the waterways. Growers should be encouraged to review their operation to determine if irrigation return flows are managed properly. Owners operating orchards should evaluate their aerial applications to minimize spray drift.

Coalition outreach since 2007 has included grower meetings and the mailing/distribution of information. A complete list of Coalition Outreach during 2009 is provided in the Summary of Coalition Outreach Activities section of the Management Plan Update Report. The Coalition has begun the process of contacting growers, conducting individual meetings, and compiling individual surveys for Bear Creek. The Coalition focused outreach to growers owning parcels with the potential to drain to the creek and those applying chlorpyrifos. Using these criteria, a list of 13 targeted growers was created. Individual meetings are currently being scheduled and will be completed by the end of May 2010. Individual contacts will focus mainly on chlorpyrifos exceedances however all water quality results will be reviewed and discussed including copper, DO, *E. coli*, and pH WQTLs exceedances as well as toxicity to *Ceriodaphnia dubia*.

The Coalition will compile and analyze the surveys by the end of August 2010. Based on the exact timing of each individual meeting and/or a grower's resources, some owners will not be able to implement recommended management practices until the 2011 irrigation season. In addition, long term structural BMPs (e.g. sediment ponds) will most likely take longer than two years to implement and will require additional tracking to document their implementation. The Coalition will follow up with growers between February and April of 2011 to determine what practices were implemented during the first irrigation year. Growers who have not already been contacted or have indicated that they have not yet implemented practices during the first year will be contacted from February to April of 2012 to determine what practices were implemented during the dormant season and/or the second irrigation year. To evaluate the management practice process, the Coalition will monitor for high priority constituents during months of past exceedances in 2010 and 2011. Depending on when additional management practices are implemented, the Coalition may monitor through 2012 to evaluate improvements in water quality. The Coalition anticipates that changes in management practices by members that have direct drainage and/or are have the potential for spray drift will affect downstream water quality by the end of the second year as a high priority site subwatershed.

The effectiveness of the management practices will be determined through the monitoring of water quality in the years following implementation. However, due to the presence of dairies and nonmembers in the subwatersheds, implementation of management practices by only coalition members may not result in improved water quality. If water quality fails to improve, the Coalition will identify parcels from dairies and nonmembers that could contribute to the exceedances and provide that information to the Regional Board.

Priority C Constituents

Copper

Copper exceeded the hardness based WQTL four times in this subwatershed from 2005 through 2008. Copper exceedances occurred once during the 2007 storm season, twice during the 2008 storm season and once in the 2008 irrigation season in July. Bear Creek was not scheduled for 2009 normal monitoring, but copper will be assessed again during 2010 MPM.

PUR data are reviewed for the number of monthly copper applications, pounds active ingredient (AI) applied, and acres treated (Table IV-10, Figure IV-3). The amount of copper applied within the subwatershed decreased annually from 2005 through 2008, but the amount of copper applied in 2009 increased by 200 lbs compared to 2008 (Table IV-10, Figure IV-3). The months of January, February, April and May indicate the greatest amount of copper use 2005 through 2009 (Table IV-10, Figure IV-3). Not all exceedances occurred during months of the greatest amount of use (Figure IV-3).

The Coalition also used PUR data to assess which crops receive the most applications of copper. The largest amount of copper applied from 2005 through 2009 was associated with almond,

peach, and walnut orchards (Table IV-11). The most common products containing copper used within this subwatershed were DuPont Kocide and Kocide (Table IV-11).

Table IV-10. Number of copper applications, total pounds applied and total acres treated by month for January 2005 through December 2009 in the Bear Creek @ Kibby Rd site subwatershed. If a month is not included in the table, no applications were made.

Month/Year	Number of Copper Applications	Pounds Applied	Acres Treated
January, 2005	2	1178.9	175
February, 2005	3	307.2	121
March, 2005	4	818.6	174
April, 2005	8	1748.0	325.5
May, 2005	8	1762.6	603
November, 2005	1	26.9	40
January, 2006	6	2290.7	493
February, 2006	4	1212.1	244
March, 2006	2	51.9	95
April, 2006	11	1614.5	499
May, 2006	3	323.8	186
June, 2006	2	142.0	88
October, 2006	2	150.6	140
January, 2007	3	1111.0	275
February, 2007	1	729.9	116.91
March, 2007	7	965.3	242
April, 2007	12	1772.9	374
February, 2008	3	2752.6	291.91
March, 2008	8	809.7	320
April, 2008	8	815.6	260
October, 2008	1	67.5	80
January, 2009	3	2709.1	291.91
February, 2009	1	62.9	15
March, 2009	1	122.7	38
April, 2009	2	1562.2	128
May, 2009	1	91.5	34
September, 2009	2	132.8	177.8
Summaries by Year			
2005 Total	26	5842.2	1438.5
2006 Total	30	5785.7	1745
2007 Total	23	4579.1	1007.91
2008 Total	20	4445.4	951.91
2009 Total	10	4681.1	684.71
Total	109	25,333.5	5,828.03

Figure IV-3. Pounds of copper applied within the Bear Creek @ Kibby Rd site subwatershed by month for 2005-2009. Asterisk (*) denotes months with exceedances.




Table IV-11. Total pounds AI for chopper based on PUR data from 2005-2009 within the Bear Creek @ Kibby Rd subwatershed.

Chemical Name	Commodity	Product Name	Lbs AI Applied
COPPER	ALFALFA (FORAGE - FODDER) (ALFALFA HAY)	KOCIDE 2000	48.420
	ALMOND	DUPONT KOCIDE 2000 FUNGICIDE/BACTERICIDE	1533.300
		KOCIDE 101	61.600
		KOCIDE 2000	798.930
		KOCIDE DF	337.700
		NORDOX 75 WG	62.925
	PEACH	BASIC COPPER 53	3528.000
		DUPONT KOCIDE 2000 FUNGICIDE/BACTERICIDE	895.770
		DUPONT KOCIDE 3000 FUNGICIDE/BACTERICIDE	517.242
		KOCIDE DF	2409.950
		NORDOX	395.928
		NORDOX 75 WG	2146.313
	PRUNE	NORDOX 75 WG	1141.040
	TOMATO	CHAMP FLOWABLE	3.459
		CHAMP FORMULA 2 FLOWABLE	155.832
		DUPONT KOCIDE 2000 FUNGICIDE/BACTERICIDE	142.032
		DUPONT KOCIDE 3000 FUNGICIDE/BACTERICIDE	132.786
		KOCIDE 2000	896.308
	WALNUT	DUPONT KOCIDE 2000 FUNGICIDE/BACTERICIDE	536.924
		DUPONT KOCIDE 3000 FUNGICIDE/BACTERICIDE	103.725
		HYDROX	542.080
		KOCIDE 2000	364.495
		KOCIDE DF	24.560
		NU-COP 50DF	3150.840
	WALNUT (ENGLISH WALNUT, PERSIAN WALNUT)	KOCIDE 101	323.400
		KOCIDE 2000	1310.030
KOCIDE DF		1036.432	
NORDOX		21.150	
NORDOX 75 WG		41.950	
NU-COP 50DF		2670.360	

Chemical Name	Commodity	Product Name	Lbs AI Applied
		ALFALFA – Total Pounds Copper Applied	48.420
		ALMOND – Total Pounds Copper Applied	2794.455
		PEACH – Total Pounds Copper Applied	9893.203
		PLUMS (INCLUDING PRUNES) – Total Pounds Copper Applied	1141.040
		TOMATO– Total Pounds Copper Applied	1330.418
		WALNUT – Total Pounds Copper Applied	10125.946
Total pounds copper applied (2005 - 2009)			25,333.482

The Coalition will continue with its management plan strategy outlined above under the chlorpyrifos outreach section when conducting individual contacts. Orchard and vineyard operators will be advised to consider irrigation water retention and management to prevent copper from entering the waterway with runoff. Storm runoff management relevant to copper applications will also be discussed to prevent the winter exceedances. Individual contacts occurring within this subwatershed are described under the chlorpyrifos outreach section above and will include discussions of copper exceedances and the above management practices.

Priority D Constituents

The Bear Creek Management Plan includes *Ceriodaphnia* toxicity and sediment toxicity to *Hyalella azteca* as priority D constituents. These constituents will remain low priority but will be discussed during individual contacts and annual grower meetings.

***Ceriodaphnia* toxicity**

Toxicity to *Ceriodaphnia* has occurred in May of 2005 and 2006, and in July of 2007. The 2006 and 2007 toxicities both coincided with an exceedance of chlorpyrifos. Due to this correlation, the Coalition’s strategy for eliminating *Ceriodaphnia* toxicity will involve focusing on chlorpyrifos. If chlorpyrifos can be prevented from entering waterways throughout the irrigation season, the Coalition believes that *Ceriodaphnia* toxicity can also be reduced or eliminated.

Priority E Constituents

The Bear Creek Management Plan includes DO, *E. coli*, and pH. These constituents will remain low priority but will be discussed during individual contacts and annual grower meetings.

Evaluation

Bear Creek @ Kibby Rd is one of the second four priority site subwatersheds within the ESJWQC and is in its first year of focused outreach (2010-2012). The Coalition’s strategy

for the Bear Creek subwatershed has been to target growers along or adjacent to the waterway that have the potential to discharge. Focus will be on water retention management practice implementation and minimizing spray drift. Outreach includes grower notification, management practice outreach and education, tracking of management practices implementation, and providing information on special studies of management practice efficacy.

Individual surveys to document current management practice implementations and assess future planned implementations are in the process of being filled out and will be completed by the end of May 2010. The Coalition will then assess these results and plan the final stages of outreach including re-contacting growers to identify newly implemented practices and future monitoring to evaluate water quality improvement.

V. COTTONWOOD CREEK @ RD 20

Management Plan Constituents

Priority A/B

- Diazinon
- Chlorpyrifos

Priority C

- Copper
- Diuron

Priority E

- Dissolved Oxygen
- Lead
- *E. coli*

Description of Cottonwood Creek @ Rd 20 Site Subwatershed

Cottonwood Creek @ Road 20 (37,360 irrigated acres) – This site subwatershed is at the very southern edge of the Coalition region in Madera County and drains into the Eastside Bypass. The immediate upstream agriculture is vineyards, and there are deciduous nuts farther to the east. There are only a few dairies in the Cottonwood Creek site subwatershed (Figure V-1). This site subwatershed includes an upstream location (Cottonwood Creek @ Hwy 145) which was sampled in 2008. Table V-1 includes the station name, station code and target latitude/longitudes for sites sampled within this subwatershed.


Table V-1. Coordinates of the Cottonwood Creek site subwatershed sampling locations.

Station Name	Station Code	Target Latitude	Target Longitude
Cottonwood Creek @ Rd 20*	545XCCART	36.8686	-120.1818
Cottonwood Creek @ Hwy 145 ^U	545XCCAHO	36.9002	-120.0555

* Original ESJWQC sampling site

^U Upstream sites

Figure V-1. Site subwatershed map of land use for the Cottonwood Creek @ Rd 20 sample site.



Subwatershed Monitoring History

Ambient water monitoring conducted at Cottonwood Creek @ Rd 20 was initiated during the storm season of 2005 continuing through 2009 (Table V-2). Cottonwood Creek @ Rd 20 will continue to be monitored as a Core Monitoring location under the current MRPP and will be monitored for all Assessment constituents every third year. The constituents sampled at this location from 2005-2009 are listed in Table V-3. Due to single exceedances that have occurred during previous monitoring, Cottonwood Creek @ Rd 20 is also monitored monthly for *S. capricornutum* (the two toxicities that occurred in May 2008 were considered the same event since one was a resample to test for toxicity persistence one week later), simazine and cyanazine starting in October 2009 (fall season).

Management Plan Monitoring occurred within the Cottonwood Creek site subwatershed in 2007 and 2008 for copper (Table V-4 and Table V-5). Cottonwood Creek was monitored twice a month in June and September 2007 and at Cottonwood Creek @ Hwy 145 (upstream location) monthly from May through September 2008. The Coalition will conduct MPM for copper again starting April 2010.

Cottonwood Creek is scheduled as a high priority site starting in 2010. In January 2010, the Coalition began additional MPM sampling during January, February and March (previously MPM was limited to irrigation months). The Coalition monitored for chlorpyrifos, diazinon and diuron in January and February 2010; these results will be reported on in the Management Plan Update Report to be submitted on April 1, 2011.

Cottonwood Creek is scheduled as a high priority site starting in 2010. Cottonwood Creek is listed as an impaired water body for *E. coli* and unknown toxicity in the proposed 2008 Central Valley Basin Plan 303d list. The expected TMDL completion date for the Cottonwood Creek site subwatershed is scheduled for 2021.

Table V-2. Cottonwood Creek @ Rd 20 sampling events per season and year

	2004		2005		2006		2007		2008			2009			
	Irrigation	Storm	Irrigation	Storm	Irrigation	Storm	Irrigation	Storm	Irrigation	Fall	Winter	Storm	Irrigation	Fall	
Events Sampled	NA	2	5	2	5	0	5	1	5	0	0	1	3	3	
Events Not Sampled	NA	0	0	0	0	2	1	2	1	2	2	0	3	0	
Total	NA	2	5	2	5	2	6	3	6	2	2	1	6	3	

NA indicates that this site was not sampled during this season/year.

Table V-3. Number of analyses performed per analyte in each sampling season and year for the Cottonwod Creek @ Rd 20 sample site. Only environmental samples with a sample replicate and lab replicate number of one are shown.

Method	Analyte	2005		2006		2007		2008			2009			
		Storm	Irrigation	Storm	Irrigation	Storm	Irrigation	Storm	Irrigation	Fall	Winter	Storm	Irrigation	Fall
Field and Physical Parameters														
EPA 110.2	Color	2	5	2	5		5	2	5					
EPA 160.1	Dissolved Solids	2	5	2	5		5	2	5			2	3	2
EPA 160.2	Suspended Solids											2	3	2
EPA 180.1	Turbidity	2	5	2	5		5	2	5			2	3	2
EPA 405.1	BOD				1		2							
EPA 415.1	Total Organic Carbon	2	5	2	5		5	2	5			2	3	2
Walkley-Black	Total Organic Carbon (sediment)												1	
SM 9223	E. coli	2	5	2	6		5	2	5			2	3	2
ASTM D422	Grain size (sediment)												1	
NA	Dissolved Oxygen	2	5	3	5		7	6	8	2	2	2	6	2
NA	Specific Conductivity	2	5	3	5		7	6	8	2	2	2	6	2
NA	pH	2	5	3	5		7	6	8	2	2	2	6	2
Carbamates														
EPA 8321A	Aldicarb				5		5	2	5					
EPA 8321A	Carbaryl				5		5	2	5					
EPA 8321A	Carbofuran				5		5	2	5					
EPA 8321A	Diuron				5		5	2	5					
EPA 8321A	Linuron				5		5	2	5					
EPA 8321A	Methiocarb				5		5	2	5					
EPA 8321A	Methomyl				5		5	2	5					
EPA 8321A	Oxamyl				5		5	2	5					
Organochlorines														
EPA 8081A	DDD(p,p')				5		5	2	5					
EPA 8081A	DDE(p,p')				5		5	2	5					
EPA 8081A	DDT(p,p')				5		5	2	5					
EPA 8081A	Dicofol				5		5	2	5					
EPA 8081A	Dieldrin				5		5	2	5					
EPA 8081A	Endrin				5		5	2	5					
EPA 8081A	Methoxychlor				5		5	2	5					

Method	Analyte	2005		2006		2007		2008			2009			
		Storm	Irrigation	Storm	Irrigation	Storm	Irrigation	Storm	Irrigation	Fall	Winter	Storm	Irrigation	Fall
Organophosphates														
EPA 8141A	Azinphos methyl				5		5	2	5					
EPA 8141A	Chlorpyrifos	2	5	2	5		5	2	5					
EPA 8141A	Diazinon	2	5	2	5		5	2	5					
EPA 8141A	Dimethoate				5		5	2	5					
EPA 8141A	Disulfoton				5		5	2	5					
EPA 8141A	Malathion				5		5	2	5					
EPA 8141A	Methamidophos				5		5	2	5					
EPA 8141A	Methidathion				5		5	2	5					
EPA 8141A	Molinate				5		5	2	5					
EPA 8141A	Parathion, Methyl				5		5	2	5					
EPA 8141A	Phorate				5		5	2	5					
EPA 8141A	Phosmet				5		5	2	5					
EPA 8141A	Thiobencarb				5		5	2	5					
Pyrethroids														
EPA 8081A	Bifenthrin		1	2	5		5							
EPA 8081A	Cyfluthrin, total		1	2	5		5							
EPA 8081A	Cyhalothrin, lambda, total	2	5	2	5		5							
EPA 8081A	Cypermethrin, total	2	5	2	5		5							
EPA 8081A	Esfenvalerate/ Fenvalerate, total	2	5	2	5		5							
EPA 8081A	Permethrin, total	2	5	2	5		5							
Triazines														
EPA 547M	Glyphosate				5		5	2	5					
EPA 549.2M	Paraquat dichloride				5		5	2	5					
EPA 619	Atrazine				5		5	2	5					
EPA 619	Cyanazine				5		5	2	5			1	2	2
EPA 619	Simazine				5		5	2	5			1	2	2
Metals (Total)														
EPA 200.8	Arsenic				5		5	2	5					
EPA 200.8	Boron				5		5	2	5					
EPA 200.8	Cadmium				5		5	2	5					
EPA 200.8	Copper				5		6	2	5					

Method	Analyte	2005		2006		2007		2008			2009			
		Storm	Irrigation	Storm	Irrigation	Storm	Irrigation	Storm	Irrigation	Fall	Winter	Storm	Irrigation	Fall
EPA 200.8	Lead				5		5	2	5					
EPA 200.8	Nickel				5		5	2	5					
EPA 200.8	Selenium				5		2	2	5					
EPA 200.8	Zinc				5		5	2	5					
Metals (Dissolved)														
EPA 200.8	Cadmium													
EPA 200.8	Copper													
EPA 200.8	Lead													
EPA 200.8	Nickel													
EPA 200.8	Zinc													
Nutrients														
SM 2340 C	Hardness as CaCO3				5		6	2	5					
EPA 300.0	Nitrate as N				5		5	2	5					
EPA 350.2	Ammonia as N				5		5	2	5			2	3	2
EPA 351.3	Nitrogen, Total Kjeldahl				5		5	2	5			1		
EPA 354.1	Nitrite as N				5		5	2	5					
EPA 353.2	Nitrate + Nitrite as N											2	3	2
EPA 365.2	Orthophosphate as P				5		5	2	5			1		
EPA 365.2	Phosphate as P				5		5	2	5			2	3	2
Toxicity														
EPA 821/R-02-012	Ceriodaphnia dubia	2	5	2	5		6	2	5					
EPA 821/R-02-012	Pimephales promelas	2	5	2	5		6	3	5			1	3	2
EPA 821/R-02-013	Selenastrum capricornutum	2	5	2	5		6		6			1	3	2
EPA 600/R-99-064	Hyalella azteca		5	1	1		1	1	1				1	

Table V-4. Cottonwood Creek site subwatershed. 2007 Management Plan additional (A) sampling schedule for copper. "X" indicates the site, month, and analyte sampled.

Sample Site	Month	Type	Copper
Cottonwood Creek @ Rd 20	19-Jun-07	A	X
Cottonwood Creek @ Rd 20	26-Sep-07	A	X

Table V-5. Cottonwood Creek site subwatershed. 2008 Management Plan upstream (U) sampling schedule for copper. "X" indicates the site, month and analyte sampled.

Sample Site	Date	Type	Copper
Cottonwood Creek @ Hwy 145	27-May-08	U	X
Cottonwood Creek @ Hwy 145	24-Jun-08	U	X
Cottonwood Creek @ Hwy 145	29-Jul-08	U	X
Cottonwood Creek @ Hwy 145	26-Aug-08	U	X
Cottonwood Creek @ Hwy 145	30-Sep-08	U	X

Exceedance History

During Coalition monitoring, exceedances of WQTLs for field and physical parameters, *E. coli*, nutrients, metals, pesticides, and water column and sediment toxicity occurred within the Cottonwood Creek site subwatershed (Table V-6). During ambient water monitoring within Cottonwood Creek (including the upstream sampling at Cottonwood Creek @ Hwy 145) exceedances of field, physical, inorganic and organic constituents have occurred including DO (15), pH (1), *E. coli* (14), total copper (13), total lead (3), chlorpyrifos (2), cyanazine (1), diazinon (1), simazine (1), and diuron (2, Table V-6). Copper exceedances have occurred during all irrigation months (May through September) as well as during storm sampling events (January and February); one of those exceedances occurred in June at the upstream sampling location at Hwy 145. Both chlorpyrifos exceedances occurred in the same year (2008) in January and February. In 2008, water column toxicity occurred three times, twice to *S. capricornutum* and once to *P. promelas* (Table V-6). The *S. capricornutum* toxicity occurred in a sample collected in April 2008 and again in the resample collected one week later. Sediment toxicity to *H. azteca* occurred once in March 2008 (Table V-6).

During 2009, the only WQTL exceedances that occurred were for *E. coli* (February and November) and DO (May, Table V-6).

All exceedances are listed in Table V-6 by season and date and are based on WQTLs listed in the introduction to the ESJWQC Management Plan. A management plan is required if a site experiences two or more exceedances of the particular constituent within three years or if a site has a single exceedance of a constituent that has a TMDL. Using the ESJWQC Management Plan prioritization process flow chart (Figure 4), priorities were assigned to management plan constituents and are listed in the bottom row of Table V-6. The highest priority constituents in the Cottonwood Creek subwatershed are chlorpyrifos (A/B), diazinon (A/B), copper (C), and diuron (C).

Table V-6. All exceedances experienced in samples collected from locations within the Cottonwood Creek site subwatershed between February 2005 and December 2009 (sorted by season and date). If the water quality trigger limit is based on hardness then the hardness value is shown in parenthesis; otherwise the WQTL used to evaluate the data is listed in the header after the analyte.

Station Name	Season	Sample Date	Oxygen, Dissolved, 7 mg/L	pH, 6.5-8.5 units	<i>E. coli</i> , 235 MPN/100 mL	Copper, Total 1,300 µg/L	Lead, Total, 15 µg/L	Chlorpyrifos, 0.015 µg/L	Cyanazine, 1 µg/L	Diazinon, 0.1 µg/L	Diuron, 2 µg/L	Simazine, 4 µg/L	<i>P. promelas</i> , Survival (%)	<i>S. capricornutum</i> , Total Cell Count	<i>H. azteca</i> , Survival (%)
Cottonwood Creek @ Rd 20	Storm	2/16/2005			1600										
Cottonwood Creek @ Rd 20	Storm	3/21/2005	5.6		1600										
Cottonwood Creek @ Rd 20	Irrigation	5/10/2005			540										
Cottonwood Creek @ Rd 20	Irrigation	6/14/2005	5.7												
Cottonwood Creek @ Rd 20	Irrigation	7/12/2005	5.17												
Cottonwood Creek @ Rd 20	Irrigation	8/16/2005			300										
Cottonwood Creek @ Rd 20	Irrigation	9/20/2005	6.5												
Cottonwood Creek @ Rd 20	Storm	2/28/2006			300										
Cottonwood Creek @ Rd 20	Storm	3/15/2006			1600										
Cottonwood Creek @ Rd 20	Irrigation	5/16/2006	5.71			4.4 (3.5)									
Cottonwood Creek @ Rd 20	Irrigation	6/13/2006	6.9			8 (6.9)	0.73 (0.63)								
Cottonwood Creek @ Rd 20	Irrigation	7/11/2006	6.51												
Cottonwood Creek @ Rd 20	Irrigation	8/8/2006	6.95												
Cottonwood Creek @ Rd 20	Irrigation	9/12/2006	6.11			5.5 (4.4)									
Cottonwood Creek @ Rd 20	Irrigation	5/29/2007	6.55			6.7 (5.5)									
Cottonwood Creek @ Rd 20	Irrigation	6/19/2007				6.7 (4.1)									
Cottonwood Creek @ Rd 20	Irrigation	6/26/2007				4.3 (4.1)									
Cottonwood Creek @ Rd 20	Irrigation	7/24/2007		9.04		5.4 (4.6)									

Station Name	Season	Sample Date	Oxygen, Dissolved, 7 mg/L	pH, 6.5-8.5 units	<i>E. coli</i> , 235 MPN/100 mL	Copper, Total 1,300 µg/L	Lead, Total, 15 µg/L	Chlorpyrifos, 0.015 µg/L	Cyanazine, 1 µg/L	Diazinon, 0.1 µg/L	Diuron, 2 µg/L	Simazine, 4 µg/L	<i>P. promelas</i> , Survival (%)	<i>S. capricornutum</i> , Total Cell Count	<i>H. azteca</i> , Survival (%)
Cottonwood Creek @ Rd 20	Irrigation	8/21/2007	6.81			5.2 (4.6)									
Cottonwood Creek @ Rd 20	Irrigation	8/23/2007	3.95												
Cottonwood Creek @ Rd 20	Storm	1/25/2008			1200	24 (3.0)	5.4 (0.6)	0.02			68		45		
Cottonwood Creek @ Rd 20	Storm	2/25/2008				21 (6.5)	1.9 (1.9)	0.04		0.2	65	5			
Cottonwood Creek @ Rd 20	Sediment	3/4/2008													96
Cottonwood Creek @ Rd 20	Irrigation	4/29/2008			580	8 (6.9)								58382	
Cottonwood Creek @ Rd 20	Irrigation	5/7/2008												130025	
Cottonwood Creek @ Rd 20	Irrigation	5/27/2008			250				1						
<i>Cottonwood Creek @ Hwy 145</i>	<i>Irrigation</i>	<i>6/24/2008</i>				<i>39 (5.5)</i>									
Cottonwood Creek @ Rd 20	Irrigation	6/24/2008			1300										
Cottonwood Creek @ Rd 20	Irrigation	7/29/2008			1000										
<i>Cottonwood Creek @ Hwy 145</i>	<i>Irrigation</i>	<i>8/26/2008</i>	<i>6.45</i>												
Cottonwood Creek @ Rd 20	Irrigation	8/26/2008	6.83		390	4.4 (3.7)									
Cottonwood Creek @ Rd 20	Storm	2/7/2009			>2400										
Cottonwood Creek @ Rd 20	Irrigation	5/19/2009	6.72												
Cottonwood Creek @ Rd 20	Fall	11/17/2009			770										
Constituent Priority			E	NP	E	C	E	A/B	NP	A/B	C	NP	NP	NP¹	NP

NP- Not prioritized. Fewer than two exceedances for this constituent at this site within three years and currently no TMDL for constituent
NP¹ – Not prioritized; both toxic samples were from the same sampling event (sample and resample to test for persistence).

2007 - 2009 Management Plan Monitoring Results

Management Plan monitoring results are included in Table V-7 for copper for the years 2007 and 2008; MPM did not occur for copper during 2009. Chlorpyrifos, diazinon, and diuron first exceeded their respective WQTLs during 2008 storm sampling (January and February) and MPM monitoring for these constituents was initiated in January 2010.

2007

In 2007, Cottonwood Creek @ Rd 20 was monitored monthly for copper from April through September plus additional copper monitoring was conducted in June and September. The copper WQTL was exceeded in all samples analyzed from May through August during normal monitoring including the additional sample collected in June 2007 (Table V-6). The April sample had detectable amounts of copper (3.9µg/L) however the concentration was not an exceedance. Cottonwood Creek was dry in September and therefore there are no results for NM or MPM (Table V-7).

2008

In accordance with the 2008 upstream MPM schedule, Cottonwood Creek @ Hwy 145 was monitored for copper from May through September 2008 (previous section, Table IV-5). Copper was detected in all samples collected during the irrigation season except in September when the site was dry. The copper WQTL was exceeded in samples collected during April and August at the normal monitoring site, but never at the upstream monitoring site (Table V-7).

2009

Cottonwood Creek @ Rd 20 was not scheduled for 2009 MPM.

Management Plan Monitoring for chlorpyrifos, diazinon and diuron (exceedances occurred during storm monitoring in 2008) took place in January and February of 2010 and will be reported in the 2011 Management Plan Update Report.

Management Plan Monitoring for copper will be conducted during the 2010 irrigation season. Management Plan Monitoring during January and February 2011 is also scheduled for copper, chlorpyrifos, diuron, and diazinon.

Table V-7. Cottonwood Creek site subwatershed. Normal monitoring (NM) and Management Plan Monitoring (MPM) results where 'A' indicates additional MPM (2007) and 'US' indicates upstream MPM (2008) for copper. Exceedance values are in bold.

	Month:	April	May	June	July	August	September
2007 NM (@ Rd 20)	Date	4/24/07	5/29/07	6/26/07	7/24/07	8/21/07	9/18/07
	Copper (µg/L)	3.9	6.7	4.3	5.4	5.2	Dry
2007 MPM A (@Rd 20)	Date	NA	NA	6/19/07	NA	NA	9/25/07
	Copper (µg/L)	NA	NA	6.7	NA	NA	Dry
2008 NM (@ Rd 20)	Date	4/29/08	5/27/08	6/24/08	7/29/08	8/26/08	9/30/08
	Copper (µg/L)	8 (6.9)	4.9	4.5	4.8	4.4 (3.7)	Dry
2008 MPM US (@ Hwy 145)	Date	NA	5/27/08	6/24/08	7/29/08	8/26/08	9/30/08
	Copper (µg/L)	NA	2.4	39	2.3	2.1	Dry

NA - Not applicable. This site was not sampled during this month.

Load Calculations

Loads have been calculated for all chlorpyrifos detections in the site subwatershed (Table V-8) based on the following formula:

$$\text{Load} = \text{Discharge (cfs)} \times 28.317\text{L/ft}^3 \times \text{Concentration (milligram/L} \times 1000 \text{ or } \mu\text{g/L)}.$$

The load values for constituents in this report represent instantaneous loads only. These values should not be used to extrapolate loading over any period of time (e.g. weekly, monthly, seasonal or annual). The primary purpose for reporting instantaneous loads is to provide a normalization of the concentrations by flow for various constituents at the time the samples were collected.

Table V-8. Cottonwood Creek site subwatershed. Instantaneous load calculations for chlorpyrifos, copper, diazinon and diuron where discharge was measured (sorted by analyte, site and date).

Station Name	Analyte Name	Sample Date	Discharge cfs	Concentration $\mu\text{g/L}$	Loading Rate $\mu\text{g/sec}$
Cottonwood Creek @ Rd 20*	Chlorpyrifos	15-Mar-06	0	0.012	0
Cottonwood Creek @ Rd 20	Chlorpyrifos	15-Mar-06	0	0.011	0
Cottonwood Creek @ Rd 20	Chlorpyrifos	11-Jul-06	1.96	0.014	0.78
Cottonwood Creek @ Rd 20	Chlorpyrifos	24-Jul-07	0	0.007	0
Cottonwood Creek @ Rd 20	Chlorpyrifos	25-Jan-08	3.05	0.019	1.64
Cottonwood Creek @ Rd 20	Chlorpyrifos	25-Feb-08	0.22	0.036	0.22
Cottonwood Creek @ Rd 20	Copper	16-May-06	2.33	4.4	290.31
Cottonwood Creek @ Rd 20*	Copper	16-May-06	2.33	4.8	316.70
Cottonwood Creek @ Rd 20	Copper	13-Jun-06	2.66	8	602.59
Cottonwood Creek @ Rd 20	Copper	11-Jul-06	1.96	5.3	294.16
Cottonwood Creek @ Rd 20	Copper	8-Aug-06	12.92	4.1	1500.01
Cottonwood Creek @ Rd 20	Copper	12-Sep-06	38.2	5.5	5949.40
Cottonwood Creek @ Rd 20	Copper	24-Apr-07	28.24	3.9	3118.72
Cottonwood Creek @ Rd 20	Copper	19-Jun-07	0.59	6.7	111.94
Cottonwood Creek @ Rd 20	Copper	24-Jul-07	0	5.4	0
Cottonwood Creek @ Rd 20*	Copper	24-Jul-07	0	5.1	0
Cottonwood Creek @ Rd 20	Copper	25-Jan-08	3.05	24	2072.80
Cottonwood Creek @ Rd 20	Copper	25-Feb-08	0.22	21	130.82
Cottonwood Creek @ Rd 20	Copper	29-Apr-08	8.26	8	1871.19
Cottonwood Creek @ Rd 20	Copper	27-May-08	17.41	4.9	2415.69
Cottonwood Creek @ Rd 20	Copper	24-Jun-08	0.08	4.5	10.19
Cottonwood Creek @ Rd 20	Copper	29-Jul-08	0.16	4.8	21.75
Cottonwood Creek @ Rd 20	Copper	26-Aug-08	0.79	4.4	98.43
<i>Cottonwood Creek at Highway 145</i>	<i>Copper</i>	<i>27-May-08</i>	<i>20.01</i>	<i>2.4</i>	<i>1359.90</i>
<i>Cottonwood Creek at Highway 145</i>	<i>Copper</i>	<i>24-Jun-08</i>	<i>60.24</i>	<i>39</i>	<i>66526.83</i>

Station Name	Analyte Name	Sample Date	Discharge cfs	Concentration µg/L	Loading Rate µg/sec
<i>Cottonwood Creek at Highway 145</i>	<i>Copper</i>	<i>29-Jul-08</i>	<i>35.03</i>	<i>2.3</i>	<i>2281.47</i>
<i>Cottonwood Creek at Highway 145</i>	<i>Copper</i>	<i>26-Aug-08</i>	<i>10.85</i>	<i>2.1</i>	<i>645.20</i>
Cottonwood Creek @ Rd 20	Diazinon	28-Feb-06	0.26	0.02	0.15
Cottonwood Creek @ Rd 20*	Diazinon	28-Feb-06	0.26	0.023	0.17
Cottonwood Creek @ Rd 20	Diazinon	25-Jan-08	3.05	0.06	5.18
Cottonwood Creek @ Rd 20	Diazinon	25-Feb-08	0.22	0.24	1.50
Cottonwood Creek @ Rd 20	Diuron	24-Apr-07	28.24	0.85	679.72
Cottonwood Creek @ Rd 20	Diuron	25-Jan-08	3.05	68	5872.95
Cottonwood Creek @ Rd 20	Diuron	25-Feb-08	0.22	65	404.93
Cottonwood Creek @ Rd 20	Diuron	29-Apr-08	8.26	0.63	147.36
Cottonwood Creek @ Rd 20	Diuron	27-May-08	17.41	0.23	113.39

*Field Duplicate

Source Identification and Outreach

Priority A/B Constituents

Chlorpyrifos and diazinon are the only priority A/B constituents listed under the Cottonwood Creek Management Plan.

Chlorpyrifos

Chlorpyrifos has exceeded the WQTL of 0.015µg/L two times in this subwatershed from 2005 through 2009 (January and February 2008). Chlorpyrifos was omitted from the monitoring schedule in October 2008 when the Coalition began monitoring under a new MRPP. Starting in 2010, as part of the MPM schedule, chlorpyrifos will be monitored for in January and February. The Coalition uses a combination of monitoring data and evaluation of Pesticide Use Report (PUR) data to identify possible sources.

PUR data are reviewed for the number of monthly chlorpyrifos applications, pounds active ingredient (AI) applied, and acres treated (Table V-9, Figure V-2). The greatest chlorpyrifos use occurred in 2006 (7,978 lbs) with the lowest use occurring in 2007 (406 lbs) and 2009 (280 lbs, Table V-9, Figure V-2). In 2005, 2007, and 2008 the greatest amount of chlorpyrifos use occurred in July whereas in 2006, the greatest use occurred in May and in 2009 during October (Table V-9, Figure V-2). The most recent chlorpyrifos use preceding the 2008 January and February chlorpyrifos exceedances occurred in August 2007; no use was reported from September 2007 through February 2008 (Figure V-2).

The Coalition also used PUR data to assess which crops receive the most applications of chlorpyrifos. The highest application rates are associated with almond orchards, orange groves and vineyards/grapes (Table V-10). The largest amount of chlorpyrifos from 2005 through 2009 was to almond orchards (10,012 lbs) followed by orange groves (2,875 lbs) and

vineyards/grapes (1,794 lbs, Table V-10). The most common product containing chlorpyrifos used within this subwatershed was Lorsban (Table V-10).

Table V-9. Number of chlorpyrifos applications, total pounds applied and total acres treated by month for January 2005 through December 2009 in the Cottonwood Creek @ Rd 20 site subwatershed. If a month is not included in the table, no applications were made.

Month/Year	Number of Chlorpyrifos Applications	Pounds Applied	Acres Treated
February, 2005	2	405.8	203
March, 2005	1	0.4	0.1
May, 2005	1	245.2	41
June, 2005	1	6.0	80
July, 2005	17	1631.3	865
August, 2005	11	1198.8	311.1
September, 2005	16	628.0	212
February, 2006	3	353.1	181
March, 2006	2	131.9	71
May, 2006	21	4107.6	1179
June, 2006	4	277.4	166.5
July, 2006	17	1586.3	792.46
August, 2006	10	784.6	220.75
September, 2006	7	244.4	88.5
October, 2006	4	492.5	110
March, 2007	1	326.1	163
May, 2007	1	14.9	8
July, 2007	8	369.9	197.2
August, 2007	3	75.8	38
March, 2008	1	326.0	163
July, 2008	20	1128.8	482
August, 2008	14	662.6	344
October, 2008	2	40.4	10
July, 2009	2	156.9	67
August, 2009	2	146.8	73
October, 2009	5	262.9	140
Summaries by Year			
2005 Total	49	4115.6	1712.2
2006 Total	68	7977.9	2809.21
2007 Total	13	786.6	406.2
2008 Total	37	2157.9	999
2009 Total	9	566.7	280
Total	176	15,604.6	6,206.61

Figure V-2. Pounds of chlorpyrifos applied within the Cottonwood Creek @ Rd 20 site subwatershed by month for 2005-2009. Asterisk (*) denotes months with exceedances.




Table V-10. Total pound AI for chlorpyrifos based on PUR data from 2005-2009 within the Cottonwood Creek @ Rd 20 subwatershed.

Chemical Name	Commodity	Product Name	Lbs AI Applied
CHLORPYRIFOS	ALFALFA (FORAGE - FODDER) (ALFALFA HAY)	LORSBAN-4E	23.657
	ALMOND	CHLORPYRIFOS 4E AG	0.404
		GOVERN 4E INSECTICIDE	2014.002
		LORSBAN 4E-HF	970.009
		LORSBAN ADVANCED	75.120
		LORSBAN-4E	6696.266
		NUFOS 4E	256.225
	CHERRY	LORSBAN 4E INSECTICIDE	27.072
	CORN (FORAGE - FODDER)	LORSBAN 4E-HF	117.637
	GRAPES	LORSBAN 4E INSECTICIDE	1304.188
		LORSBAN 4E-HF	79.754
	GRAPES, RAISIN	LORSBAN ADVANCED	75.120
		LORSBAN-4E	14.870
	GRAPES, WINE	LORSBAN ADVANCED	187.800
		LORSBAN-4E	131.945
	ORANGE	GOVERN 4E INSECTICIDE	409.063
	ORANGE (ALL OR UNSPEC)	CHLORPYRIFOS 4E AG	6.289
		GOVERN 4E INSECTICIDE	303.010
		LORSBAN 4E-HF	936.712
		LORSBAN-4E	1219.588
	TANGELO	LORSBAN-4E	306.633
	TANGERINE (MANDARIN, SATSUMA, MURCOTT, ETC.)	GOVERN 4E INSECTICIDE	84.843
		LORSBAN-4E	160.815
	TANGERINE, SEEDL	GOVERN 4E INSECTICIDE	40.401
	WALNUT	GOVERN 4E INSECTICIDE	80.803
		LORSBAN-4E	61.325
	WALNUT (ENGLISH WALNUT, PERSIAN WALNUT)	LORSBAN 4E-HF	0.997
		LORSBAN-4E	20.010
ALFALFA – Total Pounds Chlorpyrifos Applied			23.657
ALMOND – Total Pounds Chlorpyrifos Applied			10012.024
CHERRY – Total Pounds Chlorpyrifos Applied			27.072
CORN – Total Pounds Chlorpyrifos Applied			117.637
GRAPES – Total Pounds Chlorpyrifos Applied			1793.677
ORANGE – Total Pounds Chlorpyrifos Applied			2874.662
TANGELO – Total Pounds Chlorpyrifos Applied			306.633
TANGERINE – Total Pounds Chlorpyrifos Applied			286.058
WALNUT – Total Pounds Chlorpyrifos Applied			163.134
Total pounds chlorpyrifos applied (2005 - 2009)			15,604.557

Based on results from general management practice surveys, approximately 82% of the member acreage has no runoff and over 40% of the member acreage uses micro and/or drip

irrigation. Cottonwood Creek has sandy soils and a large percentage of orchards. Therefore it is likely that the exceedances of chlorpyrifos that occurred in 2008 were due to spray drift. Although there was no reported use in December, January or February preceding the exceedances, applications of chlorpyrifos during this time of year would be either for Vine Mealybug control (applied to grapes) or dormant sprays (applied to orchards like cherries).

Coalition outreach since 2007 has included grower meetings and the mailing/distribution of information. A complete list of Coalition Outreach during 2009 is provided in the Summary of Coalition Outreach Activities section of the Management Plan Update Report. The Coalition has begun the process of contacting growers, conducting individual meetings, and compiling individual surveys. The Coalition focused outreach to growers owning parcels with the potential to drain to the creek and those applying chlorpyrifos. Using these criteria, a list of 25 targeted growers was created. Targeted growers were made aware of the requirement to schedule an individual meeting with Coalition representatives through conference call meetings in October 2009. Individual meetings are currently ongoing and will be completed by the end of May 2010. Individual contacts will focus mainly on chlorpyrifos exceedances however all water quality results will be reviewed and discussed including diazinon, copper, diuron, *E. coli*, DO, and lead exceedances.

The Coalition will compile and analyze the surveys by the end of August 2010. Based on the exact timing of each individual meeting and/or a grower's resources, some owners will not be able to implement recommended management practices until the 2011 irrigation season. In addition, long term structural BMPs (e.g. sediment ponds) will most likely take longer than two years to implement and will require additional tracking to document their implementation. The Coalition will follow up with growers between February and April of 2011 to determine what practices were implemented during the first irrigation year. Growers who have not already been contacted or have indicated that they have not yet implemented practices during the first year will be contacted from February to April of 2012 to determine what practices were implemented during the dormant season and/or the second irrigation year. To evaluate the management practice process, the Coalition will monitor for high priority constituents during months of past exceedances in 2010 and 2011. Depending on when additional management practices are implemented, the Coalition may monitor through 2012 to evaluate improvements in water quality. The Coalition anticipates that changes in management practices by members that have direct drainage and/or have the potential for spray drift will affect downstream water quality by the end of the second year as a high priority site subwatershed.

The effectiveness of the management practices will be determined through the monitoring of water quality in the years following implementation. However, due to the presence of dairies and nonmembers in the subwatersheds, implementation of management practices by only coalition members may not result in improved water quality. If water quality fails to improve, the Coalition will identify parcels from dairies and nonmembers that could contribute to the exceedances and provide that information to the Regional Board.

Diazinon

Concentrations of diazinon in samples have exceeded the WQTL of 0.1 µg/L once in this subwatershed from 2005 through 2009 (February 25, 2008, Table V-6). Diazinon was omitted from the monitoring schedule in October 2008 however MPM for diazinon will begin in January 2010. The Coalition uses a combination of monitoring data and evaluation of Pesticide Use Report (PUR) data to identify possible sources.

PUR data are reviewed for the number of monthly diazinon applications, pounds active ingredient (AI) applied, and acres treated (Table V-11, Figure V-3). Overall, the amount of diazinon applied within the subwatershed has decreased since 2005. The amount of diazinon used in 2005 was more than thirteen times the amount of diazinon used in 2009 (976 lbs in 2005 compared to 71 lbs in 2009, Table V-11, Figure V-3). Diazinon use generally occurs between December and March during dormant sprays (Table V-11, Figure V-3). Exceedances of diazinon have not occurred during the months of highest use (i.e. January 2006 and January 2007, Figure V-3). However, the exceedance that occurred in February 2008 was also the month of greatest diazinon use in 2008 (Table V-11, Figure V-3).

The Coalition also used PUR data to assess which crops receive the most applications of diazinon. The greatest pounds applied are associated with peaches, plums, almonds, and nectarines (Table V-12). The largest amount of diazinon from 2005 through 2009 was to almond and peach orchards (Table V-12). Diazinon was applied to peach, plum, almond, cherry and fig orchards prior to the February 2008 exceedance.

Table V-11. Number of diazinon applications, total pounds applied and total acres treated by month for January 2005 through December 2009 in the Cottonwood Creek @ Rd 20 site subwatershed. If a month is not included in the table, no applications were made.

Month/Year	Number of Diazinon Applications	Pounds Applied	Acres Treated
January, 2005	7	367.9	184
February, 2005	8	399.8	200
May, 2005	6	5.8	4.2
June, 2005	2	162.5	50
July, 2005	1	4.0	4
August, 2005	1	35.8	35.8
January, 2006	18	816.8	498.8
February, 2006	3	86.1	42.5
May, 2006	7	8.1	32.2
June, 2006	1	31.2	40
January, 2007	13	778.0	394.71
February, 2007	1	20.0	10
March, 2007	1	42.0	28
January, 2008	6	222.7	114.5

Month/Year	Number of Diazinon Applications	Pounds Applied	Acres Treated
February, 2008	8	383.8	300.5
March, 2008	1	42.0	28
December, 2009	1	71.4	36
Summaries by Year			
2005 Total	25	975.7	478
2006 Total	29	942.2	613.5
2007 Total	15	840.0	432.71
2008 Total	15	648.4	443
2009 Total	1	71.4	36
Total	85	3,477.7	2,003.21

Figure V-3. Pounds of diazinon applied within the Cottonwood Creek @ Rd 20 site subwatershed by month for 2005-2009. Asterisk (*) denotes months with exceedances.




Table V-12. Total pound AI for diazinon based on PUR data from 2005-2009 within the Cottonwood Creek @ Rd 20 subwatershed.

Chemical Name	Commodity	Product Name	Lbs AI Applied
DIAZINON	ALMOND	CLEAN CROP DIAZINON AG500 INSECTICIDE	417.882
		DIAZINON AG 500	170.659
		GOWAN DIAZINON 4E	105.958
	APPLE	DIAZINON 50W	0.431
	APRICOT	DIAZINON 50W	4.681
	CHERRY	CLEAN CROP DIAZINON AG500 INSECTICIDE	39.393
		DIAZOL 50 W	20.000
		GOWAN DIAZINON 4E	19.992
	FIG	DIAZOL AG 500	66.994
	GRAPES, WINE	GOWAN DIAZINON 50 WSB	39.800
	NECTARINE	DIAZINON 50W	4.250
		DIAZINON AG 500	79.376
		GOWAN DIAZINON 4E	212.358
	PEACH	CLEAN CROP DIAZINON AG500 INSECTICIDE	661.860
		DIAZINON 50W	4.250
		DIAZINON AG 500	156.768
		DIAZOL 50 W	73.000
		GOWAN DIAZINON 4E	757.987
	PLUM	CLEAN CROP DIAZINON AG500 INSECTICIDE	86.124
	PLUM (INCLUDES WILD PLUMS FOR HUMAN CONSUMPTION)	DIAZINON 50W	4.750
		GOWAN DIAZINON 4E	59.976
		CLEAN CROP DIAZINON AG500 INSECTICIDE	60.792
	PRUNE	DIAZINON 50W	93.750
DIAZINON AG500 INSECTICIDE		182.232	
DIAZOL 50 W		42.000	
WALNUT (ENGLISH WALNUT, PERSIAN WALNUT)	GOWAN DIAZINON 4E	112.455	
		ALMOND – Total Pounds Diazinon Applied	694.498
		APPLE – Total Pounds Diazinon Applied	0.431
		APRICOT – Total Pounds Diazinon Applied	4.681
		CHERRY – Total Pounds Diazinon Applied	79.385
		FIG – Total Pounds Diazinon Applied	66.994
		GRAPES – Total Pounds Diazinon Applied	39.800
		NECTARINE – Total Pounds Diazinon Applied	295.984
		PEACH – Total Pounds Diazinon Applied	1653.865
		PLUM (INCLUDING PRUNES) – Total Pounds Diazinon Applied	529.624
		WALNUT – Total Pounds Diazinon Applied	112.455
		Total pounds diazinon applied (2005 - 2009)	3,477.717

Ten applications of diazinon occurred within four weeks prior to the single February 2008 exceedance; however other months with high application rates did not have exceedances. The exceedance is most likely due to spray drift from parcels directly next to the creek. Therefore the Coalition has focused its outreach to target specific growers with the potential to discharge to the creek as well as growers with parcels next to the creek with the potential for spray drift; this is the same strategy used with chlorpyrifos outreach. Orchard operators will be advised to consider storm runoff relevant to diazinon applications to prevent the winter exceedances of the pesticide. Individual contacts occurring within this subwatershed are described under the chlorpyrifos outreach section above and include discussions of diazinon exceedances as well as spray drift and storm runoff management practices.

Priority C Constituents

Priority C constituents for Cottonwood Creek are copper and diuron.

Copper

Copper has exceeded the hardness based WQTL numerous times in this subwatershed from 2005 through 2009 based on results from total copper analysis . Additional and upstream MPM confirmed that copper is a water quality concern throughout the subwatershed during most times of the year. The Coalition will begin MPM monitoring for copper (both total and dissolved) in April 2010. PUR data are reviewed for the number of monthly copper applications, pounds active ingredient (AI) applied, and acres treated (Table V-13, Figure V-4). Since 2006 there has been a general decrease in the pounds of copper applied within the subwatershed; the amount of copper used in 2005 was almost twice the amount of copper used in 2009 (23,732 lbs in 2005 compared to 12,388 lbs in 2009, Table V-13, Figure V-4). In 2005 and 2006 the greatest amount of copper use occurred in April whereas in 2007 the greatest use occurred in March, in 2008 the greatest use occurred in January, and in 2009 it was September (Table V-13, Figure V-4). Not all exceedances occurred during months of the greatest amount of use (Figure V-4).

The largest amount of copper applied from 2005 through 2009 was associated with grapes (including wine and raisin), almonds, oranges and peaches (Table V-14). The most common products containing copper used within this subwatershed are DuPont Kocide and Kocide (Table V-14).

Table V-13. Number of copper applications, total pounds applied and total acres treated by month for January 2005 through December 2009 in the Cottonwood Creek @ Rd 20 site subwatershed. If a month is not included in the table, no applications were made.

Month/Year	Number of Copper Applications	Pounds Applied	Acres Treated
January, 2005	21	3147.4	704.3
February, 2005	17	2121.8	391
March, 2005	78	5544.1	4534.66
April, 2005	111	6509.6	5140.24
May, 2005	8	133.6	142
June, 2005	10	710.2	561
August, 2005	9	1473.7	487
September, 2005	3	705.0	235
October, 2005	8	1464.9	278
November, 2005	21	1645.3	420.7
December, 2005	1	276.3	60
January, 2006	48	8582.9	2205.8
February, 2006	24	1357.6	307.2

Month/Year	Number of Copper Applications	Pounds Applied	Acres Treated
March, 2006	16	1144.0	993.6
April, 2006	237	12968.4	13377.61
May, 2006	45	2657.9	2294.1
June, 2006	9	640.3	450
July, 2006	3	281.6	107
August, 2006	3	393.8	190
September, 2006	1	108.0	36
October, 2006	9	1567.0	380.8
November, 2006	14	1212.7	478.25
December, 2006	5	677.2	183.5
January, 2007	23	2444.0	829.71
February, 2007	7	715.2	363
March, 2007	77	3404.8	3981.36
April, 2007	49	2094.1	2428.06
May, 2007	8	120.8	138.8
July, 2007	18	2712.0	1368.42
August, 2007	4	220.0	60
September, 2007	8	1170.1	382
October, 2007	1	94.1	16
January, 2008	45	7245.7	3127.76
February, 2008	12	875.5	313.5
March, 2008	46	1328.0	2369.98
April, 2008	102	4659.1	5422.97
June, 2008	14	269.3	136
August, 2008	10	405.4	344
October, 2008	10	812.2	288
November, 2008	12	2920.6	1202
January, 2009	9	1380.5	576
February, 2009	1	22.6	10
March, 2009	39	851.5	1764.4
April, 2009	64	2168.9	3290.46
May, 2009	7	184.8	186
August, 2009	1	120.0	40
September, 2009	1	4890.5	163
October, 2009	13	1387.5	800
November, 2009	6	1381.3	425
Summaries by Year			
2005 Total	287	23731.9	12953.9
2006 Total	414	31591.4	21003.86
2007 Total	195	12975.1	9567.35

Month/Year	Number of Copper Applications	Pounds Applied	Acres Treated
2008 Total	251	18515.7	13204.21
2009 Total	141	12387.5	7254.86
Total	1,288	99,201.6	63,984.18

Figure V-4. Pounds of copper applied within the Cottonwood Creek @ Rd 20 site subwatershed by month for 2005-2009. Asterisk (*) denotes months with exceedances.




Table V-14. Total pound AI for copper based on PUR data from 2005-2009 within the Cottonwood Creek @ Rd 20 subwatershed.

Chemical Name	Commodity	Product Name	Lbs AI Applied
COPPER	ALFALFA	DUPONT KOCIDE 2000 FUNGICIDE/BACTERICIDE	2.421
	ALMOND	CHAMP FORMULA 2 FLOWABLE	7475.489
		DUPONT KOCIDE 2000 FUNGICIDE/BACTERICIDE	2182.662
		DUPONT KOCIDE 3000 FUNGICIDE/BACTERICIDE	1418.903
		KOCIDE 101	649.880
		KOCIDE 2000	1329.622
		KOCIDE DF	2124.440
		NORDOX 75 WG	834.803
		TRIANGLE BRAND COPPER SULFATE CRYSTAL	217.800
	APPLE	KOCIDE 2000	0.336
		NU-COP 50DF	8.855
	APRICOT	KOCIDE 2000	3.564
		NU-COP 50DF	27.720
	CHERRY	KOCIDE DF	122.800
	GRAPES	BRITZ COPPER SULFUR 15-25 DUST	11101.734
		CHAMP FORMULA 2 FLOWABLE	1372.442
		CHAMPION WETTABLE POWDER	676.160
		CLEAN CROP C-O-C-S 15 SULFUR 25 DUST	498.000
		CSC COPPER SULFUR DUST	1645.512
		DUPONT KOCIDE 2000 FUNGICIDE/BACTERICIDE	1660.537
		DUPONT KOCIDE 3000 FUNGICIDE/BACTERICIDE	938.011
		HYDROX	92.400
		KOCIDE 101	1221.066
		KOCIDE 2000	4876.932
		KOCIDE DF	1715.958
		NORDOX	680.184
		NORDOX 75 WG	1915.020
		NU-COP 50 WP	1.540
		NU-COP 50DF	170.940
		STRETCH FUNGICIDE	0.546
	GRAPES, RAISIN	CHAMP FORMULA 2 FLOWABLE	81.720
		CHAMPION FLOWABLE	8.380
		CHAMPION WETTABLE POWDER	231.000

Chemical Name	Commodity	Product Name	Lbs AI Applied
		CLEAN CROP C-O-C-S 15 SULFUR 25 DUST	2508.000
		COPPER SULFATE CRYSTALS	269.280
		CSC COPPER SULFUR DUST	108.400
		DUPONT KOCIDE 2000 FUNGICIDE/BACTERICIDE	697.679
		DUPONT KOCIDE 3000 FUNGICIDE/BACTERICIDE	1521.028
		DUPONT KOCIDE DF FUNGICIDE/BACTERICIDE	33.402
		KOCIDE 2000	740.330
		KOCIDE DF	42.980
		NORDOX 75 WG	443.344
		NU-COP 50 WP	311.080
		ZINC COPOSIL SULFUR 15-30 DUST	120.000
	GRAPES, WINE	CHAMP FORMULA 2 FLOWABLE	8525.606
		CHAMPION WETTABLE POWDER	1275.505
		CLEAN CROP C-O-C-S 15 SULFUR 25 DUST	1179.000
		CSC COPPER SULFUR DUST	92.140
		DUPONT KOCIDE 2000 FUNGICIDE/BACTERICIDE	1473.998
		DUPONT KOCIDE 3000 FUNGICIDE/BACTERICIDE	1762.458
		DUPONT KOCIDE DF FUNGICIDE/BACTERICIDE	151.290
		KOCIDE 101	726.880
		KOCIDE 2000	4733.920
		KOCIDE 4.5 LF	1142.426
		KOCIDE DF	1624.644
		NORDOX	37.224
		NORDOX 75 WG	248.554
		NU-COP 50 WP	246.400
		NU-COP 50DF	154.000
	STRETCH FUNGICIDE	1.364	
	NECTARINE	DUPONT KOCIDE 2000 FUNGICIDE/BACTERICIDE	267.200
		KOCIDE 2000	288.239
		KOCIDE DF	491.200
		NU-COP 50DF	107.800
	N-OUTDR GRWN TRNSPLNT/PRPGTV MTRL	CHAMP FORMULA 2 FLOWABLE	371.851
	OLIVE (ALL OR UNSPEC)	DUPONT KOCIDE 2000 FUNGICIDE/BACTERICIDE	96.840

Chemical Name	Commodity	Product Name	Lbs AI Applied
	ONION (DRY, SPANISH, WHITE, YELLOW, RED, ETC.)	NORDOX 75 WG	9.095
	ORANGE	BASIC COPPER 53	948.640
		BASIC COPPER SULFATE	693.000
		COP-O-ZINC	231.000
		NORDOX	1453.428
	ORANGE (ALL OR UNSPEC)	BASIC COPPER 53	1051.344
		BLUE SHIELD DF	30.800
		COP-O-ZINC	432.250
		HYDROX	9.625
		KOCIDE 20/20	1697.480
		KOCIDE DF	2101.354
		NORDOX	1103.748
		NU-COP 50DF	1027.565
	PEACH	CHAMP FORMULA 2 FLOWABLE	77.183
		DUPONT KOCIDE 2000 FUNGICIDE/BACTERICIDE	1443.292
		KOCIDE 2000	487.385
		KOCIDE DF	2683.180
		NORDOX 75 WG	201.360
		NU-COP 50DF	266.420
		STRETCH FUNGICIDE	0.682
	PISTACHIO	KOCIDE 2000	242.100
		TRIANGLE BRAND COPPER SULFATE CRYSTAL	79.200
	PLUM	DUPONT KOCIDE 2000 FUNGICIDE/BACTERICIDE	169.470
	PLUM (INCLUDES WILD PLUMS FOR HUMAN CONSUMPTION)	KOCIDE 2000	3.228
		KOCIDE DF	368.400
		NU-COP 50DF	129.360
	PRUNE	KOCIDE 101	77.000
	TANGELO	BASIC COPPER 53	250.880
		COP-O-ZINC	175.000
		KOCIDE 20/20	359.190
		KOCIDE DF	225.952
		NORDOX	591.072
		NU-COP 50DF	215.600
	TANGERINE (MANDARIN, SATSUMA, MURCOTT, ETC.)	CHAMPION WETTABLE POWDER	431.200
	TANGERINE, SEEDL	NORDOX	1316.940
	Unknown	DUPONT KOCIDE 2000 FUNGICIDE/BACTERICIDE	46.806

Chemical Name	Commodity	Product Name	Lbs AI Applied
	WALNUT (ENGLISH WALNUT, PERSIAN WALNUT)	KOCIDE DF	337.700
		NORDOX 75 WG	1152.283
		ALFALFA – Total Pounds Copper Applied	2.421
		ALMOND – Total Pounds Copper Applied	16233.599
		APPLE – Total Pounds Copper Applied	9.191
		APRICOT – Total Pounds Copper Applied	31.284
		CHERRY – Total Pounds Copper Applied	122.800
		GRAPES – Total Pounds Copper Applied	59059.012
		NECTARINE – Total Pounds Copper Applied	1154.439
		N-OUTDR PLANTS – Total Pounds Copper Applied	371.851
		OLIVE – Total Pounds Copper Applied	96.840
		ONION – Total Pounds Copper Applied	9.095
		ORANGE – Total Pounds Copper Applied	10780.233
		PEACH – Total Pounds Copper Applied	5159.502
		PISTACHIO – Total Pounds Copper Applied	321.300
		PLUM (INCLUDING PRUNES) – Total Pounds Copper Applied	747.458
		TANGELO – Total Pounds Copper Applied	1817.694
		TANGERINE – Total Pounds Copper Applied	1748.140
		UNKNOWN – Total Pounds Copper Applied	46.806
		WALNUT – Total Pounds Copper Applied	1489.983
		Total pounds copper applied (2005 - 2009)	99,201.649

The Coalition will continue with its management plan strategy outlined above under the chlorpyrifos outreach section when conducting individual contacts. Orchard and vineyard operators will be advised to consider irrigation water retention and management to prevent copper from entering the waterway with runoff. Storm runoff management relevant to copper applications will also be discussed to prevent the winter exceedances. Individual contacts occurring within this subwatershed are described under the chlorpyrifos outreach section above and include discussions of copper exceedances and the above management practices.

Diuron

Concentrations of diuron have exceeded the WQTL twice in this subwatershed from 2005 through 2009 (January and February 2008). Diuron MPM will begin in January 2010.

PUR data are reviewed for the number of monthly diuron applications, pounds active ingredient (AI) applied, and acres treated (Table V-15, Figure V-5). There has been a general decrease in diuron use since 2006. The amount of diuron used in 2006 was three times the amount of diuron used in 2009 (4,056 lbs in 2006 compared to 966 lbs in 2009, Table V-15, Figure V-5).

Diuron is a general herbicide and is applied throughout the year, generally after high amounts of rain during the winter and/or spring (Table V-15, Figure V-5). None of the exceedances occurred during months of the greatest amount of use, although February was the month with the second highest amount of use in 2008 (Figure V-5).

The highest application rates and largest amount of diuron applied from 2005 through 2009 are associated with grapes (including wine and raisin) and oranges (Table V-16).

Table V-15. Number of diuron applications, total pounds applied and total acres treated by month for January 2005 through December 2009 in the Cottonwood Creek @ Rd 20 site subwatershed. If a month is not included in the table, no applications were made.

Month/Year	Number of Diuron Applications	Pounds Applied	Acres Treated
January, 2005	4	103.7	83.3
February, 2005	19	550.6	473
March, 2005	28	613.7	659.21
April, 2005	12	258.0	328.32
May, 2005	1	9.6	6
October, 2005	3	6.7	109
November, 2005	4	446.4	279
December, 2005	25	827.7	671.4
January, 2006	9	464.7	392.74
February, 2006	37	1140.0	1740.18
March, 2006	17	769.1	545.05
April, 2006	3	100.8	83
November, 2006	16	482.8	957.21
December, 2006	7	314.0	338
January, 2007	7	160.7	370
February, 2007	8	372.9	360.4
March, 2007	16	557.1	556
April, 2007	2	12.8	26
August, 2007	1	8.0	5
September, 2007	2	44.4	806
February, 2008	15	586.7	532.25
March, 2008	17	668.3	647.91
April, 2008	2	296.0	240
November, 2008	16	210.4	1052
January, 2009	12	225.7	339.95
February, 2009	12	295.6	330.34
March, 2009	5	54.4	222
November, 2009	8	359.3	327.7
December, 2009	3	30.8	44

Month/Year	Number of Diuron Applications	Pounds Applied	Acres Treated
Summaries by Year			
2005 Total	96	2816.5	2609.23
2006 Total	89	3271.4	4056.18
2007 Total	36	1155.8	2123.4
2008 Total	50	1761.4	2472.16
2009 Total	40	965.8	1263.99
Total	311	9,970.9	12,524.96

Figure V-5. Pounds of diuron applied within the Cottonwood Creek @ Rd 20 site subwatershed by month for 2005-2009. Asterisk (*) denotes months with exceedances.




Table V-16. Total pounds AI for diuron based on PUR data from 2005-2009 within the Cottonwood Creek @ Rd 20 subwatershed.

Chemical Name	Commodity	Product Name	Lbs AI Applied
DIURON	ALFALFA	DIURON 80 DF HERBICIDE	200.000
		DREXEL DIURON 80 HERBICIDE	244.800
		DUPONT DIREX 4L HERBICIDE	119.977
	ALFALFA (FORAGE - FODDER) (ALFALFA HAY)	DREXEL DIURON 80 HERBICIDE	695.200
	COTTON	ADIOS COTTON DEFOLIANT	44.420
	COTTON, GENERAL	GINSTAR EC COTTON DEFOLIANT	6.722
		DIREX 4L	13.157
	GRAPES	DIREX 80DF	431.872
		DIURON 80 WP WEED KILLER	4.800
		DREXEL DIURON 4L HERBICIDE	11.136
		DREXEL DIURON 80 HERBICIDE	411.288
		DU PONT KARMEX	13.200
		DU PONT KARMEX DF HERBICIDE	533.296
		KARMEX DF HERBICIDE	276.832
		KARMEX XP HERBICIDE	471.736
		GRAPES, RAISIN	DIREX 80DF
	DIURON 4L		33.969
	DIURON 80 DF HERBICIDE		9.600
	DIURON 80 WP WEED KILLER		4.800
	DREXEL DIURON 4L HERBICIDE		103.359
	DREXEL DIURON 80 HERBICIDE		122.696
	DU PONT KARMEX DF HERBICIDE		150.400
	DUPONT KARMEX DF HERBICIDE		98.200
	DUPONT KARMEX XP HERBICIDE		44.808
	KARMEX DF HERBICIDE		24.000
	KARMEX XP HERBICIDE		927.380
	GRAPES, WINE	DIREX 4L	25.422
		DIREX 80DF	90.400
		DIURON 4L	9.146
		DIURON 4L HERBICIDE	6.650
		DIURON 80 DF HERBICIDE	61.600
		DREXEL DIURON 4L HERBICIDE	106.053
		DREXEL DIURON 80 HERBICIDE	127.440
		DU PONT KARMEX	16.000
		DU PONT KARMEX DF HERBICIDE	175.520
		DUPONT KARMEX DF HERBICIDE	66.000
		KARMEX DF HERBICIDE	229.176
		KARMEX XP HERBICIDE	572.760
	OLIVE (ALL OR UNSPEC)	DU PONT KARMEX	12.800
	ORANGE	DIREX 80DF	246.400

Chemical Name	Commodity	Product Name	Lbs AI Applied
		DU PONT KROVAR I DF HERBICIDE	252.800
		DUPONT KARMEX XP HERBICIDE	240.000
		KARMEX DF HERBICIDE	81.600
	ORANGE (ALL OR UNSPEC)	DIREX 80DF	1128.320
		DU PONT KROVAR I DF HERBICIDE	387.200
		KARMEX XP HERBICIDE	153.000
	TANGELO	DU PONT KROVAR I DF HERBICIDE	504.000
		KARMEX XP HERBICIDE	9.600
	TANGERINE (MANDARIN, SATSUMA, MURCOTT, ETC.)	DU PONT KROVAR I DF HERBICIDE	9.600
	TANGERINE, SEEDL	DU PONT KROVAR I DF HERBICIDE	190.800
		DUPONT KARMEX XP HERBICIDE	180.320
	WALNUT (ENGLISH WALNUT, PERSIAN WALNUT)	DIREX 4L	65.036
ALFALFA – Total Pounds Diuron Applied			1259.977
COTTON – Total Pounds Diuron Applied			51.142
GRAPES – Total Pounds Diuron Applied			5198.295
OLIVE – Total Pounds Diuron Applied			12.800
ORANGE – Total Pounds Diuron Applied			2489.32
TANGELO – Total Pounds Diuron Applied			513.600
TANGERINE – Total Pounds Diuron Applied			380.720
WALNUT – Total Pounds Diuron Applied			65.036
Total pounds diuron applied (2005 -2009)			9,970.890

Priority E Constituents

The following priority E constituents are listed under the Cottonwood Creek Management Plan: DO, *E. coli*, and lead. These constituents will remain low priority but will be discussed during individual contacts and annual grower meetings.

Evaluation

Cottonwood Creek @ Rd 20 is one of the second four priority site subwatersheds within the ESJWQC and is in its first year of focused outreach (2010-2012). The Coalition's strategy for the Cottonwood Creek subwatershed is to target growers along or adjacent to the waterway that have the potential to discharge. Focus will be on water retention and minimizing spray drift. Outreach includes grower notification, management practice outreach and education, tracking of management practices implementation, and providing information on special studies of management practice efficacy.

Individual surveys to document current management practices and assess future implementations are in the process of being filled out and will be completed by the end of May 2010. The Coalition will then assess these results and plan the final stages of outreach including re-contacting growers to identify newly implemented practices and future monitoring to evaluate water quality improvement.

VI. DUCK SLOUGH @ GURR RD

Management Plan Constituents

Priority A/B

- Chlorpyrifos

Priority C

- Copper

Priority D

- *Selenastrum capricornutum* water column toxicity
- *Hyalella azteca* sediment toxicity

Priority E

- Dissolved Oxygen
- *E. coli*
- Lead
- pH
- Specific Conductivity
- Total Dissolved Solids
- *Ceriodaphnia dubia* water column toxicity

Description of Duck Slough @ Gurr Rd Site Subwatershed

Duck Slough @ Gurr Road (21,082 irrigated acres) – This site subwatershed is a monitoring location downstream from Duck Slough @ Hwy 99. Located to the south and west of Merced, this site drains field crops immediately upstream and deciduous nuts further upstream as well as some irrigated pasture. The city of Merced delivers treated water to Duck Slough a few miles upstream of the Gurr Road site. Duck Slough flows west eventually becoming Deadman Creek in the western portion of the Coalition region. The slough eventually flows into the San Joaquin River via Deadman Creek and Deep Slough (Figure VI-1). This site subwatershed also includes upstream locations (Duck Slough @ Hwy 59 and North Slough @ Hwy 59) that were sampled in 2008. Table VI-1 includes the station name, station code and target latitude/longitudes for sites sampled within this subwatershed.


Table VI-1. Coordinates of the Duck Slough site subwatershed sampling locations.

Station Name	Station Code	Target Latitude	Target Longitude
Duck Slough @ Gurr Rd*	535XDSAGR	37.2142	-120.5596
Duck Slough @ Hwy 59 ^U	535XDSHFN	37.2345	-120.4881
North Slough @ Hwy 59 ^U	535XNSHFN	37.2277	-120.4880

* Original ESJWQC sampling site

^U Upstream sites

Figure VI-1. Site subwatershed map of land use for the Duck Slough @ Gurr Rd sample site.



Subwatershed Monitoring History

Ambient water monitoring conducted at Duck Slough @ Gurr Rd was initiated during the irrigation season of 2004 and the site has been monitored through 2009 (Table VI-2). Duck Slough @ Gurr Rd will continue to be monitored as a Core Site in 2010 under the current MRPP. The constituents sampled at this location from 2004-2009 are listed in Table VI-3.

Management Plan Monitoring occurred within the Duck Slough @ Gurr Rd subwatershed in 2007 for chlorpyrifos, thiobencarb and copper and in 2008 for copper and *Selenastrum capricornutum* toxicity (Table VI-4 and VI-5). Duck Slough was monitored twice a month in June and July 2007 (Table VI-4). 2008 upstream MPM occurred at Duck Slough @ Hwy 59 for copper in June, for copper and *Selenastrum* toxicity in July, and for *Selenastrum* toxicity in September, and at North Slough @ Hwy 59 for copper and *Selenastrum* toxicity in July and for *Selenastrum* toxicity in September (Table VI-5). The Coalition will conduct MPM for copper (June-July 2010) and toxicity to *Selenastrum* (July 2010).

Starting in January 2010, the Coalition began additional MPM sampling during January, February and March (previously MPM was limited to irrigation months). The Coalition monitored for copper in January and February 2010; these results will be reported on in the Management Plan Update Report to be submitted on April 1, 2011.

Duck Slough @ Gurr Rd is scheduled as a high priority site starting in 2010. Duck Slough is considered impaired under the proposed 2008 Basin Plan 303d and is listed for chlorpyrifos, *E. coli*, sediment toxicity and unknown toxicity. Agriculture is listed as the potential source for the chlorpyrifos impairment; all other impairments have unknown sources.

Table VI-2. Duck Slough @ Gurr Rd sampling events per season and year.

	2004		2005		2006		2007		2008			2009		
	Irrigation	Storm	Irrigation	Storm	Irrigation	Storm	Irrigation	Storm	Irrigation	Fall	Winter	Storm	Irrigation	Fall
Events Sampled	3	2	5	2	5	2	6	2	6	0	2	1	6	3
Events Not Sampled	0	0	0	0	0	0	0	0	0	2	0	1	0	0
Total	3	2	5	2	5	2	6	2	6	2	2	2	6	3

NA indicates that this site was not sampled during this season/year.

Table VI-3. Number of analyses performed per analyte in each sampling season and year for the Duck Slough @ Gurr Rd sample site. Only environmental samples with a sample replicate and lab replicate number of one are shown.

Method	Analyte	2004		2005		2006		2007		2008			2009		
		Irrigation	Storm	Irrigation	Storm	Irrigation	Storm	Irrigation	Storm	Irrigation	Fall	Winter	Storm	Irrigation	Fall
Field and Physical Parameters															
EPA 110.2	Color	3	2	5	2	6	3	6	2	6					
EPA 160.1	Dissolved Solids	3	2	5	2	6	3	6	3	6		2	2	6	2
EPA 160.2	Suspended Solids								1			2	2	6	2
EPA 180.1	Turbidity	3	2	5	2	6	3	6	3	6		2	2	6	2
EPA 405.1	BOD					2	3	2							
EPA 415.1	Total Organic Carbon	2	2	5	1	6	3	6	3	6		2	2	6	2
SM 9223	E. coli	3	2	5	2	6	3	6	3	6		2	2	6	2
NA	Dissolved Oxygen	3	2	5	5	5	3	10	4	8	2	2	2	6	2
NA	Specific Conductivity	3	2	5	5	5	3	10	4	8	2	2	2	6	2
NA	pH	3	2	5	5	5	3	10	4	8	2	2	2	6	2
Carbamates															
EPA 8321A	Aldicarb					5	2	6	2	6					
EPA 8321A	Carbaryl					5	2	6	2	6					
EPA 8321A	Carbofuran					5	2	6	2	6			1	4	2
EPA 8321A	Diuron					5	2	6	2	6					
EPA 8321A	Linuron					5	2	6	2	6					
EPA 8321A	Methiocarb					5	2	6	2	6					
EPA 8321A	Methomyl					5	2	6	2	6					
EPA 8321A	Oxamyl					5	2	6	2	6					
Organochlorines															
EPA 8081A	DDD(p,p')					5	2	6	2	6					
EPA 8081A	DDE(p,p')					5	2	6	2	6					
EPA 8081A	DDT(p,p')					5	2	6	2	6					
EPA 8081A	Dicofol					5	2	6	2	6					
EPA 8081A	Dieldrin					5	2	6	2	6					
EPA 8081A	Endrin					5	2	6	2	6					
EPA 8081A	Methoxychlor					5	2	6	2	6					
Organophosphates															
EPA 8141A	Azinphos methyl					5	2	7	3	6		2	1	2	

Method	Analyte	2004	2005		2006		2007		2008			2009			
		Irrigation	Storm	Irrigation	Storm	Irrigation	Storm	Irrigation	Storm	Irrigation	Fall	Winter	Storm	Irrigation	Fall
EPA 8141A	Chlorpyrifos	3	2	5	2	5	2	7	3	6		2	1	2	
EPA 8141A	Demeton-s								1			2	1	2	
EPA 8141A	Diazinon	3	2	5	2	5	2	7	3	6		2	1	2	
EPA 8141A	Dichlorvos								1			2	1	2	
EPA 8141A	Dimethoate	1				5	2	7	3	6		2	1	2	
EPA 8141A	Disulfoton	1				5	2	7	3	6		2	1	2	
EPA 8141A	Malathion	1				5	2	7	3	6		2	1	2	
EPA 8141A	Methamidophos					5	2	6	3	6		2	1	2	
EPA 8141A	Methidathion					5	2	7	3	6		2	1	2	
EPA 8141A	Molinate					5	2	7	2	6					
EPA 8141A	Parathion, Methyl	1				5	2	7	3	6		2	1	2	
EPA 8141A	Phorate	1				5	2	7	3	6		2	1	2	
EPA 8141A	Phosmet					5	2	7	3	6		2	1	2	
EPA 8141A	Thiobencarb					5	2	9	2	6					
Pyrethroids															
EPA 8081A	Bifenthrin			1	2	5	2	9	2	6					
EPA 8081A	Cyfluthrin, total			1	2	5	2	7	2	6					
EPA 8081A	Cyhalothrin, lambda, total	3	2	5	2	5	2	7	2	6					
EPA 8081A	Cypermethrin, total	3	2	5	2	5	2	7	2	6					
EPA 8081A	Esfenvalerate/Fenvalerate, total	3	2	5	2	5	2	7	2	6					
EPA 8081A	Permethrin, total	3	2	5	2	5	2	7	2	6					
Triazines															
EPA 547M	Glyphosate					5	2	6	2	6					
EPA 549.2M	Paraquat dichloride					5	2	6	2	6					
EPA 619	Atrazine					5	2	6	2	6					
EPA 619	Cyanazine					5	2	6	2	6					
EPA 619	Simazine					5	2	6	2	6					
Metals (Total)															
EPA 200.8	Arsenic					6	3	6	3	6		2	1	1	
EPA 200.8	Boron					6	3	6	3	6		2	1	2	
EPA 200.8	Cadmium					6	3	6	3	6		2	1	1	
EPA 200.8	Copper					6	3	9	3	6		2	1	2	
EPA 200.8	Lead					6	3	6	3	6		2	1	1	

Method	Analyte	2004	2005		2006		2007		2008			2009			
		Irrigation	Storm	Irrigation	Storm	Irrigation	Storm	Irrigation	Storm	Irrigation	Fall	Winter	Storm	Irrigation	Fall
EPA 200.8	Molybdenum								1			2	1	1	
EPA 200.8	Nickel					6	3	6	3	6		2	1	2	
EPA 200.8	Selenium					6	3	2	3	6		2	1	2	
EPA 200.8	Zinc					6	3	7	3	6		2	1	2	
Metals (Dissolved)															
EPA 200.8	Cadmium								1			2	1	1	
EPA 200.8	Copper								1			2	1	2	
EPA 200.8	Lead								1			2	1	1	
EPA 200.8	Nickel								1			2	1	2	
EPA 200.8	Zinc								1			2	1	2	
Nutrients															
SM 2340 C	Hardness as CaCO3					6	3	9	3	6		2	1	2	
EPA 300.0	Nitrate as N					6	3	6	2	6					
EPA 350.2	Ammonia as N					6	3	6	3	6		2	2	6	2
EPA 351.3	Nitrogen, Total Kjeldahl					6	3	6	3	6		2	1	1	
EPA 354.1	Nitrite as N					6	3	6	2	6					
EPA 353.2	Nitrate + Nitrite as N								1			2	2	6	2
EPA 365.2	Orthophosphate as P					6	3	6	3	6		2	1	1	
EPA 365.2	Phosphate as P					6	3	6	3	6		2	2	6	2
Toxicity															
EPA 821/R-02-012	Ceriodaphnia dubia	3	2	5	4	5	3	7	2	7					
EPA 821/R-02-012	Pimephales promelas	3	2	5	2	5	3	7	2	7					
EPA 821/R-02-013	Selenastrum capricornutum	3	2	5	2	5	3	8	2	7					
EPA 600/R-99-064	Hyalella azteca	2		5	1	1	1	1	1	2					

Table VI-4. Duck Slough @ Gurr Rd. 2007 Management Plan additional (A) sampling schedule for chlorpyrifos, thiobencarb and copper. "X" indicates the site, month, and analyte sampled.

Sample Site	Month	Type	Chlorpyrifos	Thiobencarb	Copper
Duck Slough @ Gurr Rd	19-Jun-07	A		X	X
Duck Slough @ Gurr Rd	31-Jul-07	A	X	X	X

Table VI-5. Duck Slough site subwatershed. 2008 Management Plan upstream (U) sampling schedule for chlorpyrifos and copper. "X" indicates the site, month and analyte sampled.

Sample Site	Date	Type	Copper	<i>Selenastrum capricornutum</i> Toxicity
Duck Slough @ Hwy 59	24-Jun-08	U	X	
Duck Slough @ Hwy 59	29-Jul-08	U	X	X
North Slough @ Hwy 59	29-Jul-08	U	X	X
North Slough @ Hwy 59	30-Sep-08	U		X
Duck Slough @ Hwy 59	30-Sep-08	U		X

Exceedance History

During Coalition monitoring, exceedances of WQTLs for field and physical parameters, *E. coli*, nutrients, metals, pesticides, and water column and sediment toxicity occurred within the Duck Slough @ Gurr Rd site subwatershed (Table VI-6). Samples collected over the entire six years of monitoring resulted in exceedances at the normal monitoring and upstream MPM sites, including DO (6), pH (4), SC (3), TDS (1), *E. coli* (19), nitrate + nitrite (91), copper (9), lead (4), Carbofuran (1), chlorpyrifos (1), and thiobencarb (2). Water column toxicity has occurred five times, three times to *Ceriodaphnia dubia* and twice to *Selenastrum capricornutum*. Sediment toxicity to *Hyalella azteca* has occurred five times.

During 2009, exceedances detected at the normal monitoring site included pH (2), specific conductivity (1), *E. coli* (3), nitrate + nitrite (1), and copper (1).

All exceedances are listed in Table VI-6 by season and date and are based on WQTLs listed in the introduction to the ESJWQC Management Plan. A management plan is required if a site experiences two or more exceedances of a particular constituent within three years or if a site has a single exceedance of a constituent that has a TMDL. Using the ESJWQC Management Plan prioritization process flow chart (Figure 4), priorities were assigned to management plan constituents and are listed in the bottom row of Table VI-6. The highest priority constituents in the Duck Slough @ Gurr Road subwatershed are chlorpyrifos (A/B), copper (C), and toxicity to *Selenastrum* and *Hyalella* (D).

Table VI-6. All exceedances experienced in samples collected from locations within the Duck Slough site subwatershed between February 2005 and December 2009 (sorted by season and date). If the water quality trigger limit is based on hardness then the hardness value is shown in parenthesis; otherwise the WQTL used to evaluate the data is listed in the header after the analyte.

Station Name	Season	Sample Date	Oxygen, Dissolved, 7 mg/L	pH, 6.5-8.5 units	Specific Conductivity, 700 µS/cm	<i>E. coli</i> , 235 MPN/100 mL	Dissolved Solids, 450 mg/L	Nitrate + Nitrite as N, 10 mg/L	Copper, Dissolved, µg/L	Copper, Total, 1,300 µg/L	Lead, Total, 15 µg/L	Carbofuran, 0 µg/L	Chlorpyrifos, 0.015 µg/L	Thiobencarb, 0 µg/L	<i>C. dubia</i> , Survival (%)	<i>H. azteca</i> , Survival (%)	<i>S. capricornutum</i> , Total Cell Count
Duck Slough @ Gurr Rd	Irrigation	7/31/2004				350							0.045				
Duck Slough @ Gurr Rd	Irrigation	8/31/2004														33.75	
Duck Slough @ Gurr Rd	Irrigation	9/29/2004			701		540										1220000
Duck Slough @ Gurr Rd	Storm	2/16/2005				1600											
Duck Slough @ Gurr Rd	Storm	3/21/2005				1600											
Duck Slough @ Gurr Rd	Irrigation	5/10/2005				1600											
Duck Slough @ Gurr Rd	Irrigation	6/14/2005				300											
Duck Slough @ Gurr Rd	Irrigation	7/12/2005				300										58.8	
Duck Slough @ Gurr Rd	Irrigation	8/16/2005				240											
Duck Slough @ Gurr Rd	Irrigation	9/20/2005														3.8	
Duck Slough @ Gurr Rd	Storm	2/28/2006													35		
Duck Slough @ Gurr Rd	Storm	3/10/2006													35		
Duck Slough @ Gurr Rd	Storm	3/15/2006				300									40		
Duck Slough @ Gurr Rd	Irrigation	5/17/2006		8.6		2000											
Duck Slough @ Gurr Rd	Irrigation	6/14/2006				690				120 (10.9)			5.8				
Duck Slough @ Gurr Rd	Irrigation	7/12/2006	6.18							14 (9.3)			0.29				
Duck Slough @ Gurr Rd	Irrigation	9/13/2006	5.53														
Duck Slough @ Gurr Rd	Storm	2/12/2007				2400				47 (12.4)	13 (4.88)						
Duck Slough @ Gurr Rd	Storm	2/28/2007				2000				11 (8.8)							
Duck Slough @ Gurr Rd	Storm	3/7/2007		9.17													
Duck Slough @ Gurr Rd	Irrigation	5/29/2007				820											

Station Name	Season	Sample Date	Oxygen, Dissolved, 7 mg/L	pH, 6.5-8.5 units	Specific Conductivity, 700 µS/cm	E. coli, 235 MPN/100 mL	Dissolved Solids, 450 mg/L	Nitrate + Nitrite as N, 10 mg/L	Copper, Dissolved, µg/L	Copper, Total, 1,300 µg/L	Lead, Total, 15 µg/L	Carbofuran, 0 µg/L	Chlorpyrifos, 0.015 µg/L	Thiobencarb, 0 µg/L	C. dubia, Survival (%)	H. azteca, Survival (%)	S. capricornutum, Total Cell Count
Duck Slough @ Gurr Rd	Irrigation	6/19/2007	5.85							5.4 (3.0)							
Duck Slough @ Gurr Rd	Irrigation	6/26/2007								4.6 (3.7)	1 (0.81)						
Duck Slough @ Gurr Rd	Irrigation	7/24/2007															413341
Duck Slough @ Gurr Rd	Irrigation	9/18/2007				370											
Duck Slough @ Gurr Rd	Storm	1/25/2008				>2400				13 (9.0)	2.7 (2.4)						
Duck Slough @ Gurr Rd	Storm	2/25/2008				>2400				17 (9.3)	3.7 (3.2)						
Duck Slough @ Gurr Rd	Irrigation	4/29/2008										0.05					
Duck Slough @ Hwy 59	Irrigation	6/24/2008	4.22		841												
Duck Slough @ Hwy 59	Irrigation	7/29/2008	4.83														
Duck Slough @ Gurr Rd	Sediment	8/28/2008														62	
Duck Slough @ Hwy 59	Irrigation	9/30/2008	3.33														
Duck Slough @ Gurr Rd	Sediment	10/2/2008														90	
Duck Slough @ Gurr Rd	Storm	2/7/2009						13									
Duck Slough @ Gurr Rd	Winter	3/17/2009		9.7													
Duck Slough @ Gurr Rd	Irrigation	5/19/2009				>2400			7.3 (6.12)								
Duck Slough @ Gurr Rd	Irrigation	9/22/2009		9.03													
Duck Slough @ Gurr Rd	Fall	11/17/2009			1215	340											
Duck Slough @ Gurr Rd	Storm	12/15/2009				>2400											
Constituent Priority			E	E	E	E	E	NP	NP	C	E	NP	A/B	NP¹	E	D	D

NP- Not prioritized. Fewer than two exceedances for this constituent at this site within three years and currently no TMDL for constituent
NP¹ - Not Prioritized; thiobencarb is used only by rice and therefore all exceedances are turned over to the Rice Coalition.

2007 - 2009 Management Plan Monitoring Results

Management Plan Monitoring results for Duck Slough @ Gurr Rd as well as normal monitoring results are included in Table VI-7 for 2007 through 2009. Management Plan Monitoring results from 2007 are for thiobencarb, copper and chlorpyrifos and from 2008 are for copper and *Selenastrum capricornutum* toxicity. The Coalition began sampling for both dissolved and total fractions of metals in October 2008; results reported from 2009 are for total and dissolved copper.

2007

In 2007 thiobencarb was being managed by the Coalition and was under the Duck Slough Management Plan along with chlorpyrifos and copper. Duck Slough @ Gurr Rd was monitored monthly for thiobencarb, chlorpyrifos, and copper and additional MPM also occurred for these constituents in June (except for chlorpyrifos) and July. There were no exceedances of chlorpyrifos during the irrigation season of 2007; there were, however, three detections including two in July during normal monitoring and MPM and one in September during normal monitoring (Table VI-7). Copper was detected in all normal monitoring samples collected during the irrigation season. Exceedances of the copper WQTL based on hardness occurred in the MPM and the normal monitoring samples collected in June (Table VI-7). Thiobencarb was not detected in any of the samples collected during the irrigation season.

Duck Slough @ Gurr Rd was sampled both upstream and downstream of the bridge on Gurr Rd on June 19, 2007 to determine if a difference in water quality existed between the two water bodies (Table VI-7). The water upstream of the bridge was dammed and was not flowing downstream although a pipe was discharging water into the pool. Just downstream of this dam is Dean Drain which inputs water from the north. Sampling always occurs downstream of Dean Drain. Both samples collected from Duck Slough on June 16, 2007 had detections of copper and no thiobencarb.

2008

In 2008, normal monitoring at was conducted at Duck Slough @ Gurr Rd as well as MPM at both upstream locations for copper and *Selenastrum capricornutum* toxicity. The upstream location North Slough @ Hwy 59 was dry during both its schedule MPM, in July and September (Table VI-7). Copper was detected in every sample collected during 2008, including from the upstream site Duck Slough @ Hwy 59. There was no toxicity to *Selenastrum* experienced during normal monitoring and MPM.

2009

Monitoring for both total and dissolved copper occurred at Duck Slough @ Gurr Rd April and May 2009. Copper was detected in both the total and dissolved copper samples in April and

exceeded the WQTL and hardness based WQTL for both total and dissolved copper, respectively, in May. Duck Slough @ Gurr Rd was not scheduled for 2009 MPM.

Management Plan Monitoring for copper, chlorpyrifos, *Selenastrum* toxicity, and *Hyalella* toxicity are planned for the 2010 irrigation season. Management Plan Monitoring during January, February, and March 2011 will also occur for copper and toxicity to *Ceriodaphnia*.

Table VI-7. Duck Slough @ Gurr Rd. Normal monitoring (NM) and Management Plan Monitoring (MPM) results where 'A' indicates additional MPM (2007) and 'US' indicates upstream monitoring for chlorpyrifos, copper and thiobencarb from the 2007 irrigation season. NM and MPM (2008) for copper and *Selenastrum*. NM for copper (2009). Exceedance values are in bold.

Month:		April	May	June	July	August	September
2007 NM (@ Gurr Rd)	Date	4/24/07	5/29/07	6/26/07	7/24/07	8/21/07	9/18/07
	Chlorpyrifos (µg/L)	<0.003	<0.003	<0.003	0.007	<0.003	0.009
	Copper (µg/L)	4.6	5.3	4.6	4	4.8	5.5
	Thiobencarb (µg/L)	<0.06	<0.06	<0.06	<0.06	<0.06	<0.06
2007 MPM A (@ Gurr Rd)	Date	NA	NA	6/19/07	6/19/07 ^{US}	7/31/07	NA
	Chlorpyrifos (µg/L)	NA	NA	NA	NA	0.007	NA
	Copper (µg/L)	NA	NA	5.4	15	8.6	NA
	Thiobencarb (µg/L)	NA	NA	<0.06	<0.06	<0.06	NA
2008 NM (@ Gurr Rd)	Date	4/29/08	5/27/08	6/24/08	7/29/08	8/26/08	9/30/08
	Copper (µg/L)	2.7	7.1	4.0	5.4	3.5	6.1
	<i>S. capricornutum</i> toxicity (% Control)	175	309	459	350	262	263
2008 MPM US (@ Hwy 59^{US})	Date	NA	NA	6/24/08	7/29/08	NA	9/30/08
	Copper (µg/L)	NA	NA	12	18	NA	NA
	<i>S. capricornutum</i> toxicity (% Control)	NA	NA	NA	271	NA	131
2008 MPM US (North Slough @ Hwy 59^{US})	Date	NA	NA	NA	7/29/08	NA	9/30/08
	Copper (µg/L)	NA	NA	NA	Dry	NA	Dry
	<i>S. capricornutum</i> toxicity (% Control)	NA	NA	NA	Dry	NA	Dry
2009 NM (@ Gurr Rd)	Date	4/21/09	5/19/09	6/16/09	7/21/09	8/18/09	9/22/09
	Copper, total (µg/L)	3.3	15	NA	NA	NA	NA
	Copper, dissolved (µg/L)	2.6	7.3	NA	NA	NA	NA

^UUpstream of bridge on Gurr Rd

NA - Not applicable. This site was not sampled during this month.

Load Calculations

Loads have been calculated for all chlorpyrifos detections in the site subwatershed (Table VI-8) based on the following formula:

$$\text{Load} = \text{Discharge (cfs)} \times 28.317\text{L/ft}^3 \times \text{Concentration (milligram/L} \times 1000 \text{ or } \mu\text{g/L)}.$$

The load values for constituents in this report represent instantaneous loads only. These values should not be used to extrapolate loading over any period of time (e.g. weekly, monthly, seasonal or annual). The primary purpose for reporting instantaneous loads is to provide a normalization of the concentrations by flow for various constituents at the time the samples were collected.

Table VI-8. Duck Slough site subwatershed. Instantaneous load calculations for chlorpyrifos and copper where discharge was measured (sorted by analyte, site and date).

Station Name	Analyte Name	Sample Date	Discharge cfs	Concentration $\mu\text{g/L}$	Loading Rate $\mu\text{g/sec}$
Duck Slough @ Gurr Rd	Chlorpyrifos	8-Aug-06	6.62	0.014	2.62
Duck Slough @ Gurr Rd	Chlorpyrifos	24-Jul-07	1.78	0.007	0.35
Duck Slough @ Gurr Rd	Chlorpyrifos	31-Jul-07	3.25	0.007	0.64
Duck Slough @ Gurr Rd	Chlorpyrifos	18-Sep-07	5.2	0.009	1.33
Duck Slough @ Gurr Rd	Chlorpyrifos	29-Jul-08	0.4	0.011	0.12
Duck Slough @ Gurr Rd	Chlorpyrifos	20-Jan-09	0.23	0.012	0.08
Duck Slough @ Gurr Rd	Copper	17-May-06	2.29	2.9	188.05
Duck Slough @ Gurr Rd	Copper	14-Jun-06	1.61	120	5470.84
Duck Slough @ Gurr Rd	Copper	12-Jul-06	10.93	14	4333.07
Duck Slough @ Gurr Rd	Copper	8-Aug-06	6.62	3.8	712.34
Duck Slough @ Gurr Rd	Copper	13-Sep-06	0.96	4.1	111.46
Duck Slough @ Gurr Rd*	Copper	13-Sep-06	0.96	4	108.74
Duck Slough @ Gurr Rd*	Copper	13-Sep-06	0.96	4	108.74
Duck Slough @ Gurr Rd	Copper	12-Feb-07	298.79	47	397659.31
Duck Slough @ Gurr Rd*	Copper	12-Feb-07	298.79	46	389198.48
Duck Slough @ Gurr Rd*	Copper	12-Feb-07	298.79	44	372276.80
Duck Slough @ Gurr Rd	Copper	28-Feb-07	321.45	11	100127.50
Duck Slough @ Gurr Rd	Copper	24-Apr-07	1.88	4.6	244.89
Duck Slough @ Gurr Rd	Copper	29-May-07	1.02	5.3	153.08
Duck Slough @ Gurr Rd	Copper	19-Jun-07	0.99	5.4	151.38
Duck Slough @ Gurr Rd	Copper	19-Jun-07	0	15	0
Duck Slough @ Gurr Rd	Copper	26-Jun-07	1.12	4.6	145.89
Duck Slough @ Gurr Rd	Copper	24-Jul-07	1.78	4	201.62
Duck Slough @ Gurr Rd	Copper	31-Jul-07	3.25	8.6	791.46
Duck Slough @ Gurr Rd	Copper	21-Aug-07	1.38	4.8	187.57
Duck Slough @ Gurr Rd	Copper	18-Sep-07	5.2	5.5	809.87

Station Name	Analyte Name	Sample Date	Discharge cfs	Concentration µg/L	Loading Rate µg/sec
Duck Slough @ Gurr Rd*	Copper	18-Sep-07	5.2	6.4	942.39
Duck Slough @ Gurr Rd	Copper	29-Apr-08	0.56	2.7	42.82
Duck Slough @ Gurr Rd*	Copper	27-May-08	0.43	7.3	88.89
Duck Slough @ Gurr Rd	Copper	27-May-08	0.43	7.1	86.45
Duck Slough @ Gurr Rd	Copper	24-Jun-08	0.65	4	73.62
Duck Slough @ Gurr Rd	Copper	29-Jul-08	0.4	5.4	61.16
Duck Slough @ Gurr Rd	Copper	26-Aug-08	1.1	3.5	109.02
Duck Slough @ Gurr Rd	Copper	30-Sep-08	0.21	6.1	36.27
Duck Slough @ Gurr Rd	Copper (Dissolved)	16-Dec-08	0.31	0.3	2.63
Duck Slough @ Gurr Rd	Copper	16-Dec-08	0.31	1.2	10.53
Duck Slough @ Gurr Rd	Copper (Dissolved)	20-Jan-09	0.23	0.2	1.30
Duck Slough @ Gurr Rd	Copper	20-Jan-09	0.23	0.4	2.61
Duck Slough @ Gurr Rd	Copper (Dissolved)	7-Feb-09	2.57	7.6	553.09
Duck Slough @ Gurr Rd	Copper	7-Feb-09	2.57	11	800.52
Duck Slough @ Gurr Rd	Copper (Dissolved)	17-Mar-09	0	4.6	0
Duck Slough @ Gurr Rd	Copper	17-Mar-09	0	6.3	0
Duck Slough @ Gurr Rd	Copper	21-Apr-09	0.67	3.3	62.61
Duck Slough @ Gurr Rd	Copper (Dissolved)	21-Apr-09	0.67	2.6	49.33
Duck Slough @ Gurr Rd	Copper (Dissolved)	19-May-09	0.19	7.3	39.28
Duck Slough @ Gurr Rd	Copper	19-May-09	0.19	15	80.70
<i>Duck Slough @ Hwy 59</i>	<i>Copper</i>	<i>24-Jun-08</i>	<i>0.64</i>	<i>12</i>	<i>217.47</i>
<i>Duck Slough @ Hwy 59</i>	<i>Copper</i>	<i>29-Jul-08</i>	<i>0.47</i>	<i>18</i>	<i>239.56</i>

*Field Duplicate

Source Identification and Outreach

Priority A/B Constituents

Chlorpyrifos is the only priority A/B constituent listed under the Duck Slough @ Gurr Rd Management Plan.

Chlorpyrifos

Chlorpyrifos has exceeded the WQTL of 0.015µg/L once in this subwatershed from 2004 through 2009 (July, 2004). The Coalition uses a combination of monitoring data and evaluation of Pesticide Use Report (PUR) data to identify possible sources.

PUR data are reviewed for the number of monthly chlorpyrifos applications, pounds active ingredient (AI) applied, and acres treated (Table VI-9, Figure VI-2). The amount of chlorpyrifos applied within the subwatershed has fluctuated over the years. The highest chlorpyrifos use occurred in 2005 (5,383 lbs), and the lowest use was in 2004 (1,214 lbs); in 2009, 1,820 lbs were

applied (Table VI-9, Figure VI-2). In 2006, 2008, and 2009 the greatest amount of chlorpyrifos was applied in May, whereas the greatest amount was applied in August of 2004, July of 2005, and March of 2007 (Table VI-9, Figure VI-2). Exceedances did not always occur during months of the greatest amount of use (Figure VI-2).

The Coalition also used PUR data to assess which crops receive the most applications of chlorpyrifos. The highest application rates are associated with alfalfa, almonds, and corn (Table VI-10). The largest amount of chlorpyrifos applied from 2004 through 2009 was to almonds, followed by alfalfa and corn (Table VI-10). The most common product containing chlorpyrifos applied within the subwatershed was Lorsban (Table VI-10).

Table VI-9. Number of chlorpyrifos applications, total pounds applied and total acres treated by month for January 2004 through December 2009 in the Duck Slough @ Gurr Road site subwatershed. If a month is not included in the table, no applications were made.

Month/Year	Number of Chlorpyrifos Applications	Pounds Applied	Acres Treated
January, 2004	4	88.0	43.6
April, 2004	7	94.1	222
May, 2004	1	77.9	60
June, 2004	2	152.1	76
July, 2004	6	333.0	214
August, 2004	8	403.2	544
September, 2004	2	66.6	66
January, 2005	4	76.9	38.1
March, 2005	18	585.0	984.5
June, 2005	5	209.0	188
July, 2005	18	4282.9	868
August, 2005	4	131.5	260.5
September, 2005	2	97.9	105
January, 2006	6	119.7	59
March, 2006	5	105.9	336
May, 2006	13	706.5	433
June, 2006	1	137.6	69
July, 2006	6	585.1	528
August, 2006	9	402.2	448
September, 2006	8	425.9	413.1
January, 2007	2	29.9	15
March, 2007	40	999.4	2112.61
April, 2007	3	28.8	57.42
June, 2007	1	19.9	10
July, 2007	3	139.0	185
March, 2008	26	416.7	846.53

Month/Year	Number of Chlorpyrifos Applications	Pounds Applied	Acres Treated
May, 2008	14	1015.3	669
June, 2008	1	10.2	8
July, 2008	10	339.1	739.3
August, 2008	23	632.7	1283.2
September, 2008	2	34.6	98
March, 2009	14	298.6	754
April, 2009	1	31.7	42.2
May, 2009	3	815.4	624
July, 2009	10	370.5	563
August, 2009	7	246.7	403
September, 2009	4	57.2	227.4
Summaries by Year			
2004 Total	30	1214.8	1225.6
2005 Total	51	5383.2	2444.1
2006 Total	48	2482.8	2286.1
2007 Total	49	1216.9	2380.03
2008 Total	76	2448.5	3644.03
2009 Total	39	1820.1	2613.6
Total	293	14,566.3	14,593.46

Figure VI-2. Pounds of chlorpyrifos applied within the Duck Slough @ Gurr Rd site subwatershed by month for 2004-2009. Asterisk (*) denotes months with exceedances.




Table VI-10. Total pound AI for chlorpyrifos based on PUR data from 2004-2009 within the Duck Slough @ Gurr Rd subwatershed.

Chemical Name	Commodity	Product Name	Lbs AI Applied
CHLORPYRIFOS	ALFALFA	COBALT	139.989
		GOVERN 4E INSECTICIDE	807.143
		LOCK-ON INSECTICIDE	565.502
		LORSBAN 4E-HF	574.758
		LORSBAN-4E	559.166
		NUFOS 4E	191.737
	ALFALFA (FORAGE - FODDER) (ALFALFA HAY)	LOCK-ON INSECTICIDE	54.801
		LORSBAN 4E INSECTICIDE	63.921
		LORSBAN 4E-HF	420.474
		LORSBAN 4E-HF INSECTICIDE	63.435
		LORSBAN-4E	26.133
		NUFOS 4E	281.380
	ALMOND	CHLORPYRIFOS 4E AG	129.830
		GOVERN 4E INSECTICIDE	498.956
		LORSBAN 4E INSECTICIDE	175.710
		LORSBAN 4E-HF	1480.635
		LORSBAN-4E	3619.113
		NUFOS 4E	20.335
		WHIRLWIND	379.940
	CORN (FORAGE - FODDER)	CHLORPYRIFOS 4E AG	116.990
		GOVERN 4E INSECTICIDE	85.853
		LORSBAN 15G GRANULAR INSECTICIDE	77.852
		LORSBAN 4E-HF	161.382
		NUFOS 4E	100.660
	CORN FOR/FOD	GOVERN 4E INSECTICIDE	202.007
		LORSBAN 15G GRANULAR INSECTICIDE	1781.326
		LORSBAN 4E-HF	19.939
		LORSBAN-4E	31.666
		NUFOS 4E	100.660
	COTTON	NUFOS 4E	13.218
COTTON, GENERAL	CHLORPYRIFOS 4E AG	246.170	
	GOVERN 4E INSECTICIDE	59.087	
	LORSBAN 4E-HF	89.589	
PEACH	CHLORPYRIFOS 4E AG	164.852	
	LORSBAN 4E-HF	121.625	
SORGHUM FOR/FOD	GOVERN 4E INSECTICIDE	50.502	
SUGARBEET, GENERAL	LORSBAN 4E-HF	69.865	
	LORSBAN-4E	215.107	
WALNUT	GOVERN 4E INSECTICIDE	15.150	
	LORSBAN-4E	352.616	
WALNUT (ENGLISH WALNUT, PERSIAN WALNUT)	LORSBAN 4E-HF	71.779	
	LORSBAN-4E	365.468	

Chemical Name	Commodity	Product Name	Lbs AI Applied
		ALFALFA – Total Pounds Chlorpyrifos Applied	3748.437
		ALMOND – Total Pounds Chlorpyrifos Applied	6304.519
		CORN – Total Pounds Chlorpyrifos Applied	2678.334
		COTTON– Total Pounds Chlorpyrifos Applied	408.064
		PEACH – Total Pounds Chlorpyrifos Applied	286.477
		SORGHUM – Total Pounds Chlorpyrifos Applied	50.502
		SUGARBEET – Total Pounds Chlorpyrifos Applied	284.971
		WALNUT – Total Pounds Chlorpyrifos Applied	805.013
Total pounds chlorpyrifos applied (2004 - 2009)			14,566.317

Based on results from general management practice surveys, approximately 23% of the member acreage have no runoff occurs, and 36% of the member acreage uses recirculation and/or tailwater return systems. Because the highest applications of chlorpyrifos occur to almonds, alfalfa and corn, the Coalition believes that outreach to members applying chlorpyrifos and owning parcels with the potential to drain to the creek will result in increased implementation of irrigation water management as well as practices to minimize spray drift which will yield improvements in water quality. Growers are encouraged to review their operation to determine if irrigation return flows are managed properly and owners operating orchards should evaluate their aerial applications to minimize spray drift.

Coalition outreach since 2007 has included grower meetings and the mailing/distribution of information. A complete list of Coalition Outreach during 2009 is provided in the Summary of Coalition Outreach Activities section of the Management Plan Update Report. The Coalition has begun the process of contacting growers, conducting individual meetings, and compiling individual surveys for Duck Slough. The Coalition focused outreach to growers owning parcels with the potential to drain to the creek and those applying chlorpyrifos. Using these criteria, a list of 6 targeted growers was created. Individual meetings are currently being scheduled and will be completed by the end of May 2010. Individual contacts will focus mainly on chlorpyrifos exceedances however all water quality results will be reviewed and discussed including copper, DO, *E. coli*, pH, SC, TDS, and lead WQTLs exceedances as well as toxicity to *Ceriodaphnia dubia*, *Selenastrum capricornutum*, and *Hyalella azteca*.

The Coalition will compile and analyze the surveys by the end of August 2010. Based on the exact timing of each individual meeting and/or a grower's resources, some owners will not be able to implement recommended management practices until the 2011 irrigation season. In addition, long term structural BMPs (e.g. sediment ponds) will most likely take longer than two years to implement and will require additional tracking to document their implementation. The Coalition will follow up with growers between February and April of 2011 to determine what practices were implemented during the first irrigation year. Growers who have not already been contacted or have indicated that they have not yet implemented practices during the first year will be contacted from February to April of 2012 to determine what practices were implemented during the dormant season and/or the second irrigation year. To evaluate the management practice process, the Coalition will monitor for high priority constituents during

months of past exceedances in 2010 and 2011. Depending on when additional management practices are implemented, the Coalition may monitor through 2012 to evaluate improvements in water quality. The Coalition anticipates that changes in management practices by members that have direct drainage and/or are have the potential for spray drift will affect downstream water quality by the end of the second year as a high priority site subwatershed.

The effectiveness of the management practices will be determined through the monitoring of water quality in the years following implementation. However, due to the presence of dairies and nonmembers in the subwatersheds, implementation of management practices by only coalition members may not result in improved water quality. If water quality fails to improve, the Coalition will identify parcels from dairies and nonmembers that could contribute to exceedances and provide that information to the Regional Board.

Priority C Constituents

Duck Slough @ Gurr Rd is listed for copper.

Copper

Copper has exceeded the WQTL times in this subwatershed from 2006 through 2008. As of October 2008, the Coalition began to analyze for both total and dissolved fractions of copper to better characterize the subwatershed. Total copper exceedances occurred during June and July 2006, February and June 2007, and in January and February 2008, while the single dissolved copper exceedance occurred in May 2009 (Table VI-6).

PUR data are reviewed for the number of monthly copper applications, pounds active ingredient (AI) applied, and acres treated (Table VI-11, Figure VI-3). The amount of copper applied within the subwatershed has decreased since 2005 (Table VI-11, Figure VI-3). The largest amount of copper applied was in the year of 2005 (12,952 lbs) while the smallest amount applied occurred in 2009 (2,644 lbs). In 2004, 2006, and 2007, the greatest amount of copper use occurred in January, whereas the greatest use amount occurred in December during 2008 and 2009; the greatest amount of copper use occurred in June in 2005 (Table VI-11, Figure VI-3). Not all exceedances occurred during months of the greatest amount of use (Figure VI-3).

The Coalition also used PUR data to assess which crops receive the most applications of copper. The largest amount of copper applied from 2004 through 2009 was associated with almond, peach, rice, tomato and walnut crops (Table VI-12). The most common products containing copper applied withing the subwatershed were DuPont Kocide and Kocide (Table VI-12).

Table VI-11. Number of copper applications, total pounds applied and total acres treated by month for January 2004 through December 2009 in the Duck Slough @ Gurr Rd site subwatershed. If a month is not included in the table, no applications were made.

Month/Year	Number of Copper Applications	Pounds Applied	Acres Treated
January, 2004	21	2826.7	707.6
February, 2004	6	428.8	194
March, 2004	3	758.7	135
April, 2004	4	86.0	37.25
August, 2004	3	214.3	191.2
September, 2004	1	263.4	143
October, 2004	4	1328.3	575
November, 2004	2	550.1	224
January, 2005	12	1568.7	376.1
February, 2005	5	514.8	159.7
March, 2005	3	369.9	81
April, 2005	26	1363.2	483.25

Month/Year	Number of Copper Applications	Pounds Applied	Acres Treated
May, 2005	12	641.9	414.5
June, 2005	10	7053.8	475
August, 2005	3	139.8	127.8
September, 2005	2	82.4	74.7
October, 2005	1	25.8	21
November, 2005	1	69.1	45
December, 2005	5	1122.8	153
January, 2006	12	1872.3	321
February, 2006	9	814.0	251
March, 2006	4	288.0	360
April, 2006	12	1038.3	263.5
May, 2006	14	1197.8	613.7
June, 2006	1	105.2	100
August, 2006	1	113.0	103
October, 2006	2	255.6	166
December, 2006	2	539.0	70
January, 2007	9	1502.7	223
February, 2007	8	533.7	280
March, 2007	3	309.2	115
April, 2007	10	422.9	147.5
May, 2007	2	145.0	163
January, 2008	6	826.5	136
February, 2008	2	650.3	42
March, 2008	1	8.1	5
April, 2008	2	108.3	65
May, 2008	5	318.9	165.5
June, 2008	1	17.9	25.84
September, 2008	1	42.4	92
November, 2008	1	166.0	60
December, 2008	3	901.0	100
January, 2009	2	847.7	102
February, 2009	1	35.6	58
April, 2009	12	596.1	294.5
May, 2009	5	157.9	125.5
August, 2009	1	69.3	30
November, 2009	1	77.0	10
December, 2009	4	861.0	73
Summaries by Year			
2004 Total	44	6456.4	2207.05
2005 Total	80	12952.2	2411.05

Month/Year	Number of Copper Applications	Pounds Applied	Acres Treated
2006 Total	57	6223.3	2248.2
2007 Total	32	2913.6	928.5
2008 Total	22	3039.2	691.34
2009 Total	26	2644.6	693
Total	261	34,229.3	9,179.14

Figure VI-3. Pounds of copper applied within the Duck Slough @ Gurr Rd site subwatershed by month for 2004-2009. Asterisk (*) denotes months with exceedances.




Table VI-12. Total pound AI for copper based on PUR data from 2004-2009 within the Duck Slough @ Gurr Rd subwatershed.

Chemical Name	Commodity	Product Name	Lbs AI Applied	
COPPER	ALFALFA (FORAGE - FODDER) (ALFALFA HAY)	KOCIDE 2000	63.753	
		BASIC COPPER 53	220.500	
	ALMOND	BASICOP	307.400	
		CHAMPION WETTABLE POWDER	80.080	
		COOKE BLUESTONE	8.250	
		CUPROFIX ULTRA 40 DISPERS	602.217	
		DUPONT KOCIDE 2000 FUNGICIDE/BACTERICIDE	117.822	
		DUPONT KOCIDE 3000 FUNGICIDE/BACTERICIDE	67.076	
		DUPONT KOCIDE DF FUNGICIDE/BACTERICIDE	35.612	
		KOCIDE 101	887.040	
		KOCIDE 2000	750.510	
		KOCIDE DF	1899.716	
		KOP-HYDROXIDE 50W	776.160	
		NORDOX 75 WG	1969.972	
		APRICOT	BASIC COPPER 53	58.800
			KOCIDE 2000	118.360
	KOCIDE DF		122.800	
	CHERRY	NU-COP 50DF	154.000	
	NECTARINE	BASIC COPPER 53	98.000	
		CHAMPION WETTABLE POWDER	24.063	
		DUPONT KOCIDE 2000 FUNGICIDE/BACTERICIDE	42.368	
		DUPONT KOCIDE 3000 FUNGICIDE/BACTERICIDE	23.050	
		KOCIDE 101	1155.000	
		KOCIDE 2000	201.750	
	N-OUTDOOR PLANT	BASIC COPPER 53	1519.000	
		DUPONT KOCIDE 3000 FUNGICIDE/BACTERICIDE	331.920	
	N-OUTDR CONTAINER/FLD GRWN PLANTS	KOCIDE 101	473.550	
	PEACH	BASIC COPPER 53	2528.400	
		DUPONT KOCIDE 2000 FUNGICIDE/BACTERICIDE	172.160	
		KOCIDE 101	154.000	
		KOCIDE DF	788.744	
PEPPERS (FRUITING VEGETABLE), (BELL, CHILI, ETC.)	KOCIDE 101	121.275		
	KOCIDE DF	27.016		
PISTACHIO (PISTACHE NUT)	CUPROFIX DISPERS DRY FLOWABLE FUNGICIDE/BACTERICIDE FOR	8.303		

Chemical Name	Commodity	Product Name	Lbs AI Applied
		TURF AND ORNAMENTAL USE	
	PLUM	DUPONT KOCIDE 2000 FUNGICIDE/BACTERICIDE	80.700
	PLUM (INCLUDES WILD PLUMS FOR HUMAN CONSUMPTION)	KOCIDE 101	154.000
	PRUNE	KOCIDE DF	110.520
		NORDOX 75 WG	1056.133
	RICE	COPPER-Z 4/4	136.770
	RICE (ALL OR UNSPEC)	COPPER-Z 4/4	355.287
		TRIANGLE BRAND COPPER SULFATE CRYSTAL	7053.750
	TOMATO	BLUE SHIELD WP	127.820
		CHAMPION WETTABLE POWDER	51.398
		DUPONT KOCIDE 2000 FUNGICIDE/BACTERICIDE	19.368
		DUPONT KOCIDE 3000 FUNGICIDE/BACTERICIDE	85.861
		KOCIDE 101	422.153
		KOCIDE 2000	1011.687
		KOCIDE DF	892.449
		NU-COP 50 WP	127.820
		NU-COP 50DF	1328.250
	TOMATO PROCESSING	CHAMPION WETTABLE POWDER	69.300
		DUPONT KOCIDE 3000 FUNGICIDE/BACTERICIDE	67.541
	TOMATOES, FOR PROCESSING/CANNING	CHAMPION WETTABLE POWDER	34.650
		KOCIDE 2000	29.052
	WALNUT	DUPONT GX-569 FUNGICIDE/BACTERICIDE	212.982
		DUPONT KOCIDE 2000 FUNGICIDE/BACTERICIDE	503.568
		DUPONT KOCIDE 3000 FUNGICIDE/BACTERICIDE	436.106
		KOCIDE 101	309.232
		KOCIDE 2000	350.238
	WALNUT (ENGLISH WALNUT, PERSIAN WALNUT)	DUPONT KOCIDE 2000 FUNGICIDE/BACTERICIDE	167.453
		KOCIDE 101	1410.640
		KOCIDE 2000	1168.267
		KOP-HYDROXIDE 50W	396.550
		NORDOX 75 WG	151.020
ALFALFA – Total Pounds Copper Applied			63.753
ALMOND – Total Pounds Copper Applied			7722.355
APRICOT – Total Pounds Copper Applied			299.960
CHERRY – Total Pounds Copper Applied			154.000
NECTARINE – Total Pounds Copper Applied			1544.230

Chemical Name	Commodity	Product Name	Lbs AI Applied
	N-OUTDOOR PLANT – Total Pounds Copper Applied		2324.47
	PEACH – Total Pounds Copper Applied		3643.304
	PEPPERS – Total Pounds Copper Applied		148.291
	PISTACHIO – Total Pounds Copper Applied		8.303
	PLUM (INCLUDES PRUNES) – Total Pounds Copper Applied		1401.353
	RICE – Total Pounds Copper Applied		7545.807
	TOMATO – Total Pounds Copper Applied		4267.349
	WALNUT – Total Pounds Copper Applied		5106.056
Total pounds copper applied (2004 - 2009)			34,229.230

The Coalition will continue with its management plan strategy outlined above under the chlorpyrifos outreach section when conducting individual contacts. Orchard and vineyard operators will be advised to consider irrigation water retention and management to prevent copper from entering the waterway with runoff. Storm runoff management relevant to copper applications will also be discussed to prevent the winter exceedances. Individual contacts occurring within this subwatershed are described under the chlorpyrifos outreach section above and will include discussions of copper exceedances and the above management practices.

Priority D Constituents

The following priority D constituents are listed under the Duck Slough @ Gurr Road Management Plan: *Selenastrum capricornutum* and sediment toxicity to *Hyalella azteca*. Toxicity was not experienced at all during 2009 monitoring. These constituents will remain low priority but will be discussed during individual contacts and annual grower meetings.

***Selenastrum* toxicity**

Toxicity to *Selenastrum* occurred in September 2004 and July 2007. The July toxicity is associated with a copper exceedance that occurred in the prior month, but toxicity was not persistent in the resample on July 31, 2007.

The Coalition’s strategy for eliminating *Selenastrum* toxicity involves focusing on copper. If copper can be prevented from entering waterways throughout the irrigation season, the Coalition believes that *Selenastrum* toxicity can also be reduced or eliminated. If copper is no longer detected in the slough but *Selenastrum* toxicity persists, the Coalition will consider other herbicides.

***Hyalella* toxicity**

Sediment toxicity to *Hyalella* occurred in August 2004, July 2005, and in August 2008. Toxicity was persistent in the 2008 resample that occurred in October 2008. Toxicity was not experienced during 2009 sampling.

The Coalition has initiated management plan sampling for sediment toxicity. As of 2009, the Coalition now tests for total organic carbon and grain size in all sediments and pyrethroids and chlorpyrifos in sediment that exhibits toxicity. Management Plan Monitoring is scheduled for this site in 2010.

Priority E Constituents

The following priority E constituents are listed under the Duck Slough @ Gurr Road Management Plan: DO, pH, SC (new as of 2009 sampling), TDS, *E. coli*, lead, and water column toxicity to *Ceriodaphnia dubia*. Exceedances of pH, SC, and *E. coli* continued in 2009.

Toxicity to *Ceriodaphnia* has not been associated with the presence of pesticides or metals in the past. The Coalition did not perform testing for *Ceriodaphnia* toxicity in 2009. The Coalition will monitor this site for *Ceriodaphnia* toxicity during March 2011.

The toxicity and other constituents will remain low priority but will be discussed during individual contacts and annual grower meetings.

Evaluation

Duck Slough @ Gurr Rd is one of the second four priority site subwatersheds within the ESJWQC and is in its first year of focused outreach (2010-2012). The Coalition's strategy for the Duck Slough subwatershed has been to target growers along or adjacent to the waterway that have the potential to discharge. Focus will be on tail water management and minimizing spray drift. Outreach includes grower notification, management practice outreach and education, tracking of management practices implementation, and providing information on special studies of management practice efficacy.

Individual surveys to document current management practice and assess future planned implementations will be completed by the end of May 2010. The Coalition will then assess these results and plan the final stages of outreach including re-contacting growers to identify newly implemented practices and future monitoring to evaluate water quality improvement.

VII. HIGHLINE CANAL @ HWY 99

Management Plan Constituents

Priority A/B

- Chlorpyrifos

Priority C

- Copper
- Diuron

Priority D

- *Ceriodaphnia dubia* water column toxicity
- *Selenastrum capricornutum* water column toxicity
- *Hyalella azteca* sediment toxicity

Priority E

- Ammonia
- pH
- Lead
- Specific Conductivity
- Total Dissolved Solids
- *E. coli*


Description of Highline Canal @ Hwy 99 Site Subwatershed

Highline Canal @ Highway 99 (35,220 irrigated acres) – The Highline Canal is a conveyance of the Turlock Irrigation District (TID) and carries both clean irrigation water and irrigation return flow during the summer, and storm water runoff during the winter. This site was selected as a downstream companion site to the Highline Canal @ Lombardy Road site. This site subwatershed is monitored to determine the relative contribution of the upstream and downstream site subwatersheds to water quality impairments. The sampling site is located just south of Delhi as the canal crosses the highway. The irrigated agriculture is primarily deciduous nuts, and these are located at the lower end of the site subwatershed. A small number of vineyards are also present (Figure VII-1). The sample location for Highline Canal @ Hwy 99 is provided in Table VII-1.

Table VII-1. Coordinates of the Highline Canal site subwatershed sampling location.

Station Name	Station Code	Target Latitude	Target Longitude
Highline Canal @ Hwy 99	535XHCHNN	37.4153	-120.7557

Figure VII-1. Site subwatershed map of land use for the Highline Canal @ Hwy 99 sample site.



Subwatershed Monitoring History

Ambient water monitoring conducted at the Highline Canal @ Highway 99 site was initiated in the 2005 irrigation season and monitoring has continued through 2009 (Table VII-2). Highline Canal @ Hwy 99 will remain a Core Monitoring location under the current MRPP until 2011 when it will rotate to an Assessment Monitoring location. Specific information on the analysis conducted across each of the monitoring seasons is provided below (Table VII-3).

Management Plan Monitoring occurred at Highline Canal during 2007 and 2008 for *Ceriodaphnia dubia* toxicity only in 2007 then for *Ceriodaphnia dubia* toxicity, chlorpyrifos, and copper in 2008 (Table VII-4, Table VII-5). The site was also monitored for these constituents monthly April through September in 2007 and 2008. Management Plan Monitoring in 2009 occurred for *Selenastrum capricornutum* toxicity and chlorpyrifos (Table VII-6). The upstream location, Highline Canal @ Lombardy Rd, allows for an analysis of the contribution of each portion of the watershed to loads measured at Highline Canal @ Hwy 99. Highline Canal @ Lombardy Rd is sampled as a separate monitoring location and is not considered an upstream MPM location to Highline Canal @ Hwy 99.

Highline Canal @ Hwy 99 is scheduled as a high priority site starting in 2010. Highline Canal is considered an impaired water body under the proposed 2008 Basin Plan and is listed on the 303d list for chlorpyrifos, simazine, unknown toxicity, and sediment toxicity. Agriculture is listed as the potential source for the chlorpyrifos, simazine and unknown toxicity impairments; sediment toxicity is listed with an unknown potential source.

Table VII-2. Highline Canal @ Hwy 99 sampling events per season and year.

	2004		2005		2006		2007		2008			2009		
	Irrigation	Storm	Irrigation	Storm	Irrigation	Storm	Irrigation	Storm	Irrigation	Fall	Winter	Storm	Irrigation	Fall
Events Sampled	NA	NA	5	2	5	1	6	2	6	0	0	1	6	2
Events Not Sampled	NA	NA	0	0	0	1	0	1	0	2	2	0	0	1
Total	NA	NA	5	2	5	2	6	3	6	2	2	1	6	3

NA indicates that this site was not sampled during this season/year.

Table VII-3. Number of analyses performed per analyte in each sampling season and year for the Highline Canal @ Hwy 99 sample site. Only environmental samples with a sample replicate and lab replicate number of one are shown.

Method	Analyte	2005	2006		2007		2008			2009			
		Irrigation	Storm	Irrigation	Storm	Irrigation	Storm	Irrigation	Fall	Winter	Storm	Irrigation	Fall
Field and Physical Parameters													
EPA 110.2	Color	5	2	5	1	6	2	6					
EPA 160.1	Dissolved Solids	5	2	5	1	6	2	6			2	6	1
EPA 160.2	Suspended Solids										2	6	1
EPA 180.1	Turbidity	5	2	5	1	6	2	6			2	6	1
EPA 405.1	BOD			1	1	2							
EPA 415.1	Total Organic Carbon	5	2	5	1	6	2	6			2	6	1
SM 9223	E. coli	5	2	5	1	6	2	6			2	6	1
NA	Dissolved Oxygen	6	5	5	1	8	5	15	2	2	2	6	2
NA	Specific Conductivity	6	5	5	1	8	5	15	2	2	2	6	2
NA	pH	6	5	5	1	8	5	15	2	2	2	6	2
Carbamates													
EPA 8321A	Aldicarb			5	1	6	2	6					
EPA 8321A	Carbaryl			5	1	6	2	6					
EPA 8321A	Carbofuran			5	1	6	2	6					
EPA 8321A	Diuron			5	1	6	2	6					
EPA 8321A	Linuron			5	1	6	2	6					
EPA 8321A	Methiocarb			5	1	6	2	6					
EPA 8321A	Methomyl			5	1	6	2	6					
EPA 8321A	Oxamyl			5	1	6	2	6					
Organochlorines													
EPA 8081A	DDD(p,p')			5	1	6	2	6					
EPA 8081A	DDE(p,p')			5	1	6	2	6					
EPA 8081A	DDT(p,p')			5	1	6	2	6					
EPA 8081A	Dicofol			5	1	6	2	6					
EPA 8081A	Dieldrin			5	1	6	2	6					

Method	Analyte	2005	2006		2007		2008			2009			
		Irrigation	Storm	Irrigation	Storm	Irrigation	Storm	Irrigation	Fall	Winter	Storm	Irrigation	Fall
EPA 8081A	Endrin			5	1	6	2	6					
EPA 8081A	Methoxychlor			5	1	6	2	6					
Organophosphates													
EPA 8141A	Azinphos methyl			5	1	6	2	6					
EPA 8141A	Chlorpyrifos	5	2	5	1	6	2	7				1	
EPA 8141A	Diazinon	5	2	5	1	6	2	6					
EPA 8141A	Dimethoate			5	1	6	2	6					
EPA 8141A	Disulfoton			5	1	6	2	6					
EPA 8141A	Malathion			5	1	6	2	6					
EPA 8141A	Methamidophos			5	1	6	2	6					
EPA 8141A	Methidathion			5	1	6	2	6					
EPA 8141A	Molinate			5	1	6	2	6					
EPA 8141A	Parathion, Methyl			5	1	6	2	6					
EPA 8141A	Phorate			5	1	6	2	6					
EPA 8141A	Phosmet			5	1	6	2	6					
EPA 8141A	Thiobencarb			5	1	6	2	6					
Pyrethroids													
EPA 8081A	Bifenthrin	1	2	5	1	6	2	6					
EPA 8081A	Cyfluthrin, total	1	2	5	1	6	2	6					
EPA 8081A	Cyhalothrin, lambda, total	5	2	5	1	6	2	6					
EPA 8081A	Cypermethrin, total	5	2	5	1	6	2	6					
EPA 8081A	Esfenvalerate/Fenvalerate, total	5	2	5	1	6	2	6					
EPA 8081A	Permethrin, total	5	2	5	1	6	2	6					
Triazines													
EPA 547M	Glyphosate			5	1	6	2	6					
EPA 549.2M	Paraquat dichloride			5	1	6	2	6					
EPA 619	Atrazine			5	1	6	2	6					
EPA 619	Cyanazine			5	1	6	2	6					

Method	Analyte	2005	2006		2007		2008			2009			
		Irrigation	Storm	Irrigation	Storm	Irrigation	Storm	Irrigation	Fall	Winter	Storm	Irrigation	Fall
EPA 619	Simazine			5	1	6	2	6					
Metals (Total)													
EPA 200.8	Arsenic			5	1	6	2	6					
EPA 200.8	Boron			5	1	6	2	6					
EPA 200.8	Cadmium			5	1	6	2	6					
EPA 200.8	Copper			5	1	6	2	10					
EPA 200.8	Lead			5	1	6	2	6					
EPA 200.8	Nickel			5	1	6	2	6					
EPA 200.8	Selenium			5	1	2	2	6					
EPA 200.8	Zinc			5	1	6	2	6					
Nutrients													
SM 2340 C	Hardness as CaCO3			5	1	6	2	10					
EPA 300.0	Nitrate as N			5	1	6	2	6					
EPA 350.2	Ammonia as N			5	1	6	2	6			2	6	1
EPA 351.3	Nitrogen, Total Kjeldahl			5	1	6	2	6			1	1	
EPA 353.2	Nitrate + Nitrite as N										2	6	1
EPA 354.1	Nitrite as N			5	1	6	2	6					
EPA 365.2	Orthophosphate as P			5	1	6	2	6			1	1	
EPA 365.2	Phosphate as P			5	1	6	2	6			2	6	1
Toxicity													
EPA 0821/R-02-012	Ceriodaphnia dubia	6	3	6	1	8	2	7					
EPA 0821/R-02-012	Pimephales promelas	5	2	5	1	7	2	7					
EPA 0821/R-02-013	Selenastrum capricornutum	5	3	5	1	7	2	7				1	
EPA 600/R-99-064	Hyalella azteca	5	1	2		1	1	2					

Table VII-4. Highline Canal @ Hwy 99. 2007 Management Plan additional (A) sampling schedule for *Ceriodaphnia*. "X" indicates the site, month, and analyte sampled.

Sample Site	Date	Type	<i>Ceriodaphnia dubia</i>
Highline Canal @ Hwy 99	25-Sep-07	A	X

Table VII-5. Highline Canal @ Hwy 99. 2008 Management Plan additional (A) sampling schedule for chlorpyrifos, copper, and *Ceriodaphnia*. "X" indicates the site, month, and analyte sampled.

Sample Site	Date	Type	Chlorpyrifos	Copper	<i>Ceriodaphnia dubia</i>
Highline Canal @ Hwy 99	29-Apr-09	A		X	
Highline Canal @ Hwy 99	07-May-09	A			X
Highline Canal @ Hwy 99	03-Jun-09	A		X	
Highline Canal @ Hwy 99	08-Jul-09	A	X	X	
Highline Canal @ Hwy 99	05-Aug-09	A		X	
Highline Canal @ Hwy 99	09-Sep-09	A			X

Table VII-6. Highline Canal @ Hwy 99. 2009 Management Plan sampling schedule for chlorpyrifos and *Selenastrum*. "X" indicates the site, month and analyte sampled.

Site Name	Date	Chlorpyrifos	<i>Selenastrum capricornutum</i>
Highline Canal @ Hwy 99	21-Apr-09		X
Highline Canal @ Hwy 99	19-May-09		X
Highline Canal @ Hwy 99	21-Jul-09	X	

Exceedance History

During Coalition monitoring, exceedances of WQTLs for field and physical parameters, *E. coli*, nutrients, metals, pesticides, and water column and sediment toxicity occurred within the Highline Canal site subwatershed (Table VII-7). Samples collected over the entire five years of monitoring at the normal monitoring and upstream MPM sites resulted in several exceedances, including DO (1), pH (14), SC (1), TDS (2), *E. coli* (8), ammonia (2), copper (7), lead (7), chlorpyrifos (5), and diuron (2). Water column toxicity has occurred eight times; four times each to *Ceriodaphnia dubia* and to *Selenastrum capricornutum*. Sediment toxicity to *Hyalella azteca* has occurred six times.

During 2009, the only WQTL exceedances that occurred were for pH (February, June, August, September, and December), *E. coli* (May) and chlorpyrifos (July, Table VII-7).

All exceedances are listed in Table VII-7 by season and date and are based on WQTLs listed in the introduction to the ESJWQC Management Plan. The priority level (A-E) assigned to each constituent is listed in the bottom row of Table VII-7 and is determined using the ESJWQC Management Plan prioritization process flow chart (Figure 4). Constituents that are the highest priority in the Highline Canal @ Highway 99 subwatershed are chlorpyrifos (A/B), copper (C), diuron (C), and toxicity to *Selenastrum capricornutum*, *Ceriodaphnia dubia*, and *Hyalella azteca* (D).

Table VII-7. All exceedances experienced in samples collected within the Highline Canal @ Hwy 99 site subwatershed between March 2005 and December 2009 (sorted by season and date). If the water quality trigger limit is based on hardness then the hardness value is shown in parenthesis; otherwise the WQTL used to evaluate the data is listed in the header after the analyte.

Station Name	Season	Sample Date	Oxygen, Dissolved, 7 mg/L	pH, 6.5-8.5 units	Specific Conductivity, 700 μ S/cm	<i>E. coli</i> , 235 MPN/100 mL	Dissolved Solids, 450 mg/L	Ammonia, 1.5 mg/L	Copper, Total, 1,300 μ g/L	Lead, Total, 15 μ g/L	Chlorpyrifos, 0.015 μ g/L	Diuron, 2 μ g/L	<i>C. dubia</i> , Survival (%)	<i>H. azteca</i> , Survival (%)	<i>S. capricornutum</i> , Total Cell Count
Highline Canal @ Hwy 99	Irrigation	5/10/2005											45		
Highline Canal @ Hwy 99	Irrigation	5/19/2005											0		
Highline Canal @ Hwy 99	Irrigation	9/20/2005												87.5	
Highline Canal @ Hwy 99	Storm	3/1/2006	5.6								0.021				12750
Highline Canal @ Hwy 99	Storm	3/16/2006				300							0		
Highline Canal @ Hwy 99	Storm	5/2/2006		8.73											
Highline Canal @ Hwy 99	Irrigation	5/17/2006								0.42 (0.36)					
Highline Canal @ Hwy 99	Irrigation	8/9/2006								0.39 (0.31)				88.75	
Highline Canal @ Hwy 99	Irrigation	9/5/2006												80	
Highline Canal @ Hwy 99	Irrigation	9/13/2006											60		
Highline Canal @ Hwy 99	Storm	2/11/2007							3 (2.2)	0.52 (0.36)		25			
Highline Canal @ Hwy 99	Irrigation	4/17/2007							11 (10.1)	5.1 (3.59)					
Highline Canal @ Hwy 99	Irrigation	5/15/2007		8.56		250									
Highline Canal @ Hwy 99	Irrigation	6/19/2007				320			2.4 (1.9)	0.5 (0.31)					
Highline Canal @ Hwy 99	Irrigation	7/17/2007				440			3.2 (2.2)	1 (0.36)	0.017				
Highline Canal @ Hwy 99	Irrigation	8/14/2007		8.62					1.9 (1.7)	0.44 (0.26)					
Highline Canal @ Hwy 99	Irrigation	9/25/2007		8.73											
Highline Canal @ Hwy 99	Storm	1/24/2008				>2400	500	3.3	37 (14.7)		0.019	3.2			
Highline Canal @ Hwy 99	Storm	2/26/2008			747	>2400	520	8.3	81 (81)						817147

Station Name	Season	Sample Date	Oxygen, Dissolved, 7 mg/L	pH, 6.5-8.5 units	Specific Conductivity, 700 µS/cm	E. coli, 235 MPN/100 mL	Dissolved Solids, 450 mg/L	Ammonia, 1.5 mg/L	Copper, Total, 1,300 µg/L	Lead, Total, 15µg/L	Chlorpyrifos, 0.015 µg/L	Diuron, 2 µg/L	C. dubia, Survival (%)	H. azteca, Survival (%)	S. capricornutum, Total Cell Count
Highline Canal @ Hwy 99	Sediment	3/4/2008												90	
Highline Canal @ Hwy 99	Storm	3/4/2008		9.32											
Highline Canal @ Hwy 99	Irrigation	4/22/2008													791095
Highline Canal @ Hwy 99	Irrigation	5/7/2008		8.69											
Highline Canal @ Hwy 99	Irrigation	5/20/2008				240									1022306
Highline Canal @ Hwy 99	Irrigation	6/3/2008		8.61											
Highline Canal @ Hwy 99	Irrigation	7/22/2008									0.021				
Highline Canal @ Hwy 99	Irrigation	8/19/2008		9.24											
Highline Canal @ Hwy 99	Sediment	8/28/2008												91	
Highline Canal @ Hwy 99	Irrigation	9/9/2008		8.54											
Highline Canal @ Hwy 99	Sediment	10/2/2008												89	
Highline Canal @ Hwy 99	Storm	2/7/2009		8.86											
Highline Canal @ Hwy 99	Irrigation	5/19/2009				340									
Highline Canal @ Hwy 99	Irrigation	6/16/2009		8.95											
Highline Canal @ Hwy 99	Irrigation	7/21/2009									0.093				
Highline Canal @ Hwy 99	Irrigation	8/18/2009		9.03											
Highline Canal @ Hwy 99	Irrigation	9/22/2009		8.61											
Highline Canal @ Hwy 99	Storm	12/15/2009		8.61											
Constituent Priority			NP	E	E	E	E	E	C	E	A/B	C	D	D	D

NP- Not prioritized. Fewer than two exceedances for this constituent at this site within three years and currently no TMDL for constituent.

2007 - 2009 Management Plan Monitoring Results

Management Plan Monitoring results are included in Table VII-8 for *Ceriodaphnia dubia* (2007), for chlorpyrifos, copper and *Ceriodaphnia dubia* (2008), for chlorpyrifos and *Selenastrum capricornutum* (2009).

2007

Highline Canal @ Hwy 99 was sampled monthly for toxicity to *Ceriodaphnia* and twice in September as a part of additional 2007 MPM (Table VII-8). No toxicity occurred in any samples collected (Table VII-8). Chlorpyrifos exceeded the WQTL for a second time at this site in 2007 and copper exceeded the hardness based WQTL five times during 2007 normal monitoring.

2008

In 2008, normal monitoring occurred for all constituents monthly from April through September. Management Plan Monitoring occurred at the Highline Canal @ Hwy 99 monitoring site for copper, chlorpyrifos and *Ceriodaphnia dubia* toxicity (Table VII-8). Copper was detected in every sample collected during the irrigation season of 2008 without exceeding the WQTL. Chlorpyrifos exceeded the WQTL once in 2008 during July normal monitoring (0.021µg/L, Table VII-8). Toxicity to *Ceriodaphnia* was not experienced in any samples collected in 2008. Toxicity to *Selenastrum capricornutum* occurred in February, April, and May of 2008 and are all associated with copper detections.

2009

Highline Canal @ Hwy 99 was a Core site during 2009. Management Plan Monitoring schedule for chlorpyrifos in July and toxicity to *Selenastrum* in April and May revealed no detections of chlorpyrifos and no toxicity to *Selenastrum*.

Management Plan Monitoring for copper, chlorpyrifos, *Selenastrum* toxicity, and *Hyaella* toxicity are planned for the 2010 irrigation season. Management Plan Monitoring during January, February, and March 2011 will also occur for copper, chlorpyrifos, diuron and toxicity to *Hyaella*.

Table VII-8. Highline Canal @ Hwy 99. Normal monitoring (NM) and Management Plan Monitoring (MPM) results for *Ceriodaphnia* from the 2007 irrigation season. Normal monitoring (NM) and Management Plan Monitoring (MPM) results for copper, chlorpyrifos and *Ceriodaphnia* for the 2008 irrigation season and Management Plan Monitoring (MPM) results for chlorpyrifos and *Selenastrum* for 2009. Exceedance values are in bold.

		Month:	April	May	June	July	August	September
2007 NM (@ Hwy 99)	Date	4/17/07	5/15/07	6/19/07	7/17/07	8/14/07	9/11/07	
	<i>C. dubia</i> toxicity (% Control)	90	100	100	95	100	100	
2007 MPM (@ Hwy 99)	Date	NA	NA	NA	NA	NA	9/25/07	
	<i>C. dubia</i> toxicity (% Control)	NA	NA	NA	NA	NA	100	
2008 NM (@ Hwy 99)	Date	4/22/08	5/20/08	6/17/08	7/22/08	8/19/08	9/23/08	
	Copper (µg/L)	1.8	1.6	1.2	1.2	1.0	1.1	
	Chlorpyrifos (µg/L)	<0.003	<0.003	<0.003	0.021	<0.003	<0.003	
	<i>C. dubia</i> toxicity (% Control)	100	100	100	100	100	100	
2008 MPM (@ Hwy 99)	Date	4/29/08	5/7/08	6/3/08	7/8/08	8/5/08	9/9/08	
	Copper (µg/L)	1.4	NA	1.5	1.7	1.6	NA	
	Chlorpyrifos (µg/L)	NA	NA	NA	<0.003	NA	NA	
	<i>C. dubia</i> toxicity (% Control)	NA	100	NA	NA	NA	100	
2009 MPM (@ Hwy 99)	Date	4/21/09	5/19/09	NA	7/21/09	NA	NA	
	Chlorpyrifos (µg/L)	NA	NA	NA	<0.003	NA	NA	
	<i>S. capricornutum</i> toxicity (% Control)	428	222	NA	NA	NA	NA	

NA - Not applicable. This site was not sampled during this month.

Load Calculations

Loads have been calculated for all chlorpyrifos detections in the site subwatershed (Table VII-9) based on the following formula:

$$\text{Load} = \text{Discharge (cfs)} \times 28.317\text{L/ft}^3 \times \text{Concentration (milligram/L} \times 1000 \text{ or } \mu\text{g/L)}.$$

The load values for constituents in this report represent instantaneous loads only. These values should not be used to extrapolate loading over any period of time (e.g. weekly, monthly, seasonal or annual). The primary purpose for reporting instantaneous loads is to provide a normalization of the concentrations by flow for various constituents at the time the samples were collected.

Table VII-9. Highline Canal @ Hwy 99. Instantaneous load calculations for chlorpyrifos, copper and diuron where discharge was measured (sorted by analyte and date).

Station Name	Analyte Name	Sample Date	Discharge cfs	Concentration µg/L	Loading Rate µg/sec
Highline Canal @ Hwy 99	Chlorpyrifos	1-Mar-06	0	0.021	0
Highline Canal @ Hwy 99	Chlorpyrifos	12-Jul-06	111.81	0.012	37.99
Highline Canal @ Hwy 99*	Chlorpyrifos	17-Jul-07	56.81	0.015	24.13
Highline Canal @ Hwy 99	Chlorpyrifos	17-Jul-07	56.81	0.017	27.35
Highline Canal @ Hwy 99	Chlorpyrifos	24-Jan-08	55.88	0.019	30.06
Highline Canal @ Hwy 99	Chlorpyrifos	26-Feb-08	6.91	0.0076	1.49
Highline Canal @ Hwy 99	Chlorpyrifos	21-Jul-09	118.93	0.093	313.20

Station Name	Analyte Name	Sample Date	Discharge cfs	Concentration µg/L	Loading Rate µg/sec
Highline Canal @ Hwy 99	Copper	17-May-06	82.45	1.8	4202.53
Highline Canal @ Hwy 99	Copper	14-Jun-06	90.82	1.7	4371.97
Highline Canal @ Hwy 99	Copper	12-Jul-06	111.81	1.6	5065.80
Highline Canal @ Hwy 99	Copper	9-Aug-06	162.04	1.3	5965.03
Highline Canal @ Hwy 99	Copper	13-Sep-06	72.64	1.3	2674.03
Highline Canal @ Hwy 99	Copper	11-Feb-07	55.43	3	4708.83
Highline Canal @ Hwy 99	Copper	17-Apr-07	22.54	11	7020.92
Highline Canal @ Hwy 99	Copper	15-May-07	60.6	1.4	2402.41
Highline Canal @ Hwy 99	Copper	19-Jun-07	84.28	2.4	5727.74
Highline Canal @ Hwy 99*	Copper	17-Jul-07	56.81	3.1	4986.94
Highline Canal @ Hwy 99	Copper	17-Jul-07	56.81	3.2	5147.80
Highline Canal @ Hwy 99	Copper	14-Aug-07	81.98	1.9	4410.71
Highline Canal @ Hwy 99	Copper	11-Sep-07	33.38	1.5	1417.83
Highline Canal @ Hwy 99	Copper	24-Jan-08	55.88	37	58547.10
Highline Canal @ Hwy 99	Copper	26-Feb-08	6.91	81	15849.31
Highline Canal @ Hwy 99	Copper	29-Apr-08	34.66	1.4	1374.05
Highline Canal @ Hwy 99	Copper	20-May-08	81.42	1.6	3688.91
Highline Canal @ Hwy 99	Copper	3-Jun-08	57.91	1.5	2459.76
Highline Canal @ Hwy 99	Copper	19-Aug-08	37.45	1	1060.47
Highline Canal @ Hwy 99	Diuron	11-Feb-07	55.43	25	39240.28
Highline Canal @ Hwy 99	Diuron	24-Jan-08	55.88	3.2	5063.53
Highline Canal @ Hwy 99	Diuron	26-Feb-08	6.91	0.43	84.14

*Field Duplicate

Source Identification and Outreach

Priority A/B Constituents

Chlorpyrifos is the only priority A/B constituent listed under the Highline Canal @ Hwy 99 Management Plan.

Chlorpyrifos

Chlorpyrifos has exceeded the WQTL of 0.015 µg/L five times in this subwatershed from 2005 through 2009 (March 2006, July 2007-2009 and January 2008). The Coalition uses a combination of monitoring data and evaluation of Pesticide Use Report (PUR) data to identify possible sources.

PUR data are reviewed for the number of monthly chlorpyrifos applications, pounds active ingredient (AI) applied, and acres treated (Table VII-10, Figure VII-2). The amount of chlorpyrifos applied within the subwatershed in 2009 has decreased significantly from the 2007

level. The amount of chlorpyrifos used in 2007 was more than 20 times the amount of chlorpyrifos used in 2009 (20,905 lbs in 2007 compared to 1010 lbs in 2009, Table VII-10, Figure VII-2). The greatest amount of chlorpyrifos applied in 2005 and 2009 was during May, while July received the most chlorpyrifos applied in 2006 and 2007, and in 2008 the month of August received the most pounds applied (Table VII-10, Figure VII-2). Exceedances did not always occur during months of the greatest amount of use (Figure VII-2).

The Coalition also used PUR data to assess which crops receive the most applications of chlorpyrifos. The highest application rates are associated with almonds, corn, and walnuts (Table VII-11). The largest amount of chlorpyrifos from 2005 through 2009 was to almonds and walnuts, followed by grain/sorghum/milo and corn (Table VII-11). The most common product containing chlorpyrifos applied within the subwatershed was Lorsban (Table VII-11).

Table VII-10. Number of chlorpyrifos applications, total pounds applied and total acres treated by month for January 2005 through December 2009 in the Highline Canal @ Hwy 99 site subwatershed. If a month is not included in the table, no applications were made.

Month/Year	Number of Chlorpyrifos Applications	Pounds Applied	Acres Treated
January, 2005	14	768.3	549.2
March, 2005	5	231.9	492.2
April, 2005	2	360.2	360
May, 2005	17	6208.7	1686
June, 2005	19	1313.8	778
July, 2005	42	3632.3	2094
August, 2005	20	2815.7	2271
September, 2005	2	33.9	17
November, 2005	4	1444.7	724.6
January, 2006	7	2041.1	1035
May, 2006	13	2535.4	2007
June, 2006	11	1753.6	1538.9
July, 2006	28	6231.2	3615.2
August, 2006	30	2606.1	2461.5
September, 2006	2	85.1	42
March, 2007	1	110.6	217.4
April, 2007	1	6.2	20
May, 2007	25	3250.0	2869
June, 2007	8	3971.9	1435
July, 2007	37	9124.0	5450.3
August, 2007	3	4442.6	386.4
May, 2008	5	467.1	348
June, 2008	2	121.8	75
July, 2008	16	684.7	503
August, 2008	19	1035.1	654.5

Month/Year	Number of Chlorpyrifos Applications	Pounds Applied	Acres Treated
September, 2008	2	31.9	20
May, 2009	4	433.5	376
June, 2009	2	40.2	35
July, 2009	12	361.2	209
August, 2009	10	175.6	146
Summaries by Year			
2005 Total	125	16809.6	8972
2006 Total	91	15252.5	10699.6
2007 Total	75	20905.3	10378.1
2008 Total	44	2340.7	1600.5
2009 Total	28	1010.5	766
Total	363	56,318.6	32,416.2

Figure VII-2. Pounds of chlorpyrifos applied within the Highline Canal @ Hwy 99 site subwatershed by month for 2005-2009. Asterisk (*) denotes months with exceedances.




Table VII-11. Total pounds AI for chlorpyrifos based on PUR data from 2005-2009 within the Highline Canal @ Hwy 99 subwatershed.

Chemical Name	Commodity	Product Name	Lbs AI Applied
CHLORPYRIFOS	ALFALFA	LOCK-ON INSECTICIDE	2.504
		NUFOS 4E	110.624
	ALFALFA (FORAGE - FODDER) (ALFALFA HAY)	LOCK-ON INSECTICIDE	0.501
		LORSBAN 4E-HF	9.969
		NUFOS 4E	221.940
	ALMOND	CHLORPYRIFOS 4E AG	871.181
		DURSBAN 4E INSECTICIDE	20.178
		GOVERN 4E INSECTICIDE	4098.714
		LORSBAN 4E INSECTICIDE	478.782
		LORSBAN 4E-HF	5189.897
		LORSBAN 50W WETTABLE POWDER INSECTICIDE	30.000
		LORSBAN-4E	29930.115
	CORN (FORAGE - FODDER)	NUFOS 4E	1053.369
		CHLORPYRIFOS 4E AG	147.292
		LORSBAN 15G GRANULAR INSECTICIDE	81.675
		LORSBAN 4E INSECTICIDE	120.017
		LORSBAN 4E-HF	199.844
		LORSBAN-4E	639.321
	CORN FOR/FOD	NUFOS 4E	490.793
		CHLORPYRIFOS 4E AG	52.987
		LORSBAN 4E-HF	594.048
		LORSBAN-4E	318.158
		NUFOS 4E	665.900
	GRAPES, WINE	WARHAWK	326.241
		LORSBAN 4E-HF	1444.745
	PEACH	CHLORPYRIFOS 4E AG	118.666
		LORSBAN 4E-HF	41.746
		LORSBAN-4E	36.018
	SORGHUM MILO	NUFOS 4E	4148.796
	SWEET POTATO	LORSBAN 15G GRANULAR INSECTICIDE	93.150
	WALNUT	CHLORPYRIFOS 4E AG	270.192
		GOVERN 4E INSECTICIDE	30.301
LORSBAN 4E-HF		279.139	
LORSBAN-4E		757.074	
NUFOS 4E		480.930	
WALNUT (ENGLISH WALNUT, PERSIAN WALNUT)	CHLORPYRIFOS 4E AG	135.096	
	GOVERN 4E INSECTICIDE	190.896	
	LORSBAN 4E-HF	484.456	
	LORSBAN-4E	1957.075	
	NUFOS 4E	196.236	
ALFALFA – Total Pounds Chlorpyrifos Applied			345.538

Chemical Name	Commodity	Product Name	Lbs AI Applied
		ALMOND – Total Pounds Chlorpyrifos Applied	41672.236
		CORN – Total Pounds Chlorpyrifos Applied	3636.275
		GRAPES – Total Pounds Chlorpyrifos Applied	1444.745
		PEACH – Total Pounds Chlorpyrifos Applied	196.430
		SORGHUM MILO – Total Pounds Chlorpyrifos Applied	4148.796
		SWEET POTATO – Total Pounds Chlorpyrifos Applied	93.150
		WALNUT – Total Pounds Chlorpyrifos Applied	4781.395
Total pounds chlorpyrifos applied (2005 - 2009)			56,318.565

Based on results from general management practice surveys, approximately 41% of the member acreage indicates no runoff occurs and 18% of the member acreage uses recirculation and/or tailwater return systems. Chlorpyrifos exceedances often occur in July (2007-2009), but have also occurred during storm monitoring (January 2008 and March 2006). The Coalition targets growers applying chlorpyrifos and also those with the potential to drain to the canal. Growers should be encouraged to review their operation to determine if irrigation return flows are managed properly. Owners operating orchards should evaluate their aerial applications to minimize spray drift.

Coalition outreach since 2007 has included grower meetings and the mailing/distribution of information. A complete list of Coalition Outreach during 2009 is provided in the Summary of Coalition Outreach Activities section of the Management Plan Update Report. The Coalition has begun the process of contacting growers, conducting individual meetings, and compiling individual surveys for Highline Canal @ Hwy 99. The Coalition focused outreach to growers owning parcels with the potential to drain to the creek and those applying chlorpyrifos. Using these criteria, a list of 10 targeted growers was created. Individual meetings are currently ongoing and will be completed by the end of May 2010. Individual contacts will focus mainly on chlorpyrifos exceedances however all water quality results will be reviewed and discussed including copper, diuron, ammonia, pH, SC, TDS, *E. coli*, and lead WQTLs exceedances as well as toxicity to *Ceriodaphnia dubia*, *Selenastrum capricornutum* and *Hyalella azteca*.

The Coalition will compile and analyze the surveys by the end of August 2010. Based on the exact timing of each individual meeting and/or a grower's resources, some owners will not be able to implement recommended management practices until the 2011 irrigation season. In addition, long term structural BMPs (e.g. sediment ponds) will most likely take longer than two years to implement and will require additional tracking to document their implementation. The Coalition will follow up with growers between February and April of 2011 to determine what practices were implemented during the first irrigation year. Growers who have not already been contacted or have indicated that they have not yet implemented practices during the first year will be contacted from February to April of 2012 to determine what practices were implemented during the dormant season and/or the second irrigation year. To evaluate the management practice process, the Coalition will monitor for high priority constituents during months of past exceedances in 2010 and 2011. Depending on when additional management practices are implemented, the Coalition may monitor through 2012 to evaluate improvements in water quality. The Coalition anticipates that changes in management practices by members

that have direct drainage and/or are have the potential for spray drift will affect downstream water quality by the end of the second year as a high priority site subwatershed.

The effectiveness of the management practices will be determined through the monitoring of water quality in the years following implementation. However, due to the presence of dairies and nonmembers in the subwatersheds, implementation of management practices by only coalition members may not result in improved water quality. If water quality fails to improve, the Coalition will identify parcels from dairies and nonmembers that could contribute to the exceedances and provide that information to the Regional Board.

Priority C Constituents

Highline Canal @ Hwy 99 is listed for copper and diuron as priority C constituents.

Copper

Copper has exceeded the hardness based WQTL seven times in this subwatershed from 2005 through 2009. Copper exceedances occurred five times during 2007 (February, April, and June-August), and twice in 2008 (January and February).

PUR data are reviewed for the number of monthly copper applications, pounds active ingredient (AI) applied, and acres treated (Table VII-12, Figure VII-3). The amount of copper applied within the subwatershed decreased annually since 2006 (Table VII-12, Figure VII-3). The month of January consistently receives the greatest amount of copper applied (Table VII-12, Figure VII-3). Not all exceedances occurred during months of the greatest amount of use (Figure VII-3).

The Coalition also used PUR data to assess which crops receive the most applications of copper. The largest amount of copper applied from 2005 through 2009 was associated with almond, peach and walnut orchards (Table VII-13). The most common products containing copper applied withing the subwatershed were Dupont Kocide and Kocide (Table VII-13).

Table VII-12. Number of copper applications, total pounds applied and total acres treated by month for January 2005 through December 2009 in the Highline Canal @ Hwy 99 site subwatershed. If a month is not included in the table, no applications were made.

Month/Year	Number of Copper Applications	Pounds Applied	Acres Treated
January, 2005	63	21059.0	5037.3
February, 2005	101	5123.9	4325.19
March, 2005	35	3319.9	1511.23
April, 2005	41	5387.7	2321.7
May, 2005	18	2139.3	678
June, 2005	2	100.1	45
August, 2005	2	293.8	63
September, 2005	1	8.6	4

Month/Year	Number of Copper Applications	Pounds Applied	Acres Treated
October, 2005	2	9.1	8
November, 2005	20	15983.3	3993.08
December, 2005	59	14774.6	1746.1
January, 2006	79	90658.2	13310
February, 2006	96	10879.8	5316.13
March, 2006	21	1152.6	1374.1
April, 2006	25	2544.1	665
May, 2006	22	2701.2	1545.4
August, 2006	2	318.6	63
November, 2006	12	12148.1	3039.59
December, 2006	17	2809.6	267.09
January, 2007	73	28634.4	5479.75
February, 2007	50	4993.0	2280.93
March, 2007	55	3895.0	2407.83
April, 2007	49	5367.4	1172
May, 2007	17	2220.3	511.1
January, 2008	31	8127.0	865.4
February, 2008	49	3733.2	1229.56
March, 2008	42	2405.4	1163.15
April, 2008	4	139.9	93
May, 2008	1	9.2	5
July, 2008	2	135.0	63
December, 2008	25	3298.2	406.94
January, 2009	24	4650.5	576.94
February, 2009	52	3077.1	963.71
March, 2009	17	550.4	396
April, 2009	14	847.4	219
May, 2009	5	401.6	63
June, 2009	2	191.2	64
July, 2009	2	153.7	63
December, 2009	3	268.1	47
Summaries by Year			
2005 Total	344	68199.1	19732.6
2006 Total	274	123212.1	25580.31
2007 Total	244	45110.2	11851.61
2008 Total	154	17847.9	3826.05
2009 Total	119	10140.0	2392.65
Total	1,135	264,509.3	63,383.22

Figure VII-3. Pounds of copper applied within the Highline Canal @ Hwy 99 site subwatershed by month for 2005-2009. Asterisk (*) denotes months with exceedances.




Table VII-13. Total pounds AI for copper based on PUR data from 2005-2009 within the Highline Canal @ Hwy 99 subwatershed.

Chemical Name	Commodity	Product Name	Lbs AI Applied
COPPER	ALMOND	BASIC COPPER 53	74465.006
		BASIC Copper53	9472.484
		BASICOP	1229.050
		CHAMP FORMULA 2 FLOWABLE	9525.065
		CHAMPION WETTABLE POWDER	6007.540
		CUPROFIX DISPERSS	22.140
		CUPROFIX ULTRA 40 DISPERSS	489.879
		DUPONT KOCIDE 2000 FUNGICIDE/BACTERICIDE	5241.062
		HYDROX	4350.500
		KOCIDE 101	12470.965
		KOCIDE 2000	4366.704
		KOCIDE DF	8957.943
		NORDOX	56.400
		NORDOX 75 WG	19651.184
		NU-COP 50 WP	6190.800
		NU-COP 50DF	29077.302
		ALMOND, ORGANIC	CHAMPION WETTABLE POWDER
	APRICOT	KOCIDE DF	310.070
	CHERRY	BASIC COPPER 53	9.800
		CHAMPION WETTABLE POWDER	23.100
	GRAPES	BASIC Copper53	230.300
		CHAMP FORMULA 2 FLOWABLE	183.036
		CHAMPION WETTABLE POWDER	10.780
	GRAPES, WINE	CHAMP FORMULA 2 FLOWABLE	1859.947723
		CLEAN CROP C-O-C-S 15 SULFUR 25 DUST	747.328475
		IAP COPPER SULFUR 15-25 DUST	344.97
		KOCIDE 4.5 LF	318.4275093
		KOCIDE 606 FLOWABLE AGRICULTURAL FUNGICIDE	713.781873
	N-OUTDR CONTAINER/FLD GRWN PLANTS	CHAMPION WETTABLE POWDER	1135.303
		KOCIDE 2000	242.638
	N-OUTDR PLANTS	NU-COP 50 WP	71.610
		NU-COP 50DF	73.920
PEACH	BASIC COPPER 53	15082.886	
	BASIC Copper53	14828.331	
	CHAMPION WETTABLE POWDER	641.441	
	DUPONT KOCIDE 2000	535.041	

Chemical Name	Commodity	Product Name	Lbs AI Applied
		FUNGICIDE/BACTERICIDE	
		DUPONT KOCIDE 3000 FUNGICIDE/BACTERICIDE	39.254
		KOCIDE 101	1121.120
		KOCIDE 2000	104.264
		KOCIDE DF	1880.068
		NORDOX 75 WG	6706.295
		NU-COP 50DF	1398.320
	PEACH PROCESSNG	KOCIDE DF	98.240
		NU-COP 50 WP	40.040
		NU-COP 50DF	616.000
	PISTACHIO	CUPROFIX ULTRA 40 DISPERSS	81.054
		BASIC COPPER 53	911.400
		CHAMP FORMULA 2 FLOWABLE	426.002
		CHAMPION WETTABLE POWDER	49.280
		CUPROFIX ULTRA 40 DISPERSS	298.620
		DUPONT GX-569 FUNGICIDE/BACTERICIDE	9.220
		DUPONT KOCIDE 101 FUNGICIDE/BACTERICIDE	78.540
		DUPONT KOCIDE 2000 FUNGICIDE/BACTERICIDE	684.336
		KENTAN DF	279.528
		KOCIDE 101	2585.660
		KOCIDE 2000	1251.388
		KOCIDE DF	39.296
		NORDOX 75 WG	3580.852
		NU-COP 50DF	1068.760
		CHAMP FORMULA 2 FLOWABLE	744.871
		CHAMPION WETTABLE POWDER	337.260
		DUPONT KOCIDE 2000 FUNGICIDE/BACTERICIDE	645.600
		KOCIDE 101	3321.924
		KOCIDE 2000	4496.604
		KOCIDE DF	1882.524
		NORDOX 75 WG	656.098
		NU-COP 50DF	86.240
	WALNUT		
	WALNUT (ENGLISH WALNUT, PERSIAN WALNUT)		
		ALMOND – Total Pounds Copper Applied	191627.924
		APRICOT – Total Pounds Copper Applied	310.070
		CHERRY – Total Pounds Copper Applied	32.900
		GRAPES – Total Pounds Copper Applied	4408.571

Chemical Name	Commodity	Product Name	Lbs AI Applied
		N-OUTDOOR PLANTS – Total Pounds Copper Applied	1523.471
		PEACH – Total Pounds Copper Applied	43091.300
		PISTACHIO – Total Pounds Copper Applied	81.054
		WALNUT – Total Pounds Copper Applied	23434.004
Total pounds copper applied (2005 - 2009)			264,509.294

The Coalition will continue with its management plan strategy outlined above under the chlorpyrifos outreach section when conducting individual contacts. Orchard and vineyard operators will be advised to consider irrigation water retention and management to prevent copper from entering the waterway with runoff. Storm runoff management relevant to copper applications will also be discussed to prevent the winter exceedances. Individual contacts occurring within this subwatershed are described under the chlorpyrifos outreach section above and will include discussions of copper exceedances and the above management practices.

Diuron

Diuron is a soluble herbicide applied throughout the year in the Highline Canal watershed, primarily in the months of October – June. There was an exceedance of diuron in February 2007 and one in January 2008. The months with the largest amount of pounds applied are usually January (2006 and 2007) and December (2005), however April received the most pounds applied in 2009 (Table VII-14, Figure VII-4). Pounds applied and acres treated have decreased annually since 2005. The lowest amount applied between 2005 and 2009 was in April 2009; it was the only month that received applications that year (Table VII-14, Figure VII-4). Diuron was only applied to alfalfa, walnuts, and corn in this subwatershed, and alfalfa and walnuts received the most pounds applied (Table VII-15).

Table VII-14. Number of diuron applications, total pounds applied and total acres treated by month for January 2005 through December 2009 in the Highline Canal @ Hwy 99 site subwatershed. If a month is not included in the table, no applications were made.

Month/Year*	Number of Diuron Applications	Pounds Applied	Acres Treated
January, 2005	4	106.0	151
October, 2005	2	59.1	40
November, 2005	2	44.5	73
December, 2005	4	341.1	242.4
January, 2006	2	67.2	46
October, 2006	1	30.0	20
November, 2006	2	27.1	65
December, 2006	1	318.8	217.4
January, 2007	3	122.6	104.35
February, 2007	2	66.3	46

Month/Year*	Number of Diuron Applications	Pounds Applied	Acres Treated
March, 2007	1	17.8	40
April, 2007	2	33.4	75
June, 2007	2	1.1	28
April, 2009	2	15.3	104.9
Summaries by Year			
2005 Total	12	550.7	506.4
2006 Total	6	443.1	348.4
2007 Total	10	241.0	293.35
2009 Total	2	15.3	104.9
Total	30	1,250.1	1,253.05

*No applications of diuron reported for 2008.

Figure VII-4. Pounds of diuron applied within the Highline Canal @ Hwy 99 site subwatershed by month for 2005-2009. Asterisk (*) denotes months with exceedances. No use of diuron was reported for 2008.




Table VII-15. Total pounds AI for diuron based on PUR data from 2005-2009 within the Highline Canal @ Hwy 99 subwatershed.

Chemical Name	Commodity	Product Name	Lbs AI Applied	
DIURON	ALFALFA	DIREX 4L	107.207	
	ALFALFA (FORAGE - FODDER) (ALFALFA HAY)	DIREX 4L	649.476	
	CORN FOR/FOD	DU PONT KARMEX DF HERBICIDE	1.050	
	WALNUT		DIREX 4L	66.273
			DREXEL DIURON 4L HERBICIDE	66.488
			DUPONT DIREX 4L HERBICIDE	15.255
	WALNUT (ENGLISH WALNUT, PERSIAN WALNUT)		DIREX 4L	274.353
			DREXEL DIURON 4L HERBICIDE	69.972
ALFALFA – Total Pounds Diuron Applied			756.683	
CORN – Total Pounds Diuron Applied			1.050	
WALNUT – Total Pounds Diuron Applied			492.342	
Total pounds diuron applied (2005 - 2009)			1,250.074	

*No applications of diuron reported for 2008.

Management of diuron in this watershed should focus on providing information to the small number of growers whose applications could be associated with the exceedances. The Coalition’s management plan strategy for addressing diuron exceedances is to include diuron in discussions with individual growers relevant to preventing storm and irrigation discharge to waterways. Hopefully improved and additional water management practices will reduce or eliminate these exceedances.

Priority D Constituents

Highline Canal is listed for water column toxicity to *Ceriodaphnia dubia*, *Selenastrum capricornutum* and *Hyalella azteca*. *Selenastrum* toxicity was monitored during 2009 MPM and no water column toxicity experienced.

Ceriodaphnia toxicity

Toxicity to *Ceriodaphnia* occurred in May 2005 and in March and September 2006. The 2005 toxicity was persistent in the resample a week later; however, this was not the case in any of the 2006 toxicities. Both 2006 toxicities were associated with exceedances of the chlorpyrifos WQTL. Toxicity to *Ceriodaphnia* did not occur during 2008.

The Coalition's strategy for eliminating *Ceriodaphnia* toxicity will involve focusing on chlorpyrifos. If chlorpyrifos can be prevented from entering waterways throughout the irrigation season, the Coalition believes that *Ceriodaphnia* toxicity can also be reduced or eliminated. If chlorpyrifos is no longer detected in the slough but *Ceriodaphnia* toxicity persists, the Coalition will consider other pesticides to eliminate toxicity.

***Selenastrum* toxicity**

Water column toxicity to *Selenastrum capricornutum* occurred March 2006 and in February, April, and May 2008. The 2008 February storm toxicity coincided with a copper exceedance on the same day.

The Coalition's strategy for eliminating *Selenastrum* toxicity will involve focusing on copper. If copper can be prevented from entering storm water during the winter rainy season and early irrigation season, the Coalition believes that *Selenastrum* toxicity can also be reduced or eliminated. There were no exceedances of the copper WQTL and also no water column toxicity associated with *Selenastrum* during 2009 monitoring.

***Hyalella* toxicity**

Sediment toxicity to *Hyalella* occurred in the late irrigation months in of 2005 and 2006, and during both the storm and late irrigation seasons of 2008. Both the August 2006 and 2008 toxicities were persistent in the resample. Sediment toxicity was not monitored during 2009.

The Coalition has begun to conduct management plan sampling for sediment toxicities. As of 2009, the Coalition now tests for total organic carbon and grain size in all sediments and pyrethroids and chlorpyrifos in sediment that exhibits toxicity. Management Plan Monitoring is scheduled for this site in 2010.

Priority E Constituents

The following priority E constituents are listed under the Highline Canal @ Highway 99 Management Plan: ammonia, pH, SC, dissolved solids, *E. coli*, and lead. Exceedances of pH and *E. coli* continued in 2009. These constituents will remain low priority but will be discussed during individual contacts and annual grower meetings.

Evaluation

Highline Canal @ Hwy 99 is one of the second four priority site subwatersheds within the ESJWQC and is in its first year of focused outreach (2010-2012). The Coalition's strategy for the Highline Canal subwatershed has been to target growers along or adjacent to the waterway that have the potential to discharge. Focus will be on water retention management practice implementation and minimizing spray drift. Outreach

includes grower notification, management practice outreach and education, tracking of management practices implementation, and providing information on special studies of management practice efficacy.

Individual surveys to document current management practice implementations and assess future planned implementations are in the process of being filled out and will be completed by the end of May 2010. The Coalition will then assess these results and plan the final stages of outreach including re-contacting growers to identify newly implemented practices and future monitoring to evaluate water quality improvement.

Low Priority Subwatersheds

Merced River @ Santa Fe Drive

Merced River @ Santa Fe Drive (33,421 irrigated acres) – This water body is designated as a major water body and is 303d listed. It was selected as an integrator site for several of the drains and tributaries in the vicinity. The Merced River originates in the high Sierra encountering several dams and impoundments as it flows west. The Merced River eventually drains into the San Joaquin River near Hatfield State Park. Upstream agriculture includes some field crops in the immediate vicinity of the river and deciduous nuts, primarily almonds.

Merced River @ Santa Fe Drive is a Core monitoring location. The high load associated with one of the three prior exceedances of the chlorpyrifos WQTL and the concomitant toxicity to *C. dubia* (the third since 2004) initiated an effort to source the chlorpyrifos including a review of pesticide use reports and meetings with growers in the subwatershed. Based on information gained from these efforts, the Coalition decided to collect upstream samples in 2009 during months of past chlorpyrifos exceedances in the Merced River; the Coalition also collected samples for chlorpyrifos at the downstream location during the same events. It is believed that the chlorpyrifos is entering the Merced River via Dry Creek (not Dry Creek @ Wellsford Rd) several miles upstream. Since the toxicity to *Ceriodaphnia dubia* was associated with chlorpyrifos in the water, the Coalition is focusing its efforts in further identifying the sources of the chlorpyrifos which will aid in future outreach within this large subwatershed.

Other constituents that have exceeded their respective WQTLs at the Merced River sampling site are DO (September and October, 2009), lead (February 2007 and January 2008), and toxicity to *Selenastrum capricornutum* (March 2005).

Merced River @ Santa Fe Drive was monitored as a normal monitoring location from January to December 2009. Merced River @ Santa Fe Drive will be monitored as a Core Monitoring site in 2010.

Dry Creek @ Oakdale Rd (upstream of Merced River)

Dry Creek @ Oakdale Rd (13,564 irrigated acres) – Dry Creek flows into Merced River at Oakdale Ave. The agriculture draining into Dry Creek consists primarily of deciduous trees.

Dry Creek @ Oakdale Rd is a MPM location. This upstream location was selected based on a review of PUR data indicating likely upstream sources. The upstream monitoring site was selected to cut the watershed into smaller areas which allows for an analysis of the contribution of each portion of the watershed to the load measured at the Merced River @ Santa Fe Dr site. Management Plan Monitoring for chlorpyrifos occurred at this

site during the months of November and December of 2009. However, the site was dry during both sampling events and thus no sampling data exists for this site to date.

Miles Creek @ Reilly Rd

Miles Creek @ Reilly Rd (9,840 irrigated acres) – Miles Creek is located just north of Duck Slough and drains into Owen’s Creek. The primary agriculture includes field crops, deciduous nuts and fruit, pasture and truck/nursery/berry. Within the subwatershed are also urban drainages, dairies and hay, and pasture lands.

Miles Creek @ Reilly Rd was a new site in May 2007. In 2008, toxicity occurred for the second time to *Selenastrum capricornutum* and *Ceriodaphnia dubia*, and exceedances of WQTLs occurred twice for chlorpyrifos and four times for copper. The site was included in the ESJWQC Management Plan in 2009. During the irrigation season of 2009, Miles Creek was monitored for *Selenastrum capricornutum* in April, for copper in July and August, and for chlorpyrifos July through September. Chlorpyrifos was the only exceedance of the WQTL in July 2009.

Miles Creek will be monitored as an Assessment Monitoring location in 2015 and 2016.