

N O R T H A M E R I C A N JOURNAL OF FISHERIES MANAGEMENT

2002 303(d) List Update

Ref. # 75

Volume 16

November 1996

Number 4

North American Journal of Fisheries Management 16:693-727, 1996
© Copyright by the American Fisheries Society 1996

Channel Suspended Sediment and Fisheries: A Synthesis for Quantitative Assessment of Risk and Impact

CHARLES P. NEWCOMBE

Habitat Protection Branch, Ministry of Environment, Lands and Parks
Victoria, British Columbia V8V 1X4, Canada

JORGEN O. T. JENSEN

Science Branch, Pacific Biological Station, Department of Fisheries and Oceans
Nanaimo, British Columbia V9R 5K6, Canada

Abstract.—Our meta-analysis of 80 published and adequately documented reports on fish responses to suspended sediment in streams and estuaries has yielded six empirical equations that relate biological response to duration of exposure and suspended sediment concentration. These equations answer an important need in fisheries management; quantifying the response of fishes to suspended sediment pollution of streams and estuaries has been difficult historically, and the lack of a reliable metric has hindered assessment for risk and impact for fishes subjected to excess sedimentation. The six equations address various taxonomic groups of lotic, lentic, and estuarine fishes, life stages of species within those groups, and particle sizes of suspended sediments. The equations all have the form

$$z = a + b(\log_e x) + c(\log_e y);$$

z is severity of ill effect, x is duration of exposure (h), y is concentration of suspended sediment (mg SS/L), a is the intercept, and b and c are slope coefficients. The severity of ill effect (z) is delineated semiquantitatively along a 15-point scale on which is superimposed four "decision" categories ranging from no effect through behavioral and sublethal effects to lethal consequences (a category that also includes a range of para-lethal effects such as reduced growth rate, reduced fish density, reduced fish population size, and habitat damage). The study also provided best available estimates of the onset of sublethal and lethal effects, and it supported the hypothesis that susceptible individuals are affected by sediment doses (concentration \times exposure duration) lower than those at which population responses can be detected. Some species and life stages show "ultrasensitivity" to suspended sediment. When tested against data not included in the analysis, the equations were robust. They demonstrate that meta-analysis can be an important tool in habitat impact assessment.

While it is now generally accepted that the severity of effect of suspended sediment pollution on fish increases as a function of sediment concentration and duration of exposure, or dose (the product of concentration and exposure time), attempts to document the dose-response relationship

for sediment and aquatic organisms have been limited in several ways. First, initial analyses were based on pooled data (Newcombe 1986; Newcombe and MacDonald 1991). Second, the database available for those analyses embraced a wide taxonomic range from phytoplankton to fish. Third,

the database contained little information about particular species and life stages. The resulting dose-response model for aquatic ecosystems (Newcombe 1986; Newcombe and MacDonald 1991) established a general principle, but this model was held to be too imprecise to help fishery and habitat managers address local sediment problems (Gregory et al. 1993).

In an effort to refine the general dose-response model, MacDonald and Newcombe (1993) extracted and analyzed data for juvenile salmon from the recent literature. These data yielded an equation similar to the one for pooled data, but the two curves differed in important ways. This finding established a need to revisit the dose-response database so that models could be tailored to particular groups of fishes as functions of taxonomic group, natural history, life history phase, and predominant sizes of the sediment particles responsible for ill effects (Newcombe 1994). We have endeavored to meet this need and present a meta-analytic synthesis of dose-response data in this paper. Insofar as this research provides new understanding of channel sediment impacts, it leads to discussion of potential changes in the methods and goals of quantitative impact assessment. Specifically, the results (i) suggest the need to change the methods of data collection for environmental law enforcement, (ii) demonstrate the value of meta-analysis as a research method in fisheries habitat impact assessment, and (iii) prompt an expression of concern about land use practices and protection of instream, riparian, and upland zones.

Methods

This study is based on 264 data triplets consisting of (i) suspended sediment concentration, (ii) duration of exposure, and (iii) severity of ill effect for fishes. These data were taken from a comprehensive literature review (Newcombe 1994; Newcombe et al. 1995). Supporting data extracted from the review included taxonomic group, species of fish, natural history, life history phase, and sediment particle size range.

We define dose as concentration of suspended sediment (SS) times duration of exposure; dose has the units $\text{mg SS}\cdot\text{h}\cdot\text{L}^{-1}$. The natural logarithm of dose is termed the stress index (Newcombe 1986, 1994; Newcombe and MacDonald 1991; MacDonald and Newcombe 1993). Response is the severity of ill effect, described below. The dose-response matrix, which is the basis of data presentation in this report, encompasses all combinations of sediment concentration (1–500,000 mg SS/L) and ex-

TABLE 1.—Scale of the severity (SEV) of ill effects associated with excess suspended sediment.

SEV	Description of effect
Nil effect	
0	No behavioral effects
Behavioral effects	
1	Alarm reaction
2	Abandonment of cover
3	Avoidance response
Sublethal effects	
4	Short-term reduction in feeding rates; short-term reduction in feeding success
5	Minor physiological stress; increase in rate of coughing; increased respiration rate
6	Moderate physiological stress
7	Moderate habitat degradation; impaired homing
8	Indications of major physiological stress; long-term reduction in feeding rate; long-term reduction in feeding success; poor condition
Lethal and para-lethal effects	
9	Reduced growth rate; delayed hatching; reduced fish density
10	0–20% mortality; increased predation; moderate to severe habitat degradation
11	>20–40% mortality
12	>40–60% mortality
13	>60–80% mortality
14	>80–100% mortality

posure duration (1–35,000 h). Except when it refers specifically to duration, we use “exposure” broadly to include dose, particle size, and other potential contributors to stress on fishes. In most cases, data on particle shape and roughness and on water temperature were lacking.

Severity-of-Ill-Effect Scale

As before (MacDonald and Newcombe 1993; Newcombe 1994) and in a nearly identical way, we scored qualitative response data along a semi-quantitative ranking scale (Table 1). Superimposed on a 15-point scale (0–14) were four major classes of effect: (i) nil effect, (ii) behavioral effects, (iii) sublethal effects (a category that also includes effects such as short-term reduction in feeding success), and (iv) lethal effects (direct mortality, or its para-lethal surrogates—reduced growth, reduced fish density, habitat damage such as reduced porosity of spawning gravel, delayed hatching, and reduction in population size). When these various effects could be compared directly, pollution episodes associated with sublethal or lethal effects

also degra
which is
are group
between
tality, we
severity-
represent

We no
the class
ary betw
in feedi
long-ter
800 h an
to be su
can be l
as paral
change
of food

Along
from m
characte
both of
tations
of strea
ment o
1961),

(McLea
(Swens
degrad:
1977b);

structur
al. 198
Coats
excess
habitat
cess of
low sil
(Berkn

Hab
dence
life cy
result

Unive
(ii) av
1984a
and re
al. 19
ulatio:
1977)

(Ham
and V
behav
al. 19

Rel

also degraded habitat and reduced population size, which is why these seemingly disparate ill effects are grouped together in the hierarchy. For events between the extremes of nil effect and 100% mortality, we assumed for modeling purposes that the severity-of-ill effects (SEV for "severity") scale represents proportional differences in true effects.

We now incorporate all feeding reductions in the class of sublethal effects, and we set the boundary between short-term and long-term reductions in feeding success at 2 h. In practice, reports of long-term disruption of feeding rates encompass 800 h and more. We consider all feeding reductions to be sublethal effects (unless feeding reductions can be linked to slow growth when we treat them as para-lethal effects) because they reflect less a change in fish behavior than reduced availability of food and reduced visual hunting range.

Along the SEV scale, habitat damage ranges from moderate to severe. Habitat damage can be characterized in biological or physical terms or both of these in conjunction. Biological manifestations of habitat damage include underutilization of stream habitat (Birtwell et al. 1984), abandonment of traditional spawning habitat (Hamilton 1961), displacement of fish from their habitat (McLeay et al. 1987), and avoidance of habitat (Swenson 1978). Physical manifestations include degradation of spawning habitat (Slaney et al. 1977b; Cederholm et al. 1981), damage to habitat structure (Newcomb and Flagg 1983; Menzel et al. 1984), and loss of habitat (Menzel et al. 1984; Coats et al. 1985). Biophysical manifestations of excess SS are reported (in one typical example) as habitat degradation that reduces the relative success of one or more fish species that depend on low siltation rates and silt-free (<3% silt) riffles (Berkmann and Rabeni 1987).

Habitat degradation can be inferred by (i) evidence of increased mortality at any stage in a fish's life cycle (egg-to-fry survival may decrease as a result of increased sedimentation: J. LaPerriere, University of Alaska, personal communication), (ii) avoidance behavior by fishes (Suchanek et al. 1984a, 1984b), (iii) reduced abundance of insects and reduced quality of rearing habitat (Slaney et al. 1977b), (iv) decreased size of zoobenthic populations (Gammon 1970; Rosenberg and Snow 1977), (v) reduced utility of spawning habitat (Hamilton 1961), (vi) delayed hatching (Schubel and Wang 1973), and (vii) disruption of homing behavior and home water preference (Brannon et al. 1981; Whitman et al. 1982).

Relative severity of habitat damage is a contin-

uum on a two-dimensional plane (SS concentration \times duration of SS exposure) in which an event may be minor (ephemeral or low SS concentration or both), or major (long term or high SS concentration or both), or anywhere between these extremes. Severe habitat damage has been described by various authors, some of whom used aquatic invertebrates as indicators (Herbert and Richards 1963; Vaughan 1979; Vaughan et al. 1982; Menzel et al. 1984; Wagener and LaPerriere 1985). Severity of habitat damage caused by excess SS sometimes has been reported in terms of the length of time required for the stream to return to its natural state—sometimes as long as 15–20 years (estimated) after extensive coal mining (Vaughan et al. 1982).

The distinction between moderate and severe habitat damage is a matter of degree that still has not been delineated exactly. Severe habitat damage can be characterized in its extreme by the absence of fish where fish normally are found or by substantial reduction in fish population size, as was documented for brown trout by Herbert et al. (1961). (Scientific names of fish species are given in Table 2.) A pollution event that results in the deposition of suspended sediment in or on spawning habitat during egg incubation might be considered "moderately severe" if the area affected were a small portion of the total available. On the other hand, chronic or acute SS pollution that causes substantial reduction in the size of riverine fish populations (Herbert et al. 1961; Stober et al. 1981) should be considered to represent "severe" habitat damage. Likewise, major SS pollution that results in extensive deposition of sediment on spawning grounds should be characterized as severe habitat damage because its effects could reduce the strength of an entire year-class.

Habitat damage is a valid description of the harm caused by SS pollution, but it is probably an abstraction insofar as ill effects operate on one or more life stages of a fish's life cycle. Age-specific morbidity and mortality rates are fundamental to the notion of habitat damage. For example, habitat damage may manifest itself as foregone opportunity for fish to use a portion of a stream. Reduced suitability of habitat could result in increased age-specific morbidity and mortality rates, or both, depending on the focus and methods of a study. Habitat damage, therefore, should be seen as an accumulative measure of numerous (potentially undocumented) ill effects at various stages in a fish's life cycle. It is a unique phenomenon in that it can only be studied in the field (in contrast to direct

effects—age-specific morbidity and mortality, for example—that can be studied in the laboratory as well as in the field). Thus the documented harm caused by excess SS—especially when it is not known by direct observation to have caused an increase in morbidity or mortality rates—can reasonably be characterized in more general terms as habitat damage.

Model Formulation

From the expanded database (see Appendix Table A.1), six groupings of fish data were identified for which sample sizes were large enough to support modeling. The six groupings arose from various combinations of four attributes: taxonomic group, life stage, life history, and particle size of suspended sediment.

Taxonomy.—Salmonids (family Salmonidae) were distinguished from nonsalmonids, although several groupings were not exclusively one or the other.

Life stage.—Life stages were allocated among four categories: *eggs*, *larvae* (recently hatched fish, including yolk-sac fry, that had not passed through final metamorphosis); *juveniles* (fish, including fry, parr, and smolts, that had passed through larval metamorphosis but were sexually immature), and *adults* (mature).

Life history.—Estuarine species were categorized separately from anadromous and freshwater species, although these two groups were combined for early life stages.

Sediment particle size.—The predominant sizes of suspended sediment particles reported in the database literature ranged up to 250 μm . We collated sizes into two categories separated at 75 μm . *Fine* particles were smaller than 75 μm , small enough to pass through gill membranes into interlamellar spaces of gill tissue. This category includes clay, silt, and very fine sand particles (Agriculture Canada 1974). *Coarse* particles were 75–250 μm in diameter, large enough to cause mechanical abrasion of gills. This size range includes very fine to fine sand particles.

The six data groups for which we developed models follow. Species in each group are listed in Table 2.

Group 1: juvenile and adult salmonids; particle sizes 0.5–250 μm .—Group 1 ($N = 171$ studies or experimental units) includes Atlantic and Pacific salmon, trout, Arctic grayling, mountain whitefish, and rainbow smelt (a nonsalmonid). Some studies dealt with fine sediment as categorized above, some with coarse sediment, and some with both.

TABLE 2.—Common and scientific names of fish species and other taxa mentioned in this paper and the sediment effects model(s) to which they contributed. Species without a model number were not used in any model.

Common name	Scientific name	Model
Anchovy (bay)	<i>Anchoa mitchilli</i>	5 ^a
Bass (largemouth)	<i>Micropterus salmoides</i>	6
Bass (smallmouth)	<i>Micropterus dolomieu</i>	
Bass (striped)	<i>Morone saxatilis</i>	4,5
Bluegill	<i>Lepomis macrochirus</i>	6
Carp (common)	<i>Cyprinus carpio</i>	6
Cunner	<i>Tautoglabrus adspersus</i>	5
Darters	Percidae; includes <i>Semotilus</i> <i>atromaculatus</i> ^b	6
Fish	(Genus and species obscure)	5
Fish (warmwater)	(Genus and species obscure)	5,6
Goldfish	<i>Carassius auratus</i>	6
Grayling (Arctic)	<i>Thymallus arcticus</i>	1–4
Herring (Atlantic)	<i>Clupea harengus</i>	4,5 ^a
Herring (lake)	<i>Coregonus artedii</i>	4
Herring (Pacific)	<i>Clupea pallasii</i>	4
Hogchoker	<i>Trinectes maculatus</i>	5
Killifish (striped)	<i>Fundulus majalis</i>	5
Menhaden (Atlantic)	<i>Brevoortia tyrannus</i>	5 ^a
Minnnow (sheepshead)	<i>Cyprinodon variegatus</i>	5 ^a
Mummichog	<i>Fundulus heteroclitus</i>	5
Perch (white)	<i>Morone americana</i>	4,5
Perch (yellow)	<i>Perca flavescens</i>	4
Rasbora (harlequin)	<i>Rasbora heteromorpha</i>	5
Salmon	(Genus and species obscure)	1,2,4
Salmon (Atlantic)	<i>Salmo salar</i>	1,2
Salmon (chinook)	<i>Oncorhynchus tshawytscha</i>	1–3
Salmon (chum)	<i>Oncorhynchus keta</i>	1,3,4
Salmon (coho)	<i>Oncorhynchus kisutch</i>	1,3,4
Salmon (Pacific)	<i>Oncorhynchus</i> spp.	1,2
Salmon (sockeye)	<i>Oncorhynchus nerka</i>	1–3
Shad (American)	<i>Alosa sapidissima</i>	4,5
Silverside (Atlantic)	<i>Menidia menidia</i>	5 ^a
Smelt (rainbow)	<i>Osmerus mordax</i>	1,2
Spot	<i>Leiostomus xanthurus</i>	5 ^a
Steelhead	<i>Oncorhynchus mykiss</i> (anadromous)	1–4
Stickleback (fourspine)	<i>Apeltes quadracus</i>	5 ^a
Stickleback (threespine)	<i>Gasterosteus aculeatus</i>	5
Sunfish (green)	<i>Lepomis cyanellus</i>	6
Sunfish (reardear)	<i>Lepomis microlophus</i>	6
Toadfish (oyster)	<i>Opsanus tau</i>	5
Trout	(Genus and species obscure)	1,2,4
Trout (brook)	<i>Salvelinus fontinalis</i>	1–3
Trout (brown)	<i>Salmo trutta</i>	1,2
Trout (cutthroat)	<i>Oncorhynchus clarki</i>	1,2
Trout (lake)	<i>Salvelinus namaycush</i>	1,2
Trout (rainbow)	<i>Oncorhynchus mykiss</i>	1–4
Trout (sea)	(Genus and species obscure)	1,2
Whitefish (lake)	<i>Coregonus clupeaformis</i>	1,2
Whitefish (mountain)	<i>Prosopium williamsoni</i>	1,2

^a A relatively sensitive species used in the empirical model for estuarine species.

^b Creek chubs are included with darters here because the relevant study (Vaughan et al. 1978) referred to reduced fish abundance in streams where chubs and darters were reported to live.

TABLE 3
Life stage
Ecology

Life stage
Life history
Sediment

Intercept
Slope of 1
Slope of 1

Estuaries
Deterministic
Probabilistic
Sample size

Life stage
Life history
Life history
Life history

Group 1
 μm Group

Group 1
 μm Group

Life history
Life history

Group 4
salmonids,
300 μm

Group 5
very fine to
coarse particles

Life history
Life history

Group 6
life stage
both fine to
coarse

Group 7
both fine to
coarse

Group 8
both fine to
coarse

Group 9
both fine to
coarse

Group 10
both fine to
coarse

TABLE 3.—Attributes, slopes and coefficients, and statistics of six models that relate severity of ill effect on fishes (z , 15-point scale) to duration of exposure (x , h) and concentration of suspended sediment (y , mg/L) in the form $z = a + b(\log_e x) + c(\log_e y)$.

fish species
e sediment
d. Species
model.

Model	Term	Model					
		1	2	3	4	5	6
5 ^a							
6							
		Attributes					
	Taxon ^a	S	S	S	S + N	N	N
4,5	Life stage ^b	J + A	A	J	E + L	A	A
6	Life history ^c	FW	FW	FW	FW + ES	ES	FW
6	Sediment particle size ^d	F to C	F to C	F	F	F	F
5							
6							
		Slopes and coefficients					
	Intercept (a)	1.0642	1.6814	0.7262	3.7466	3.4969	4.0815
	Slope of $\log_e x$ (b)	0.6068	0.4769	0.7034	1.0946	1.9647	0.7126
5	Slope of $\log_e y$ (c)	0.7384	0.7565	0.7144	0.3117	0.2669	0.2829
		Statistics					
5,6	Coefficient of determination ^e (r^2)	0.6009	0.6173	0.5984	0.5516	0.6200	0.6998
6	F -statistic	130.28	52.37	82.00	28.03	24.50	27.42
1-4	Probability (P)	<0.01	<0.01	<0.01	<0.01	<0.01	<0.01
4,5 ^a	Sample size (N)	171	63	108	43	28	22

^a S = salmonids (predominantly); N = nonsalmonids.
^b A = adults; J = juveniles; L = larvae; E = eggs.
^c FW = freshwater and anadromous; ES = estuarine.
^d F = fine (predominantly <75 μm); C = coarse (75-250 μm).
^e Corrected for degrees of freedom.

Group 2: adult salmonids; particle sizes 0.5-250 μm .—Group 2 ($N = 63$) is a subset of group 1.

Group 3: juvenile salmonids; particle sizes 0.5-75 μm .—Group 3 ($N = 108$) is a subset of group 1. In a few cases, sediment sizes were as large as 150 μm .

Group 4: eggs and larvae of salmonids and nonsalmonids; particle sizes 0.5-75 μm .—Group 4 ($N = 43$) includes salmonids that do not bury their eggs. Nonsalmonids comprise species that spawn in rivers, lakes, and estuaries. Sediment sizes exceeded 75 μm in a few studies.

Group 5: adult estuarine nonsalmonids; particle sizes 0.5-75 μm .—Group 5 ($N = 28$) includes several species believed to be particularly sensitive to the effects of suspended sediment; these are footnoted in Table 2. Some test sediments exceeded 75 μm .

Group 6: adult freshwater nonsalmonids; particle sizes 0.5-75 μm .—Group 6 ($N = 22$) includes both lentic and lotic species. Particle sizes exceeded 75 μm in some cases.

For each group, the severity of effect (SEV, 15-point scale, 0-14) was regressed on suspended sediment dose (exposure duration [ED, h] and suspended sediment concentration [mg SS/L]). Preliminary analyses indicated that logarithmic transformations of ED and concentration provided suitably linear relations of the form

$$SEV = a + b(\log_e ED) + c(\log_e \text{mg SS/L});$$

intercepts (a) and slope coefficients (b and c) emerged from the fitting exercise. Commercial software was used for the regressions (TableCurve 3D; Jandel Scientific). Coefficients of determination (r^2) were adjusted for degrees of freedom ($r^2 = 1 - [\text{sum of squares due to error}]/[\text{sum of squares around the mean}]$). The software also generated F -statistics, P -values, and 95% confidence intervals around the SEVs. Although arithmetic values for exposure duration and concentration are also given in the Results and in the Appendix, the models we present are based on logarithmic transformations.

The regressions, having been fitted to the data, become predictive models of the form

$$z = a + b(\log_e x) + c(\log_e y),$$

for which z is calculated severity of ill effect (SEV), x is an estimate of exposure duration (ED), and y is the concentration of the (estimated) predominant suspended sediment size (mg SS/L). These predictive models are numbered 1-6 to correspond with the data groupings already described. Because of scatter even in the fitted data, the predictive equations can yield severity-of-ill-effect (z) values greater than 14, which already includes the

model for

relevant
distance in

Juvenile and Adult Salmonids

Duration of exposure to SS (\log_e hours)




FIGURE 1.—(A) Average empirical severity-of-ill-effect scores for juvenile and adult salmonids (freshwater, group 1) in the matrix of suspended sediment (SS) concentration and duration of exposure. Both matrix axes are expressed in logarithmic and absolute terms. Dashes mean "no data." Shaded bands denote inferred (by manual interpolation) thresholds of sublethal effects (shading without a border) and lethal effects (shading with a border; see Table 1 for criteria). (B, upper matrix) Severity-of-effect scores calculated by model (1) (Table 3). Severity-of-ill-effect calculations are based on the logarithmic values shown on the axes of the matrix. Shaded areas represent extrapolations beyond empirical data; extrapolations have been capped at 14 (upper limit of the effects scale; Table 1), although higher values are possible. Diagonal terraced lines denote thresholds of sublethal effects (lower left) and lethal effects (middle diagonal) delineated by the model with reference to Table 1. (B, lower matrix) Half-95% confidence intervals around calculated severity-of-effect scores. Shaded areas denote half-intervals greater than 1.0.

most serious effects to be measured (100% mortality; catastrophic habitat degradation).

Data Presentation

Empirical data.—Severity-of-ill-effect values for each of the six data groups are presented as rounded averages in the cells of dose matrixes whose axes are concentration of suspended sediment and duration of exposure (panel A of the figure for each group). Maximum possible duration of exposure in the matrix is 48 months (\log_e [hours] = 10.4999). All but one of the matrixes show a maximum possible suspended sediment concentration of 268,337 mg/L (\log_e [mg SS/L] = 12.4999). The exception—adult estuarine fishes—has a maximum possible concentration of 729,416 mg SS/L (\log_e [mg SS/L] = 13.4999).

Displayed logarithmic values of duration and

concentration are the midrange values. Thus the range of logarithmic values represented by a row or a column in the figures is approximately the value ± 0.4999 in logarithmic units (take antilogarithms for absolute values and their ranges). The accompanying confidence values are one-half the 95% confidence intervals around z .

Cells of a matrix that contain data form a cluster of "populated" cells. The imaginary "tight-string" polygon that encompasses all the populated cells in a matrix is the "data envelope." Typically, some cells within a data envelope are unpopulated. For predictive purposes, values are assigned to these cells by interpolation. Empty cells outside the envelope are given values by extrapolation. Interpolations are considered to have greater intrinsic reliability than extrapolations because they can be compared more easily with known data.

Threshold severity-of-estimation rations tha (panel A c purpose, u: lope are as Thresholds

Juvenile and Adult Salmonids

Duration of exposure to SS (log_e hours)

0	1	2	3	4	5	6	7	8	9	10
---	---	---	---	---	---	---	---	---	---	----

(B) Average severity-of-ill-effect scores (calculated)

Concentration (mg SS/L)	162755	10	11	11	12	12	13	14	14	-	-	-	12	(log _e mg SS/L)
	59874	9	10	10	11	12	12	13	13	14	-	-	11	
	22026	8	9	10	10	11	11	12	13	13	14	-	10	
	8103	8	8	9	10	10	11	11	12	13	13	14	9	
	2981	7	8	8	9	9	10	11	11	12	12	13	8	
	1097	6	7	7	8	9	9	10	10	11	12	12	7	
	403	5	6	7	7	8	9	9	10	10	11	12	6	
	148	5	5	6	7	7	8	8	9	10	10	11	5	
	55	4	5	5	6	6	7	8	8	9	9	10	4	
	20	3	4	4	5	6	6	7	8	8	9	9	3	
	7	3	3	4	4	5	6	6	7	7	8	8	2	
	3	2	2	3	4	4	5	5	6	7	7	8	1	
1	1	2	2	3	3	4	5	5	6	7	7	0		
	1	3	7	1	2	6	2	7	4	11	30			
	Hours			Days			Weeks		Months					

Half-95% confidence intervals (±) around calculated severity-of-ill-effect scores (above)

162755	0.9	0.8	0.8	0.8	0.7	0.7	0.8	0.8	-	-	-	12
59874	0.8	0.7	0.7	0.6	0.6	0.6	0.6	0.7	0.7	-	-	11
22026	0.7	0.6	0.6	0.5	0.5	0.5	0.6	0.6	0.7	0.7	-	10
8103	0.6	0.6	0.5	0.4	0.4	0.4	0.5	0.5	0.6	0.7	0.8	9
2981	0.6	0.5	0.4	0.4	0.4	0.4	0.4	0.5	0.5	0.6	0.7	8
1097	0.6	0.5	0.4	0.3	0.3	0.3	0.4	0.4	0.5	0.6	0.7	7
403	0.6	0.5	0.4	0.3	0.3	0.3	0.4	0.5	0.5	0.6	0.7	6
148	0.6	0.5	0.4	0.4	0.4	0.4	0.4	0.5	0.6	0.7	0.8	5
55	0.6	0.6	0.5	0.5	0.5	0.5	0.5	0.6	0.6	0.7	0.8	4
20	0.7	0.7	0.6	0.6	0.6	0.6	0.6	0.7	0.7	0.8	0.9	3
7	0.8	0.7	0.7	0.7	0.7	0.7	0.7	0.8	0.8	0.9	1.0	2
3	0.9	0.8	0.8	0.8	0.8	0.8	0.8	0.9	0.9	1.0	1.0	1
1	1.0	1.0	0.9	0.9	0.9	0.9	0.9	1.0	1.0	1.1	1.1	0
	1	3	7	1	2	6	2	7	4	11	30	
	Hours			Days			Weeks		Months			

FIGURE 1.—Continued.

Thresholds of ill effect.—Display of empirical severity-of-effect scores in the dose matrix permits estimation of the minimum concentrations and durations that trigger sublethal and lethal effects (panel A of the figure for each group). For this purpose, unpopulated cells within the data envelope are assigned values by manual interpolation. Thresholds thus estimated from empirical data of-

ten are lower than thresholds predicted by regressions fit to meta-analytical data. We interpret "empirical thresholds" as an approximated response of the more "sensitive" individuals within a species group.

Predictions of ill effect.—The regression equation fitted to each of the six data groups provides predictions of response within the matrix of con-

er, group
pressed
olation)
Table 1
ill-effect
extrap-
Table 1),
left) and
alf-95%
ter than

hus the
y a row
ely the
antilog-
es). The
half the

a cluster
-string"
cells in
y, some
ted. For
to these
the en-
Interpo-
insic re-
can be

Adult Salmonids

Duration of exposure to SS (log₁₀ hours)




FIGURE 2.—Empirical severity-of-ill-effect scores for adult salmonids (freshwater, group 2) and scores (with half-95% confidence intervals) predicted by model (2). Conventions are those of Figure 1.

centration and duration of exposure (panel B of the figure for each group). Each prediction is accompanied by half-95% confidence intervals.

Each prediction matrix is divided into a maximum of three zones by terraced lines separating behavioral, sublethal, and lethal responses. We compare these modeled thresholds to empirical ones to discern responses of "sensitive" individuals within each species group.

Results

Dose-response models fitted to the empirical data groups were all highly significant ($P < 0.01$) and accounted for 55-70% of the variances (Table 3). Averaged empirical data on which the models are based are displayed in panel A of Figures 1-6. Panel B of Figures 1-6 gives the model-generated responses (and confidence intervals) for each cell of the dose-response matrixes. These panels provide a set of "look-up tables" suitable for field use in impact assessment. Superimposed on them are predicted thresholds of sublethal and lethal effects based on the response categories in Table 1. Response surfaces resulting from the models are shown in Figures 7-12. Data are derived from sources listed in the Appendix.

Group 1: Juvenile and Adult Salmonids

Average empirical severity-of-ill-effect data for group 1 fill 56 of the 143 available cells (Figure 1A). Data are widely distributed, but thresholds for the onset of sublethal and lethal ill effects can be inferred within broad limits, based on manual interpolations within the data envelope (see gray-shaded zones without and with borders).

The full matrix array of severity scores predicted by model 1 (Table 3, Figure 1B) shows regular increases of response intensity with sediment dose, as expected. Predicted thresholds of sublethal and lethal effects (terraced diagonals) have similar orientations to those inferred from empirical data, but they generally occur at higher sediment doses.

Group 2: Adult Salmonids

Group 2 data fill 36 widely scattered cells of the 143 available in the empirical matrix (Figure 2A). The thresholds of lethal effect predicted by model 2 (Table 3; Figure 2B) are similar to the empirically inferred threshold (Figure 2A), but predicted sublethal effects emerge at slightly lower sediment doses than implied by empirical data.

Group 3: Juvenile and Adult Salmonids

Average 37 cells, m... ratios of 1... for adult s... 3: Table 3; thresholds... ical ones f...

Adult Salmonids

Duration of exposure to SS (log_e hours)

0	1	2	3	4	5	6	7	8	9	10
---	---	---	---	---	---	---	---	---	---	----

(B) Average severity-of-ill-effect scores (calculated)

Concentration (mg SS/L)	162755	11	11	12	12	13	13	14	14	-	-	-	12
	59874	10	10	11	11	12	12	13	13	14	14	-	11
	22026	9	10	10	11	11	12	12	13	13	14	14	10
	8103	8	9	9	10	10	11	11	12	12	13	13	9
	2981	8	8	9	9	10	10	11	11	12	12	13	8
	1097	7	7	8	8	9	9	10	10	11	11	12	7
	403	6	7	7	8	8	9	9	10	10	11	11	6
	148	5	6	6	7	7	8	8	9	9	10	10	5
	55	5	5	6	6	7	7	8	8	9	9	9	4
	20	4	4	5	5	6	6	7	7	8	8	9	3
	7	3	4	4	5	5	6	6	7	7	7	8	2
	3	2	3	3	4	4	5	5	6	6	7	7	1
	1	2	2	3	3	4	4	5	5	5	6	6	0
	1	3	7	1	2	6	2	7	4	11	30		
	Hours			Days			Weeks		Months				

(log_e mg SS/L)

(log_e mg SS/L)

with half-

Half-95% confidence intervals (±)
around calculated severity-of-ill-effect scores (above)

162755	1.4	1.4	1.4	1.4	1.4	1.4	1.4	1.4	-	-	-	12
59874	1.3	1.3	1.3	1.3	1.3	1.3	1.3	1.3	1.3	1.3	-	11
22026	1.2	1.2	1.2	1.2	1.2	1.2	1.2	1.2	1.2	1.2	1.3	10
8103	1.1	1.1	1.1	1.1	1.1	1.1	1.1	1.1	1.1	1.1	1.2	9
2981	1.0	1.0	1.0	1.0	1.0	1.0	1.0	1.0	1.0	1.0	1.1	8
1097	1.0	1.0	1.0	1.0	1.0	1.0	1.0	1.0	1.0	1.0	1.1	7
403	0.9	0.9	0.9	0.9	0.9	0.9	0.9	0.9	0.9	0.9	1.1	6
148	1.0	1.0	1.0	1.0	1.0	1.0	1.0	1.0	1.0	1.0	1.1	5
55	1.0	1.0	1.0	1.0	1.0	1.0	1.0	1.0	1.0	1.0	1.1	4
20	1.1	1.1	1.1	1.1	1.1	1.1	1.1	1.1	1.1	1.1	1.2	3
7	1.2	1.2	1.2	1.2	1.2	1.2	1.2	1.2	1.2	1.2	1.3	2
3	1.3	1.3	1.3	1.3	1.3	1.3	1.3	1.3	1.3	1.3	1.4	1
1	1.4	1.4	1.4	1.4	1.4	1.4	1.4	1.4	1.4	1.4	1.5	0
	1	3	7	1	2	6	2	7	4	11	30	
	Hours			Days			Weeks		Months			

t data for
s (Figure
resholds
ffects can
a manual
see gray-

predicted
s regular
ent dose,
ethal and
milar ori-
data, but
doses.

FIGURE 2.—Continued.

Group 3: Juvenile Salmonids

Average severity-of-effect scores for group 3 fill 37 cells, most of them clustered at exposure durations of 1 h and 2 d to 7 weeks (Figure 3A). As for adult salmonids, predicted thresholds (model 3: Table 3; Figure 3B) were similar to empirical thresholds for lethal effects but lower than empirical ones for sublethal effects.

Group 4: Eggs and Larvae of Salmonids and Nonsalmonids

Average severity scores for eggs and larvae of salmonids and freshwater and estuarine nonsalmonids fill 23 cells (Figure 4A). Most data are clustered in the exposure interval of 1 d to 7 weeks. Sublethal effects thresholds were estimated empirically, but they were not recognized by model

ells of the
gure 2A).
by model
e empiri-
predicted
sediment

Juvenile Salmonids

Duration of exposure to SS (log₁₀ hours)




FIGURE 3.—Empirical severity-of-ill-effect scores for juvenile salmonids (freshwater, group 3) and scores (with half-95% confidence intervals) predicted by model (3). Conventions are those of Figure 1.

4 (Table 3; Figure 4B), which generated no severity score lower than 4. Empirical and predicted thresholds of lethal effect agreed well and occurred at relatively low doses.

Group 5: Adult Estuarine Nonsalmonids

Average severity-of-effect scores for at least 15 species of estuarine fishes filled 23 of the available 154 matrix cells (Figure 5A). Most of the data represent 1–6-d exposures.

Model 5 (Table 3) was developed for only the seven species represented by adequate data. These seven are believed to be relatively more sensitive to the ill effects of suspended sediment than the other species in the database (Table 2). Predicted thresholds of lethal effect (Figure 5B) tracked empirical thresholds well for exposure durations less than 1 d; both estimates indicated that lethal effects on those sensitive species result from short exposures to a wide range of sediment concentrations. Sublethal effect thresholds were considerably closer the origin in the predictive matrix than in the empirical matrix.

Group 6: Adult Freshwater Nonsalmonids

A relatively small sample of stream and still-water fishes in cold, temperate, and warmwater

environments provided average severity scores for 15 scattered matrix cells of the 143 available (Figure 6A). Model 6 (Table 3) generated lethal effects thresholds that agreed well with interpolations of empirical data for exposures of 7 d to 7 weeks (Figure 6B). Although sublethal thresholds could be inferred from empirical data, the model indicated that they lay beyond the matrix—below concentrations of 1 mg/L, exposure durations of 1 h, or both.

Response Surfaces

Dose–response surfaces based on models 1–6 are shown in Figures 7–12. We think it important to emphasize that only models (1), (3), and (4) address early life stages in some form. Many studies have shown that early stages (some stages of egg development through young juveniles) are more susceptible to toxicants and other pollutants than older juveniles and adults. The response surfaces (and prediction matrixes) should be judged by the data available to develop them.

Discussion

Fisheries biologists, habitat protection specialists, and enforcement officers in many parts of the world may find that the dose–response equations

generate daily validation patterns (iii) pot enforcer findings

Juvenile Salmonids

Duration of exposure to SS (log_e hours)

	0	1	2	3	4	5	6	7	8	9	10		
(B)	Average severity-of-ill-effect scores (calculated)												
Concentration (mg SS/L)	162755	9	10	11	11	12	13	14	14	-	-	-	12
	59874	9	9	10	11	11	12	13	14	14	-	-	11
	22026	8	9	9	10	11	11	12	13	13	14	-	10
	8103	7	8	9	9	10	11	11	12	13	13	14	9
	2981	6	7	8	9	9	10	11	11	12	13	13	8
	1097	6	6	7	8	9	9	10	11	11	12	13	7
	403	5	6	6	7	8	9	9	10	11	11	12	6
	148	4	5	6	6	7	8	9	9	10	11	11	5
	55	4	4	5	6	6	7	8	8	9	10	11	4
	20	3	4	4	5	6	6	7	8	8	9	10	3
	7	2	3	4	4	5	6	6	7	8	8	9	2
	3	1	2	3	4	4	5	6	6	7	8	8	1
1	1	1	2	3	4	4	5	6	6	7	8	0	
	1	3	7	1	2	6	2	7	4	11	30		
	Hours			Days			Weeks		Months				

(log_e mg SS/L)

Half-95% confidence intervals (±)
around calculated severity-of-ill-effect scores (above)

162755	1.3	1.2	1.1	1.1	1.0	1.0	1.0	1.1	-	-	-	12
59874	1.1	1.0	0.9	0.9	0.9	0.8	0.9	0.9	1.0	-	-	11
22026	1.0	0.9	0.8	0.7	0.7	0.7	0.7	0.8	0.9	1.0	-	10
8103	0.9	0.7	0.7	0.6	0.5	0.6	0.6	0.7	0.8	0.9	1.0	9
2981	0.8	0.7	0.6	0.5	0.4	0.5	0.5	0.6	0.7	0.8	1.0	8
1097	0.7	0.6	0.5	0.4	0.4	0.4	0.5	0.6	0.7	0.8	1.0	7
403	0.7	0.6	0.5	0.5	0.4	0.5	0.5	0.6	0.7	0.9	1.0	6
148	0.8	0.7	0.6	0.6	0.5	0.6	0.6	0.7	0.8	1.0	1.1	5
55	0.9	0.8	0.7	0.7	0.7	0.7	0.8	0.9	0.9	1.1	1.2	4
20	1.0	0.9	0.9	0.8	0.8	0.9	0.9	1.0	1.1	1.2	1.3	3
7	1.1	1.1	1.0	1.0	1.0	1.0	1.1	1.2	1.2	1.3	1.4	2
3	1.3	1.2	1.2	1.2	1.2	1.2	1.3	1.3	1.4	1.5	1.6	1
1	1.4	1.4	1.4	1.4	1.4	1.4	1.4	1.5	1.6	1.6	1.7	0
	1	3	7	1	2	6	2	7	4	11	30	
	Hours			Days			Weeks		Months			

FIGURE 3.—Continued.

generated in this study are useful additions to their daily work. The discussion below focuses on (i) validation of the models, (ii) the dose-response patterns of ultrasensitive species and life stages, (iii) potential new options in environmental law enforcement, (iv) the role of meta-analysis in the findings of this study, (v) possible directions of

future research, and (vi) implications of this study for ecosystem assessment.

Validation of the Models

Validation of the models in this study will rely on new studies that add to the data now available. Creation of new data—in sufficient volume for

res (with
ores for
le (Fig-
l effects
tions of
7 weeks
ls could
el indi-
ow con-
of 1 h,
els 1-6
portant
and (4)
y stud-
ages of
les) are
llutants
nse sur-
judged
special-
s of the
uations

Eggs and Larvae of Salmonids and Nonsalmonids

Duration of exposure to SS (\log_e hours)




FIGURE 4.—Empirical severity-of-ill-effect scores for eggs and larvae of salmonids and nonsalmonids (freshwater and estuarine, group 4) and scores (with half-95% confidence intervals) predicted by model (4). Conventions are those of Figure 1, except the model (B, upper matrix) recognized no threshold of sublethal effects.

testing and refinement of these models—is bound to be a slow process. However, in the brief time since the conclusion of the data-gathering phase of this study, some new data have emerged.

First, coho salmon fry (mean weight, 1.95 g; $N = 10$ fish), when exposed to suspended sediment at a concentration of 5,471 mg SS/L for 96 h, sustained a mortality rate of 10% after they had been held in water at 18.7°C and 9.7 mg O_2/L (J.O.T.J., unpublished data). This mortality rate expressed as a severity of ill effect (with reference to Table 1) is $SEV = 10$. Severity of ill effect as predicted by model 1 ($SEV = 0.7262 + 0.7034\log_e[96 h] + 0.7144(\log_e 5,471 \text{ mg SS/L})$) is 10.09. These values agree closely and tend to validate this model. Steelhead ($N = 10$), similarly exposed, had 0% mortality. This result too is consistent with the predictions of the model, because $SEV = 10$ represents 0–20% mortality, and the test fish exhibited behaviors of severe sublethal stress.

Second, a recent laboratory study of effects of suspended bentonite clay (1–5- μm diameters) on larval nonsalmonid fishes (smallmouth bass, largemouth bass, and bluegill) in warm water (20–25°C) has produced several sets of morbidity data (re-

duced growth rate) and mortality data that are highly consistent with the predictions of model (4) (J. Sweeten, Asherwood Environmental Learning Centre, personal communication).

Third, an inverse relationship has been documented between sediment concentrations in streams and maximum salmonid densities in fluvial habitats in British Columbia (Ptolemy 1993; R. A. Ptolemy, British Columbia Ministry of Environment, Lands and Parks, personal communication). For example, the density (number of fish per unit area) of juvenile chinook salmon and steelhead that rear in the turbid main stem of the Bella Coola River (British Columbia) is lower than would be expected in clear water. Rearing occurs in June, July, and August. During this time, turbidity averages 21 nephelometric units, suspended sediment concentration averages 61 mg SS/L, particle sizes are smaller than 75 μm , and the temperature range is 8–12°C). Reduced fish density is consistent with the range of ill effects—low parallel rankings—predicted by the models. These results tacitly acknowledge the role of excess sediment exposure—particularly concentration and duration—as a factor in the productivity of salmon streams. Two extenuating factors—relatively

small particle could explain the Bella Coola.

Fourth, juvenile rainbow trout, a to seek refuge—chinook salmon the upper Fraser

Eggs and Larvae of Salmonids and Nonsalmonids

Duration of exposure to SS (log_e hours)

	0	1	2	3	4	5	6	7	8	9	10		
(B)	Average severity-of-ill-effect scores (calculated)												
Concentration (mg SS/L)	162755	7	9	10	11	12	13	14	-	-	-	-	12
	59874	7	8	9	10	12	13	14	-	-	-	-	11
	22026	7	8	9	10	11	12	13	-	-	-	-	10
	8103	7	8	9	10	11	12	13	14	-	-	-	9
	2981	6	7	8	10	11	12	13	14	-	-	-	8
	1097	6	7	8	9	10	11	12	14	-	-	-	7
	403	6	7	8	9	10	11	12	13	14	-	-	6
	148	5	6	7	9	10	11	12	13	14	-	-	5
	55	5	6	7	8	9	10	12	13	14	-	-	4
	20	5	6	7	8	9	10	11	12	13	-	-	3
	7	4	5	7	8	9	10	11	12	13	14	-	2
	3	4	5	6	7	8	10	11	12	13	14	-	1
	1	4	5	6	7	8	9	10	11	13	14	-	0
	1	3	7	1	2	6	2	7	4	11	30		
	Hours			Days			Weeks		Months				

(log_e mg SS/L)

Half-95% confidence intervals (±)
around calculated severity-of-ill-effect scores (above)

162755	2.6	2.5	2.5	2.5	2.5	2.6	2.7	-	-	-	-	12
59874	2.3	2.2	2.1	2.1	2.1	2.2	2.3	-	-	-	-	11
22026	2.0	1.9	1.8	1.8	1.8	1.9	2.0	2.1	-	-	-	10
8103	1.8	1.6	1.5	1.5	1.5	1.5	1.6	1.8	-	-	-	9
2981	1.6	1.4	1.2	1.2	1.1	1.2	1.3	1.5	-	-	-	8
1097	1.4	1.2	1.0	0.9	0.8	0.9	1.0	1.2	-	-	-	7
403	1.4	1.1	0.9	0.7	0.6	0.7	0.8	1.0	1.3	-	-	6
148	1.4	1.2	0.9	0.7	0.6	0.6	0.7	0.9	1.2	-	-	5
55	1.6	1.3	1.1	0.9	0.7	0.7	0.8	0.9	1.1	-	-	4
20	1.8	1.5	1.3	1.1	1.0	0.9	1.0	1.1	1.2	1.5	-	3
7	2.0	1.8	1.6	1.4	1.3	1.2	1.2	1.3	1.4	1.6	-	2
3	2.3	2.1	1.9	1.7	1.6	1.6	1.5	1.6	1.7	1.8	-	1
1	2.6	2.4	2.2	2.1	2.0	1.9	1.9	1.9	2.0	2.1	-	0
	1	3	7	1	2	6	2	7	4	11	30	
	Hours			Days			Weeks		Months			

FIGURE 4.—Continued.

small particle size and relatively cool water—could explain the absence of direct lethality in the Bella Coola.

Fourth, juvenile salmonids (chinook salmon, rainbow trout, and mountain whitefish) are thought to seek refuge—an average of 9 d for age-0 wild chinook salmon—in a small nonnatal tributary of the upper Fraser River, perhaps to avoid unsuitable

rearing conditions created by high, naturally occurring sediment loads found in the main stem (Scrivener et al. 1993).

Although these recent findings tend to support the predictions of the models, the well-documented good health (as indicated by acceptable rates of growth and survival) among salmon juveniles in turbid estuarine waters remains unexplained.

Adult Estuarine Nonsalmonids

Duration of exposure to SS (\log_e hours)




FIGURE 5.—Empirical severity-of-ill-effect scores for adult nonsalmonids (estuarine, group 5) and scores (with half-95% confidence intervals) predicted by model (5). Conventions are those of Figure 1.

Considerations relevant to this "anomaly" include (i) the extremely fine texture of suspended sediment (generally much smaller than 75 μ m); (ii) the relatively cold water temperatures; (iii) the potential for favorable physicochemical effects such as flocculation, which could be enhanced by the chemistry of brackish water; (iv) beneficial behavioral adaptations of juvenile salmonids; and (v) the suitability of reedy habitat, where average sediment concentrations and average particle size may be further reduced below those found in traditional sampling sites.

Ultrasensitivity of Some Species and Life Stages

Rapid escalation of ill effects on eggs, larvae, and fry (Figures 4, 10) and on some adult fishes of the estuary (Figures 5, 11) as duration of sediment exposure increases suggests that the mechanisms of self-preservation in at least some estuarine fishes are easily overwhelmed by the presence of suspended sediment. This pattern implies the existence of an abrupt threshold concentration of suspended sediment leading to ill effects in ultrasensitive species and life stages.

If this inference is correct, these dose-response patterns might be explained in terms of the time

required to reach an end point (e.g., lethality), and might indicate that the physiological and physical processes involved in homeostasis are more sensitive to exposure time than to suspended sediment concentrations. It is reasonable to speculate further that the sequence of events leading to a lethal end point (for example, severely abraded gill tissue and associated loss of capacity for ion regulation), once triggered, would not easily be halted or reversed.

Environmental Enforcement Issues

Fisheries biologists and enforcement personnel can, as part of an investigation, document the sediment concentration and duration of exposure, and they can use these data to infer the most probable severity of impact. The dose-response equations alone are sufficient for this task. But the "look-up" tables (here, Figures 1-6, panels B) simplify the task even more; they are based on the equations, and they supply ranges of interpolation and extrapolation and confidence intervals. They make it possible for field workers readily to distinguish between minor and major events in the broad context established by the dose-response matrixes. This knowledge can contribute to decisions about

the need for gather physical severity of the inferences— field of change influence the g Impacts or of excess sec

Adult Estuarine Nonsalmonids

Duration of exposure to SS (log_e hours)

	0	1	2	3	4	5	6	7	8	9	10	
(B)	Average severity-of-ill-effect scores (calculated)											
Concentration (mg SS/L)	162755	7	9	11	13	-	-	-	-	-	-	12
	59874	6	8	10	12	14	-	-	-	-	-	11
	22026	6	8	10	12	14	-	-	-	-	-	10
	8103	6	8	10	12	14	-	-	-	-	-	9
	2981	6	8	10	12	13	-	-	-	-	-	8
	1097	5	7	9	11	13	-	-	-	-	-	7
	403	5	7	9	11	13	-	-	-	-	-	6
	148	5	7	9	11	13	-	-	-	-	-	5
	55	5	7	8	10	12	14	-	-	-	-	4
	20	4	6	8	10	12	14	-	-	-	-	3
	7	4	6	8	10	12	14	-	-	-	-	2
	3	4	6	8	10	12	14	-	-	-	-	1
	1	3	5	7	9	11	13	-	-	-	-	0
	1	3	7	1	2	6	2	7	4	11	30	
	Hours			Days			Weeks		Months			

Half-95% confidence intervals (±)
around calculated severity-of-ill-effect scores (above)

162755	2.7	2.1	1.7	1.4	-	-	-	-	-	-	-	12
59874	2.5	1.9	1.5	1.2	1.2	-	-	-	-	-	-	11
22026	2.4	1.8	1.3	1.0	1.1	-	-	-	-	-	-	10
8103	2.3	1.6	1.1	0.8	1.0	-	-	-	-	-	-	9
2981	2.2	1.5	1.0	0.7	0.9	-	-	-	-	-	-	8
1097	2.1	1.5	0.9	0.6	0.9	-	-	-	-	-	-	7
403	2.1	1.5	0.9	0.7	1.0	-	-	-	-	-	-	6
148	2.1	1.5	1.0	0.8	1.1	-	-	-	-	-	-	5
55	2.1	1.5	1.1	1.0	1.3	1.8	-	-	-	-	-	4
20	2.2	1.6	1.3	1.2	1.5	2.0	-	-	-	-	-	3
7	2.2	1.8	1.5	1.5	1.7	2.2	-	-	-	-	-	2
3	2.4	1.9	1.7	1.7	1.9	2.4	-	-	-	-	-	1
1	2.5	2.1	1.9	1.9	2.2	2.6	-	-	-	-	-	0
	1	3	7	1	2	6	2	7	4	11	30	
	Hours			Days			Weeks		Months			

FIGURE 5.—Continued.

the need for additional field work by which to gather physical evidence about the nature and severity of the ill effects. This new capacity to make inferences—an unprecedented development in the field of channel sediment impacts—might also influence the goals of a prosecution.

Impacts on fish populations exposed to episodes of excess sediment may vary according to the cir-

cumstances of the event. For example, fish tend to avoid high concentrations of suspended sediment when possible. Thus, a pollution episode capable of causing high mortality (e.g., of sac fry) or gill damage or starvation or slowed maturation (e.g., of age-0 fingerlings and age-2 juveniles) among caged fish (Reynolds et al. 1989) might not cause any of these direct effects in a wild population that

(log_e mg SS/L)

(log_e mg SS/L)

scores (with

thality), and and physical e more send sediment ulate further a lethal end ill tissue and regulation), orted or re-

nt personnel rent the sed- exposure, and ost probable se equations t the "look- B) simplify on the equa- polation and . They make o distinguish e broad conse- matrixes. sions about




FIGURE 6.—Empirical severity-of-ill-effect scores for adult nonsalmonids (freshwater, group 6) and scores (with half-95% confidence intervals) predicted by model (6). Conventions are those of Figure 1, except the model (B, upper matrix) recognized no threshold of sublethal effects.

is free to move elsewhere in the stream system. Absence of dead fish (notwithstanding reduced egg-to-fry survival) is, however, not necessarily an indication of absence of harm. Indirect effects of sedimentation—loss of summer habitat for feeding and reproduction—may outweigh the direct effects seen in caged fish (Reynolds et al. 1989). This dichotomy has practical implications for enforcement. An investigation during a pollution event should attempt to document suspended sediment concentrations and durations for possible use with the models given here.

However, in the aftermath of a sediment pollution event, the investigation should switch its focus and gather evidence of sediment deposition. Changes in streambed composition resulting from excess sediment are usually manifested as changes in particle size composition. Subjective methods for assessing the extent of sedimentation exist. Objective methods are being developed (Kondolf and Li 1992; Kondolf and Wolman 1993; Potyondy and Hardy 1995) and could be used in place of or in conjunction with the traditional methods. Photographic and videographic records are invaluable regardless of the streambed survey methods chosen.

Four provisions of existing legislation and four potential goals of prosecution are convictions, fines, compensatory damages, and remediation. When the state's purpose is to secure a conviction, a single water sample may be the only evidence required. In some jurisdictions, water quality criteria may be used to identify potential episodes of SS pollution by a tandem system of thresholds. Typically these guidelines state that SS concentrations should not exceed background by more than 10 mg SS/L when background is less than 100 mg SS/L and not more than 10% when background is equal to or greater than 100 mg SS/L (Singleton 1985a, 1985b). This tandem system of thresholds—based on literature reviews specifically intended to document the nature and severity of ill effect under these conditions—is commendable because it recognizes the seasonal patterns in suspended sediment load of natural streams. However, these guidelines do not purport to deal with the inherent nature of sediment as a deleterious substance in aquatic ecosystems as defined by an act of legislation. Nor do they purport to detect the least change in concentration capable of causing ill effects. Various researchers report ill effects when concentrations exceed

background level and Schreiner and Swenson 1991. Prosecution is successful because it is known to have abundances, but pollution (fish f

Adult Freshwater Nonsalmonids

Duration of exposure to SS (\log_e hours)

	0	1	2	3	4	5	6	7	8	9	10
(B)	Average severity-of-ill-effect scores (calculated)										
162755	7	8	9	10	10	11	12	12	13	14	12
59874	7	8	9	9	10	11	11	12	13	14	11
22026	7	8	8	9	10	10	11	12	13	13	10
8103	7	7	8	9	9	10	11	12	12	13	9
2981	6	7	8	8	9	10	11	11	12	13	8
1097	6	7	7	8	9	10	10	11	12	12	7
403	6	6	7	8	9	9	10	11	11	12	6
148	5	6	7	8	8	9	10	10	11	12	5
55	5	6	7	7	8	9	9	10	11	12	4
20	5	6	6	7	8	8	9	10	11	11	3
7	5	5	6	7	7	8	9	10	10	11	2
3	4	5	6	7	7	8	9	9	10	11	1
1	4	5	6	6	7	8	8	9	10	10	0
	1	3	7	1	2	6	2	7	4	11	30
	Hours			Days			Weeks		Months		

Half-95% confidence intervals (\pm)
around calculated severity-of-ill-effect scores (above)

162755	1.6	1.6	1.5	1.5	1.4	1.5	1.5	1.6	1.7	1.8	12
59874	1.5	1.4	1.3	1.3	1.2	1.3	1.3	1.4	1.5	1.6	11
22026	1.4	1.2	1.1	1.1	1.0	1.1	1.1	1.2	1.3	1.4	10
8103	1.3	1.1	1.0	0.9	0.9	0.9	0.9	1.0	1.1	1.2	9
2981	1.2	1.1	0.9	0.8	0.7	0.7	0.8	0.8	1.0	1.1	8
1097	1.2	1.1	0.9	0.8	0.7	0.6	0.7	0.7	0.9	1.0	7
403	1.3	1.1	0.9	0.8	0.7	0.6	0.6	0.7	0.8	1.0	6
148	1.3	1.2	1.0	0.9	0.8	0.7	0.7	0.7	0.8	1.0	5
55	1.5	1.3	1.2	1.0	0.9	0.9	0.8	0.9	0.9	1.0	4
20	1.6	1.5	1.3	1.2	1.1	1.0	1.0	1.0	1.1	1.2	3
7	1.6	1.6	1.5	1.4	1.3	1.2	1.2	1.2	1.2	1.3	2
3	2.0	1.8	1.7	1.6	1.5	1.4	1.4	1.4	1.4	1.5	1
1	2.2	2.0	1.9	1.8	1.7	1.7	1.6	1.6	1.6	1.7	0
	1	3	7	1	2	6	2	7	4	11	30
	Hours			Days			Weeks		Months		

FIGURE 6.—Continued.

background levels by small amounts (see Lawrence and Scherer 1974; Swenson 1978; Gradall and Swenson 1982).

Prosecution based on these rules has been successful because the increased concentrations are known to harm aquatic life. Such evidence abounds, but pertains largely to invertebrate populations (fish food) and primary production (phy-

toplankton and periphyton, the source of energy on which invertebrates may depend) (Newcombe 1994).

However, to the extent that legislation emphasizes the existence of an impact, or the probability of an impact, its primary goal is to secure a conviction. Scope for additional penalty—fines, compensatory damages, and remediation—depends on

d scores (with the model (B,

tion and four convictions. remediation. a conviction, nly evidence r quality critical episodes of thresholds. t SS concen- und by more l is less than b when back- 100 mg SS/L em system of iews specifi- re and sever- ons—is com- seasonal pat- d of natural lo not purport sediment as a cosystems as r do they pur- concentration as researchers tions exceed

Juvenile and Adult Salmonids




FIGURE 7.—Dose-response surfaces describing the severity of ill effect for juvenile and adult salmonids (freshwater, group 1) as a function of suspended sediment concentration and duration of exposure (model 1): $z = 1.0642 + 0.6068(\log_e x) + 0.7384(\log_e y)$.

an ability to demonstrate harmful effects. Dose-response models enhance this capability.

It is difficult to overstate the value of time series water quality data, but there are some kinds of pollution episodes in which other evidence might take precedence. These instances could be classed as catastrophic events in which one or more of the following conditions prevail: (i) the pollution damage is severe, or extensive and highly visible—blanketing by silt, for example; (ii) the extent of harm is to be confirmed by field studies designed and conducted for the purpose (especially relevant for streams on which previous work has been done); or (iii) the pollution event is detected after the fact, in which case the option to sample suspended sediment is foregone already. Notwithstanding these exceptions, efforts to collect sequential water samples during a pollution episode may be the most cost-effective option, especially when court fines, compensation, and remediation are high-priority goals.

In short, the dose-response equations proposed in this report make it possible not only to identify the existence of a pollution event—this information alone being sufficient to secure a conviction—but also to document the severity of ill effect in support of additional penalties.

Meta-analysis

No single researcher could have aspired to conduct all the field work represented in our database. However, the collective works have value beyond anything the original authors could have envisaged. To the extent that this synthesis informs the science, it demonstrates the utility of meta-analysis as a way to shed new light on old problems by using existing data. Limitations of the database can be overcome with further study.

Future Research

The dose-response models in this synthesis are only a beginning. Many gaps remain. Gaps are




FIGURE 8.—Dose-response surfaces describing the severity of ill effect for juvenile and adult salmonids (freshwater, group 1) as a function of suspended sediment concentration (model 2): $z = 0.7565(\log_e y)$.

especially conspicuous (eggs through life stages requiring important thresholds of sediment). Each identified and treated developing unique dose-response p

There are practical distinctions. For example, thresholds must be clearly defined, which raises a problem that must be carefully treated. Susceptibility to sediment must be known. Thresholds of sediment must be known more in particular, thresholds are poorly delineated. Data is a challenge

Adult Salmonids




FIGURE 8.—Dose-response surface describing the severity of ill effect for adult salmonids (freshwater, group 2) as a function of suspended sediment concentration and duration of exposure (model 2): $z = 1.6814 + 0.4769(\log_e x) + 0.7565(\log_e y)$.

Is (fresh-
= 1.0642

roposed
identify
nforma-
tion—
effect in

to con-
atabase.
beyond
: envis-
irms the
ta-anal-
problems
database

esis are
aps are

especially conspicuous for the youngest age-classes (eggs through young juveniles). The pooling of life stages required for these models—eggs with larvae, young with old juveniles—doubtless masks important thresholds of susceptibility to suspended sediment. Each developmental stage should be identified and treated separately for the purpose of developing uniquely age-specific and size-specific dose-response profiles.

There are practical reasons to make such distinctions. For example, artificial spawning channels must be cleaned annually. Gravel cleaning, which raises a plume of silty water, therefore must be carefully timed to minimize the potential ill effects. Susceptibilities of resident life stages to sediment must be known.

Thresholds of sublethal and lethal effects must be known more precisely. Our analysis has shown, in particular, that sublethal effects thresholds are poorly delineated for most groups. Finding useable data is a challenge; we rejected many studies be-

cause they were too vague about sediment concentration, duration of exposure, or the exact nature of the ill effect. We undoubtedly overlooked some reports, but more directed research is warranted. Research is especially needed into particle quality (particle size, angularity, and mineralogy), particle toxicity (toxicants in and adsorbed on sediments), and temperature effects.

Particle quality and toxicology.—Ill effects increase as a function of increasing particle size (if other variables are kept constant). Pollution events often subject fish to particle sizes to which they are not normally exposed. Newcombe et al. (1995) documented that rainbow trout died rapidly when exposed to a silty water discharge (mortality, 80–100%; concentration, $\approx 4,315$ mg SS/L; duration, <57 h; particle sizes, 100–170 μm , water temperature, 10°C). These results differ from those from other pollution episodes in which the particle size was smaller; generally, the ill effects would be much less severe—on the order of 0–10% mor-

Juvenile Salmonids




FIGURE 9.—Dose-response surface describing the severity of ill effect for juvenile salmonids (freshwater, group 3) as a function of suspended sediment concentration and duration of exposure (model 3): $z = 0.7262 + 0.7034(\log_e x) + 0.7144(\log_e y)$.

tality. Some research to quantify ill effect as a function of particle size has been done with several species of Pacific salmon (Servizi and Martens 1987, 1991, 1992). Further work should make it possible to create a set of dose-response models as functions of particle size range that are unique to each relevant life stage. The growing need to explore ill effects of suspended sediment as a function of particle size imposes an obligation among fisheries biologists to use a uniform nomenclature in reference to the particle grade scale. Suitable systems exist already so there is no need to invent a more specialized one. For example, soils scientists recognize three particle size-classes—sand, silt and clay (Agriculture Canada 1974)—with formalized subdivisions, names, and sizes as follows: very coarse sand, 2.0–1.0 mm; coarse sand, 1.0–0.5 mm; medium sand, 0.5–0.25 mm; fine sand, 0.25–0.10 mm; very fine sand, 0.10–0.05 mm; silt, 0.05–0.002 mm; and clay, ≤ 0.002 mm. Fisheries

biologists would do well to adopt this or some similar particle grade scale.

The importance of particle angularity, especially in relation to gill abrasion, should be studied. The mineralogy of sediment particles may offer clues to the potential for toxicity and physiological effects. Likewise, the presence of innate or adsorbed toxicants may offer clues to latent effects on fish population health. Studies of the mineralogy and potential chemical activity of the particle itself, of particles in the colloidal size range capable of entering the fish's cells, and of particles with adsorbed toxicants may reveal common properties relating to fate and ill effect at the tissue and cellular level. If common properties do exist among these particular variables, there may be a unifying explanation in the phenomenon of phagocytosis.

Phagocytosis, the envelopment of fine particles by cells of the fish's gill and gut, transports the particles into the fish's body. Although these par-




FIGURE 10.—Dose-response surface describing the severity of ill effect for non-salmonids (freshwater, group 3) as a function of suspended sediment concentration and duration of exposure (model 3): $z = 0.7262 + 0.7034(\log_e x) + 0.7144(\log_e y)$.

ticles may end up in the fish's body, they may also be a major reposit of sediment. Fishes exposed to fine sediment to the extent that they are able to ingest it (e.g., the glass mussel, *Mytilus edulis*, Malaspina, 1987). Thus, particles could be transported into the cells of the fish, where they could be harmful. This is especially among fish that are only partially exposed to sediment. Particles may be particularly harmful when they are present in water runoff or in sediment. Such latent ill effects are a common phenomenon.

Water temperature is an important function of ambient water quality and should be explored more fully.

Eggs and Larvae of Salmonids and Nonsalmonids




FIGURE 10.—Dose-response surface describing the severity of ill effect for eggs and larvae of salmonids and nonsalmonids (freshwater and estuarine, group 4) as a function of suspended sediment concentration and duration of exposure (model 4): $z = 3.7466 + 1.0946(\log_e x) + 0.3117(\log_e y)$.

water, group
7034($\log_e x$)

s or some

especially
died. The
offer clues
logical ef-
r adsorbed
cts on fish
alogy and
e itself, of
ible of en-
with ad-
properties
e and cel-
ist among
a unifying
ocytosis.

particles
sports the
these par-

ticles may end up in various tissues, the spleen is a major repository. The spleens of some fishes exposed to fine sediment become mineralized to the extent that the tissue damages the cutting edge of the glass microtome blades (Goldes 1983; S. Goldes, Malaspina College, personal communication). Thus, phagocytosis of fine suspended sediments could trigger a sequence of harmful events within the cells of a fish's body leading to ill effects that are only partially understood today. Invasive particles may be the biological equivalent of a Trojan horse: harmless when on the outside, devastating when on the inside. Tumorigenesis, especially among groundfish that dwell in harbors where sediments may be contaminated by storm-water runoff or by industrial effluent, may be one such latent ill effect yet to be linked to this phenomenon.

Water temperature.—Severity of ill effect as a function of ambient water temperature ought to be explored more fully. Ill effects are greater in sea-

sonably warm water than would be the case for the same fishes in seasonably cold water. Mechanisms for this effect have not been systematically described. The dynamics of this variable probably have to do with the temperature-related patterns of oxygen saturation, respiration rate, and metabolic rate of fishes (slower in cool water, more rapid in warm)—all of which result in reduced risk of gill abrasion in cool water and increased risk in warm water. These mechanisms should be explored in the context of seasonal temperature ranges in a fish's natural habitat.

Ecosystem Considerations

Broad-based ecosystem research supporting stream protection is under way, but it is a relatively new science. Stream protection requires, among other things, quantitative linkages between impacts of channel sediment and the land use practices that generate the sediment. Leadership in this area will come from many disciplines, as exem-

Adult Freshwater Nonsalmonids




FIGURE 12.—Dose-response surface describing the severity of ill effect for adult nonsalmonids (freshwater, group 6) as a function of suspended sediment concentration and duration of exposure (model 6): $z = 4.0815 + 0.7126(\log_e x) + 0.2829(\log_e y)$.

arine, group
9647($\log_e x$)

protection
away from
major factor
limit is a
Unless it is
can not be
increased
might be pub-
protection
stream pro-

(Nanaimo,
rest in this
estions. We
uska Coop-
iversity of
(Fisheries
Lands and

Parks), and Jerry Sweeten (Asherwood Learning Center) for raw data; Sally Goldes (Fisheries Branch, Malaspina College, Nanaimo) for information about fate and effects of small particles on cells and tissues of fish; Mike Miles (Mike Miles and Associates, Victoria, British Columbia), Howard Singleton (Water Quality Branch, Ministry of Environment, Lands and Parks), and Mark Labelle (Institut Français de Recherche pour l'Exploitation de la Mer, Nantes Cedex) for various suggestions; Bill McLean (Qunisam River Hatchery, Campbell River, British Columbia) for field-testing some of the models; and American Fisheries Society reviewers and staff for their numerous improvements to the manuscript.

References

- Agriculture Canada, 1974. The system of soil classification for Canada. Canada Department of Agriculture, Publication 1455, Information Canada, Ottawa.
- Alabaster, J. S., and R. Lloyd. 1980. Finely divided solids. Pages 1-20 in Water quality criteria for freshwater fish. Butterworth, London.
- Auld, A. H., and J. R. Schubel. 1978. Effects of suspended sediment on fish eggs and larvae: a laboratory assessment. *Estuarine and Coastal Marine Science* 6:153-164.
- Berg, L. 1983. Effects of short term exposure to suspended sediments on the behaviour of juvenile coho salmon. Master's thesis. University of British Columbia, Vancouver.
- Berg, L., and T. G. Northcote. 1985. Changes in territorial, gill-flaring, and feeding behaviour in juvenile coho salmon (*Oncorhynchus kisutch*) following short-term pulses of suspended sediment. *Canadian Journal of Fisheries and Aquatic Sciences* 42:1410-1477.
- Berkmann, H. E., and C. F. Rabeni. 1987. Effect of siltation on stream fish communities. *Environmental Biology of Fishes* 18:285-294.
- Birtwell, I. K., G. F. Hartman, B. Anderson, D. J. McLeay, and J. G. Malick. 1984. A brief investigation of Arctic grayling (*Thymallus arcticus*) and aquatic invertebrates in the Minto Creek drainage,

- of fine sediment rates below a watal Management
- inorganic sediment nental Protection Research Series. D.C.
- idity on feeding ransactions of the 446-450.
- nd operation of a y. His Majesty's
- ultrastructural efe gills of rainbow). Master's thesis. itario.
1982. Responses turbidity. Trans- Society 111:392-
- .. Jayasuriya, and ility in mountain ts of roads from ial of Hydrology
7. Martens, 1995. ex for predicting i American Jour- 68-873.
- the tolerance of ided sediment in ogy and Mineral B):28-31. (Not i 1980.)
- nd-pit washings- Vereinigung für nologie Verhan-
- 15 years of fish- esearch to man- Chamberlin, ed- n Creek results. n Creek Steering mbia.
- C. Dart, and R. -clay wastes on of Air and Water
961. The effect urvival of trout. ater Pollution 5:
- ds. 1963. The -suspensions of onal Journal of
- rd. 1962. The urvival of rain- ent 8:608-609. ,loyd 1980.)
- Hesse, L. W., and B. A. Newcomb. 1982. Effects of flushing Spencer Hydro on water quality, fish, and insect fauna in the Niobrara River, Nebraska. *North American Journal of Fisheries Management* 2:45-52.
- Horkel, J. D., and W. D. Pearson. 1976. Effects of turbidity on ventilation rates and oxygen consumption of green sunfish, *Lepomis cyanellus*. *Transactions of the American Fisheries Society* 105:107-113.
- Hughes, G. M. 1975. Coughing in the rainbow trout (*Salmo gairdneri*) and the influence of pollutants. *Revue Suisse de Zoologie* 82:47-64.
- Johnson, D. D., and D. J. Wildish. 1982. Effect of suspended sediment on feeding by larval herring (*Clupea harengus harengus* L.). *Bulletin of Environmental Contamination and Toxicology* 29:261-267.
- Kemp, H. A. 1949. Soil pollution in the Potomac River basin. *American Water Works Association Journal* 41:792-796. (Not seen: cited by Cordone and Kelley 1961.)
- Kondlof, G. M., and S. Li. 1992. The pebble count techniques for quantifying surface bed material size in instream flow studies. *Rivers* 3:80-87.
- Kondolf, G. M., and M. G. Wolman. 1993. The sizes of salmonid spawning gravels. *Water Resources Research* 29:2275-2285.
- Langer, O. E. 1980. Effects of sedimentation on salmonid stream life. Environment Canada, Environmental Protection Service, unpublished report, North Vancouver, British Columbia.
- Lawrence, M., and E. Scherer. 1974. Behavioral responses of whitefish and rainbow trout to drilling fluids. *Canada Fisheries and Marine Service Technical Report* 502.
- LeGore, R. S., and D. M. DesVoigne. 1973. Absence of acute effects on three-spine sticklebacks (*Gasterosteus aculeatus*) and coho salmon (*Oncorhynchus kisutch*) exposed to re-suspended harbour sediment contaminants. *Journal of the Fisheries Research Board of Canada* 30:1240-1242.
- Lloyd, D. S. 1985. Turbidity in freshwater habitats of Alaska: a review of published and unpublished literature relevant to the use of turbidity as a water quality standard. Alaska Department of Fish and Game, Habitat Division, Report 85, Part 1, Juneau.
- MacDonald, D. D., and C. P. Newcombe. 1993. Utility of the stress index for predicting suspended sediment effects: response to comment. *North American Journal of Fisheries Management* 13:873-876.
- Macdonald, J. S., editor. 1994. Proceedings of the Takla fishery/forestry workshop: a two year review. *Canadian Technical Report of Fisheries and Aquatic Sciences* 2007.
- Macdonald, J. S., J. C. Scrivener, and G. Smith. 1992. The Stuart-Takla fisheries/forestry interaction project: study description and design. *Canadian Technical Report of Fisheries and Aquatic Sciences* 1899.
- MacKinlay, D. D., D. D. MacDonald, M. K. Johnson, and R. F. Fielden. 1987. Culture of chinook salmon (*Oncorhynchus tshawytscha*) in iron-rich groundwater: Stuart pilot hatchery experiences. *Canadian Manuscript Report of Fisheries and Aquatic Sciences* 1944.
- McLeay, D. J., I. K. Birtwell, G. F. Hartman, and G. L. Ennis. 1987. Responses of Arctic grayling (*Thymallus arcticus*) to acute and prolonged exposure to Yukon placer mining sediment. *Canadian Journal of Fisheries and Aquatic Sciences* 44:658-673.
- McLeay, D. J., G. L. Ennis, I. K. Birtwell, and G. F. Hartman. 1984. Effects on Arctic grayling (*Thymallus arcticus*) of prolonged exposure to Yukon placer mining sediment: laboratory study. *Canadian Technical Report of Fisheries and Aquatic Sciences* 1241.
- McLeay, D. J. and five coauthors. 1983. Effects on Arctic grayling (*Thymallus arcticus*) of short term exposure to Yukon placer mining sediments: laboratory and field studies. *Canadian Technical Report of Fisheries and Aquatic Sciences* 1171.
- Menzel, B. W., J. B. Barnum, and L. M. Antosch. 1984. Ecological alterations of Iowa prairie-agricultural streams. *Iowa State Journal of Research* 59:5-30.
- Morgan, R. P. II, J. V. Rasin, Jr., and L. A. Noe. 1973. Effects of suspended sediments on the development of eggs and larvae of striped bass and white perch. appendix 11. Final Report to U.S. Army Corps of Engineers, Contract DACW61-71-C0062, Philadelphia.
- Morgan, R. P. II, J. R. Rasin, Jr., and L. A. Noe. 1983. Sediment effects on eggs and larvae of striped bass and white perch. *Transactions of the American Fisheries Society* 112:220-224.
- Neumann, D. A., J. M. O'Connor, J. A. Sherk, and K. V. Wood. 1975. Respiratory and hematological responses of oyster toadfish (*Opsanus tau*) to suspended solids. *Transactions of the American Fisheries Association* 104:775-781.
- Newcomb, T. W., and T. A. Flagg. 1983. Some effects of Mt. St. Helens ash on juvenile salmon smolts. *U.S. National Marine Fisheries Service Marine Fisheries Review* 45(2):8-12.
- Newcombe, C. P. 1986. Fisheries and the problem of turbidity and inert sediment in water: a synthesis for environmental impact assessment. British Columbia Ministry of Environment, Environmental Impact Unit, Environmental Services Section, Waste Management Branch, Victoria.
- Newcombe, C. P. 1994. Suspended sediment in aquatic ecosystems: ill effects as a function of concentration and duration of exposure. *British Columbia Ministry of Environment, Lands and Parks, Habitat Protection Branch, Victoria.*
- Newcombe, C. P., and D. D. MacDonald. 1991. Effects of suspended sediments on aquatic ecosystems. *North American Journal of Fisheries Management* 11:72-82.
- Newcombe, C. P., B. Shepherd, G. Hoyer, and M. Ladd. 1995. Documentation of a fish kill (juvenile rainbow trout: *Oncorhynchus mykiss*) in Bellevue Creek (near Mission, Kelowna, British Columbia, Canada), caused by silty water discharge. *British Columbia Ministry of Environment, Lands and Parks,*

- Habitat Protection Branch, Habitat Protection Occasional Report. Victoria.
- Noggle, C. C. 1978. Behavioral, physiological and lethal effects of suspended sediment on juvenile salmonids. Master's thesis. University of Washington, Seattle.
- Ott, A. G. 1984. Personal communication. Alaska Department of Fish and Game, Fairbanks. (Not seen: cited as personal communication in Lloyd 1985.)
- Peters, J. C. 1967. Effects on a trout stream of sediment from agricultural practices. *Journal of Wildlife Management* 31:805-812.
- Phillips, R. W. 1970. Effects of sediment on the gravel environment and fish production. Pages 64-74 in *Proceedings of the symposium on forest land use and stream environment*. Oregon State University, Continuing Education Publications, Corvallis, Oregon.
- Potyondy, J. P. and T. Hardy. 1995. Reply to discussion by G. Mathias Kondolf. "Use of pebble counts to evaluate fine sediment increase in stream channels." *Water Resources Bulletin* 31:539-540.
- Ptolemy, R. A. 1993. Maximum salmonid densities in fluvial habitats in British Columbia. Pages 223-250 in L. Berg and P. W. Delaney, editors. *Proceedings of the Coho Workshop*. Nanaimo, British Columbia. Department of Fisheries and Oceans, Vancouver.
- Redding, J. M., and C. B. Schreck. 1982. Mount St. Helens ash causes sublethal stress responses in steelhead trout. Pages 300-307 in *Mt. St. Helens: effects on water resources*. Washington State University, Washington Water Research Center, Report 41, Pullman.
- Reynolds, J. B., R. C. Simmons, and A. R. Burkholder. 1989. Effects of placer mining discharge on health and food of Arctic grayling. *Water Resources Bulletin* 25:625-635.
- Rogers, B. A. 1969. Tolerance levels of four species of estuarine fishes to suspended mineral solids. Master's thesis. University of Rhode Island, Kingston.
- Rosenberg, D. M., and N. B. Snow. 1977. A design for environmental impact studies with special reference to sedimentation in aquatic systems of the Mackenzie and Porcupine river drainages. Pages 65-78 in *Proceedings of the circumpolar conference on northern ecology*. National Research Council, Ottawa.
- Scannell, P. A. 1988. Effects of elevated sediments levels from placer mining on survival and behavior of immature Arctic grayling. Master's thesis. University of Alaska, Fairbanks.
- Schubel, J. R., and J. C. S. Wang. 1973. The effects of suspended sediment on the hatching success of *Percia flavescens* (yellow perch), *Morone americana* (white perch), *Morone saxatilis* (striped bass) and *Alosa pseudoharengus* (alewife) eggs. Chesapeake Bay Institute, Johns Hopkins University Special Report 30, Reference 73-3, Baltimore, Maryland. (Not seen: cited by Morgan et al. 1983.)
- Scrivener, C. J., T. G. Brown, and B. C. Anderson. 1993. Juvenile chinook salmon (*Oncorhynchus tshawytscha*) utilization of Hawks Creek, a small nonnatal tributary of the upper Fraser River. *Canadian Journal of Fisheries and Aquatic Sciences* 51:1139-1146.
- Scullion, J., and R. W. Edwards. 1980. The effects of pollutants from the coal industry on fish fauna of a small river in the South Wales coalfield. *Environmental Pollution Series A* 21:141-153.
- Servizi, J. A., and D. W. Martens. 1987. Some effects of suspended Fraser River sediments on sockeye salmon (*Oncorhynchus nerka*). *Canadian Special Publication of Fisheries and Aquatic Sciences* 96:254-264.
- Servizi, J. A., and D. W. Martens. 1991. Effect of temperature, season, and fish size on acute lethality of suspended sediments to coho salmon (*Oncorhynchus kisutch*). *Canadian Journal of Fisheries and Aquatic Sciences* 48:493-497.
- Servizi, J. A., and D. W. Martens. 1992. Sublethal responses of coho salmon (*Oncorhynchus kisutch*) to suspended sediments. *Canadian Journal of Fisheries and Aquatic Sciences* 49:1389-1395.
- Shaw, P. A., and J. A. Maga. 1943. The effect of mining silt on yield of fry from salmon spawning beds. *California Fish and Game* 29:29-41.
- Sherk, J. A., J. M. O'Connor, and D. A. Neumann. 1975. Effects of suspended and deposited sediments on estuarine environments. Pages 541-558 in L. E. Cronin, editor. *Estuarine Research* 2. Academic Press, New York.
- Sigler, J. W., T. C. Bjornn, and F. H. Everest. 1984. Effects of chronic turbidity on density and growth of steelheads and coho salmon. *Transactions of the American Fisheries Society* 113:142-150.
- Simmons, R. C. 1982. Effects of placer mining on Arctic grayling of interior Alaska. Master's thesis. University of Alaska, Fairbanks.
- Singleton, H. J. 1985a. Water quality criteria for particulate matter. British Columbia Ministry of Environment, Water Management Branch, Victoria.
- Singleton, H. J. 1985b. Water quality criteria for particulate matter: technical appendix. British Columbia Ministry of Environment, Water Management Branch, Victoria.
- Slaney, P. A., T. G. Halsey, and H. A. Smith. 1977a. Some effects of forest harvesting on salmonid rearing habitat in two streams in the central interior of British Columbia. British Columbia Ministry of Recreation and Conservation, Fish and Wildlife Branch, Fisheries Management Report 71, Victoria.
- Slaney, P. A., T. G. Halsey, and A. F. Tautz. 1977b. Effects of forest harvesting practices on spawning habitat of stream salmonids in the Centennial Creek watershed, British Columbia. British Columbia Ministry of Recreation and Conservation, Fish and Wildlife Branch, Fisheries Management Report 73, Victoria.
- Slanina, K. 1962. Beitrag zur Wirkung mineralischer Suspensionen auf Fische. *Wasser und Abwasser* 1962:186-194.
- Smith, O. R. 1940. Placer mining silt and its relation to the salmon and trout on the Pacific coast. *Transactions of the American Fisheries Society* 69:230.
- Stober, Q. J., and five coauthors. 1984. Effects of suspended volcanic sediments on salmon in the Toutle River, Washington. *Fisheries Bulletin* 82:1-10.
- Suchanek, P. M., R. P. M. Schmidt, and R. P. M. Schmidt. 1984a. Juvenile salmonid survival criteria. Alaska Department of Fish and Game, Division of Hydro and Aquatic Sciences, Anchorage. (Not seen: cited by Lloyd 1985.)
- Suchanek, P. M., R. L. Schmidt, and R. P. M. Schmidt. 1984b. Resident fish populations in the Tustumene River. Alaska Department of Fish and Game, Division of Hydro and Aquatic Sciences, Anchorage. (Not seen: cited by Lloyd 1985.)
- Swenson, W. A. 1978. Turbidity and abundance in western streams. *Environmental Protection Agency Research Center Report EPA 600/3-78-067*. (Not seen: cited by Swenson 1982.)
- Swenson, W. A., and M. L. Swenson. 1982. Turbidity on survival of larval lake herring in streams of the American West. *Journal of Great Lakes Research* 8:545-555.
- Sykora, J. L., E. J. Smith, and J. L. Smith. 1982. Effects of lime-neutralized placer tailings on juvenile brook trout in a stream. *Water Research* 16:93-100.
- Townsend, A. H. 1983. Effects of placer mining on the Chatanika River, Alaska. Master's thesis. Alaska Department of Fish and Game, Division of Hydro and Aquatic Sciences, Anchorage. (Not seen: cited by Lloyd 1985.)

- Canadian Journal of Fisheries and Aquatic Sciences 51:1139-1144.
- The effects of suspended sediment on the benthic fauna of a stream. Environmental Management 10:101-106.
- Some effects of suspended sediment on sockeye salmon. Canadian Journal of Fisheries and Aquatic Sciences 96:101-106.
- Effect of temperature on the lethality of suspended sediment to rainbow trout (*Oncorhynchus mykiss*). Journal of the American Fisheries Society 114:101-106.
- Sublethal effects of suspended sediment on rainbow trout (*Oncorhynchus mykiss*). Journal of the American Fisheries Society 114:101-106.
- Effect of mining on the environment. Environmental Management 10:101-106.
- Wagener, G. L., and J. D. LaPerriere. 1985. Effects of placer mining on the invertebrate communities of interior Alaska streams. *Freshwater Invertebrate Biology* 4:208-214.
- Wallen, E. I. 1951. The direct effect of turbidity on fishes. *Oklahoma Agricultural and Mechanical College, Arts and Sciences Studies, Biological Series* 48(2). (Not seen: cited by Alabaster and Lloyd 1980.)
- Whitman, R. P., T. P. Quinn, and E. L. Brannon. 1982. Influence of suspended volcanic ash on homing behavior of adult chinook salmon. *Transactions of the American Fisheries Society* 111:63-69.
- Turnpenny, A. W. H., and R. Williams. 1980. Effects of sedimentation on the gravels of an industrial river system. *Journal of Fish Biology* 17:681-693.
- Vaughan, G. L. 1979. Effects of stripmining on fish and diatoms in streams of the New River drainage basin. *Journal of the Tennessee Academy of Science* 54:110-114.
- Vaughan, G. L., L. Minter, and J. Schiller. 1982. New River project data bases and documentation. Joint research, volume 2: biological and associated water quality data. University of Tennessee, Departments of Civil Engineering and Zoology, and U.S. Department of Energy, Knoxville, Tennessee.
- Vaughan, G. L., A. Talak, and R. J. Anderson. 1978. The chronology and character of recovery of aquatic communities from the effects of strip mining for coal in east Tennessee. *U.S. Fish and Wildlife Service FWS/OBS-78/81:119-125*.
- Vinyard, G. L., and W. J. O'Brien. 1976. Effects of light and turbidity on the reactive distance of bluegill (*Lepomis macrochirus*). *Journal of the Fisheries Research Board of Canada* 33:2845-2849.
- Wagener, G. L., and J. D. LaPerriere. 1985. Effects of placer mining on the invertebrate communities of interior Alaska streams. *Freshwater Invertebrate Biology* 4:208-214.
- Wallen, E. I. 1951. The direct effect of turbidity on fishes. *Oklahoma Agricultural and Mechanical College, Arts and Sciences Studies, Biological Series* 48(2). (Not seen: cited by Alabaster and Lloyd 1980.)
- Whitman, R. P., T. P. Quinn, and E. L. Brannon. 1982. Influence of suspended volcanic ash on homing behavior of adult chinook salmon. *Transactions of the American Fisheries Society* 111:63-69.
- actions of the American Fisheries Society 69:225-230.
- Stober, Q. J., and five coauthors. 1981. Effects of suspended volcanic sediment on coho and chinook salmon in the Toutle and Cowlitz rivers. University of Washington, Fisheries Research Institute, Technical Completion Report FRI-UW-8124, Seattle.
- Suchanek, P. M., R. P. Marshall, S. S. Hale, and D. C. Schmidt. 1984a. Juvenile salmon rearing suitability criteria. Alaska Department of Fish and Game, Susitna Hydro Aquatic Studies, 1984 Report 2, Part 3. Anchorage. (Not seen: cited by Lloyd 1985.)
- Suchanek, P. M., R. L. Sundet, and M. N. Wenger. 1984b. Resident fish habitat studies. Alaska Department of Fish and Game, Susitna Hydro Aquatic Studies, 1984 Report 2, Part 6. Anchorage. (Not seen: cited by Lloyd 1985.)
- Swenson, W. A. 1978. Influence of turbidity on fish abundance in western Lake Superior. U. S. Environmental Protection Agency, National Environmental Research Center, Ecological Research Series EPA 600/3-78-067. (Not seen: cited by Gradall and Swenson 1982.)
- Swenson, W. A., and M. L. Matson. 1976. Influence of turbidity on survival, growth, and distribution of larval lake herring (*Coregonus artedii*). *Transactions of the American Fisheries Society* 105:541-545.
- Sykora, J. L., E. J. Smith, and M. Synak. 1972. Effect of lime-neutralized iron hydroxide suspensions on juvenile brook trout (*Salvelinus fontinalis* Mitchell). *Water Research* 6:935-950.
- Townsend, A. H. 1983. Sport fishing—placer mining: Chatanika River. Memorandum to Director B. Baker, Habitat Division, Alaska Department of Fish and Game, February 2, 1983, Juneau. (Not seen: cited by Lloyd 1985.)
- Wagener, G. L., and J. D. LaPerriere. 1985. Effects of placer mining on the invertebrate communities of interior Alaska streams. *Freshwater Invertebrate Biology* 4:208-214.
- Wallen, E. I. 1951. The direct effect of turbidity on fishes. *Oklahoma Agricultural and Mechanical College, Arts and Sciences Studies, Biological Series* 48(2). (Not seen: cited by Alabaster and Lloyd 1980.)
- Whitman, R. P., T. P. Quinn, and E. L. Brannon. 1982. Influence of suspended volcanic ash on homing behavior of adult chinook salmon. *Transactions of the American Fisheries Society* 111:63-69.

Appendix follows on page 720

Appendix: Dose-Response Database

TABLE A.1.—Continued

TABLE A.1 Dose-response database for fishes exposed to suspended sediment.

Species	Life stage ^a	Sediment dose		Fish response		Reference	Species
		Exposure concentration (mg/L)	Exposure duration (h)	SEV ^b	Description ^c		
Adult salmonids and rainbow smelt (freshwater, groups 1 and 2)							
Grayling (Arctic)	A	100	0.10	3	Fish avoided turbid water	Suchanek et al. (1984a, 1984b)	Trout (brown)
Grayling (Arctic)	A	100	1,008	8	Fish had decreased resistance to environmental stresses	McLeay et al. (1984)	Trout (brown)
Grayling (Arctic)	A	100	1,008	9	Impaired feeding	McLeay et al. (1984)	Trout (brown)
Grayling (Arctic)	A	100	1,008	9	Reduced growth	McLeay et al. (1984)	Trout (brown)
Salmon	A	25	4	4	Feeding activity reduced	Phillips (1970)	Trout (brown)
Salmon	A	16.5	24	4	Feeding behavior apparently reduced	Townsend (1983); Ott (1984)	Trout (cutthroat)
Salmon	A	1,650	240	7	Loss of habitat caused by excessive sediment transport	Coats et al. (1985)	Trout (lake)
Salmon	A	75	168	7	Reduced quality of rearing habitat	Slaney et al. (1977b)	Trout (rainbow)
Salmon	A	210	24	10	Fish abandoned their traditional spawning habitat	Hamilton (1961)	Trout (rainbow)
Salmon (Atlantic)	A	2,500	24	10	Increased risk of predation	Gibson (1933)	Trout (rainbow)
Salmon (chinook)	A	650	168	5	No histological signs of damage to olfactory epithelium	Brannon et al. (1981)	Trout (rainbow)
Salmon (chinook)	A	350	0.17	7	Home water preference disrupted	Whitman et al. (1982)	Trout (rainbow)
Salmon (chinook)	A	650	168	7	Homing behavior normal, but fewer test fish returned	Whitman et al. (1982)	Trout (rainbow)
Salmon (chinook)	A	39,300	24	10	No mortality (VA, <5-100 µm; median, <15 µm)	Newcomb and Flagg (1983)	Trout (rainbow)
Salmon (chinook)	A	82,400	6	12	Mortality rate 60% (VA, <5-100 µm)	Newcomb and Flagg (1983)	Trout (rainbow)
Salmon (chinook)	A	207,000	1	14	Mortality rate 100% (VA, <5-100 µm)	Newcomb and Flagg (1983)	Trout (rainbow)
Salmon (Pacific)	A	525	588	10	No mortality (other end points not investigated)	Griffin (1938)	Trout (rainbow)
Salmon (sockeye)	A	500	96	8	Plasma glucose levels increased 39%	Servizi and Martens (1987)	Trout (rainbow)
Salmon (sockeye)	A	1,500	96	8	Plasma glucose levels increased 150%	Servizi and Martens (1987)	Trout (rainbow)
Salmon (sockeye)	A	39,300	24	10	No mortality (VA, <5-100 µm; median, <15 µm)	Newcomb and Flagg (1983)	Trout (rainbow)
Salmon (sockeye)	A	82,400	6	12	Mortality rate 60% (VA, <5-100 µm; median, <15 µm)	Newcomb and Flagg (1983)	Trout (rainbow)
Salmon (sockeye)	A	207,000	1	14	Mortality rate 100% (VA)	Newcomb and Flagg (1983)	Trout (sea)
Smelt (rainbow)	A	3.5	168	7	Increased vulnerability to predation	Swenson (1978)	Whitefish (lake)
Steelhead	A	500	3	5	Signs of sublethal stress (VA)	Redding and Schreck (1982)	Whitefish (lake)
Steelhead	A	1,650	240	7	Loss of habit caused by excessive sediment transport	Coats et al. (1985)	Whitefish (mountain)
Steelhead	A	500	9	8	Blood cell count and blood chemistry change	Redding and Schreck (1982)	Grayling (Arctic)
Trout	A	16.5	24	4	Feeding behavior apparently reduced	Townsend (1983); Ott (1984)	Grayling (Arctic)
Trout	A	75	168	7	Reduced quality of rearing habitat	Slaney et al. (1977b)	Grayling (Arctic)
Trout	A	270	312	8	Gill tissue damaged	Herbert and Merckens (1961)	Grayling (Arctic)
Trout	A	525	588	10	No mortality (other end points not investigated)	Griffin (1938)	Grayling (Arctic)
Trout	A	300	720	12	Decrease in population size	Peters (1967)	Grayling (Arctic)
Trout (brook)	A	4.5	168	3	Fish more active and less dependent on cover	Gradall and Swenson (1982)	Grayling (Arctic)

TABLE A.1.—Continued.

Reference	Species	Sediment dose			Fish response		Reference
		Life stage ^a	Exposure concentration (mg/L)	Exposure duration (h)	SEV ^b	Description ^c	
	Trout (brown)	A	1,040	17,520	8	Gill lamellae thickened (VFSS)	Herbert et al. (1961)
	Trout (brown)	A	1,210	17,520	8	Some gill lamellae became fused (VFSS)	Herbert et al. (1961)
al. (1984a, 1984b)	Trout (brown)	A	18	720	10	Abundance reduced	Peters (1967)
al. (1984)	Trout (brown)	A	100	720	11	Population reduced	Scullion and Edwards (1980)
	Trout (brown)	A	1,040	8,760	14	Population one-seventh of expected size (River Fal)	Herbert et al. (1961)
al. (1984)	Trout (brown)	A	5,838	8,760	14	Fish numbers one-seventh of expected (River Par)	Herbert et al. (1961)
al. (1984)	Trout (cutthroat)	A	35	2	4	Feeding ceased; fish sought cover	Cordone and Kelly (1961)
1983); Ott (1984)	Trout (lake)	A	3.5	168	3	Fish avoided turbid areas	Swenson (1978)
(1985)	Trout (rainbow)	A	66	1	3	Avoidance behavior manifested part of the time	Lawrence and Scherer (1974)
	Trout (rainbow)	A	665	1	3	Fish attracted to turbidity	Lawrence and Scherer (1974)
(1977b)	Trout (rainbow)	A	100	0.10	3	Fish avoided turbid water (avoidance behavior)	Suchanek et al. (1984a, 1984b)
961)	Trout (rainbow)	A	100	0.25	5	Rate of coughing increased (FSS)	Hughes (1975)
	Trout (rainbow)	A	250	0.25	5	Rate of coughing increased (FSS)	Hughes (1975)
33)	Trout (rainbow)	A	810	504	8	Gills of fish that survived had thickened epithelium	Herbert and Merkens (1961)
al. (1981)	Trout (rainbow)	A	17,500	168	8	Fish survived; gill epithelium proliferated and thickened	Slanina (1962)
al. (1982)	Trout (rainbow)	A	50	960	9	Rate of weight gain reduced (CWS)	Herbert and Richards (1963)
al. (1982)	Trout (rainbow)	A	50	960	9	Rate of weight gain reduced (WF)	Herbert and Richards (1963)
	Trout (rainbow)	A	810	504	10	Some fish died	Herbert and Merkens (1961)
and Flagg (1983)	Trout (rainbow)	A	270	3,240	10	Survival rate reduced	Herbert and Merkens (1961)
	Trout (rainbow)	A	200	24	10	Test fish began to die on the first day (WF)	Herbert and Richards (1963)
and Flagg (1983)	Trout (rainbow)	A	80,000	24	10	No mortality	D. Herbert, personal communication to Alabaster and Lloyd (1980)
and Flagg (1983)	Trout (rainbow)	A	18	720	10	Abundance reduced	Peters (1967)
38)	Trout (rainbow)	A	59	2,250	10	Habitat damage: reduced porosity of gravel	Slaney et al. (1977b)
Martens (1987)	Trout (rainbow)	A	4,250	588	12	Mortality rate 50% (CS)	Herbert and Wakeford (1962)
Martens (1987)	Trout (rainbow)	A	49,838	96	12	Mortality rate 50% (DM)	Lawrence and Scherer (1974)
	Trout (rainbow)	A	3,500	1,488	13	Catastrophic reduction in population size	Herbert and Merkens (1961)
and Flagg (1983)	Trout (rainbow)	A	160,000	24	14	Mortality rate 100%	D. Herbert, personal communication to Alabaster and Lloyd (1980)
and Flagg (1983)	Trout (sea)	A	210	24	10	Fish abandoned traditional spawning habitat	Hamilton (1961)
and Flagg (1983)	Whitefish (lake)	A	0.66	1	3	Swimming behavior changed	Lawrence and Scherer (1974)
1978)	Whitefish (lake)	A	16,613	96	12	Mortality rate 50% (DM)	Lawrence and Scherer (1974)
and Schreck (1982)	Whitefish (mountain)	A	10,000	24	10	Fish died: silt-clogged gills	Langer (1980)
(1985)							
Juvenile salmonids (freshwater, groups 1 and 3)							
and Schreck (1982)	Grayling (Arctic)	U	20	24	3	Fish avoided parts of the stream	Birtwell et al. (1984)
	Grayling (Arctic)	U	10,000	96	3	Fish swam near the surface	McLeay et al. (1987)
(1983); Ott (1984)	Grayling (Arctic)	J	86	0.42	3	78% of fish avoided turbid water (NTU, >20)	Scannell (1988)
al. (1977b)	Grayling (Arctic)	U	100	1	4	Catch rate reduced (unfamiliar prey: drosophila)	McLeay et al. (1987)
and Merkens (1961)	Grayling (Arctic)	U	100	1	4	Catch rate reduced (unfamiliar prey: tubificids)	McLeay et al. (1987)
38)	Grayling (Arctic)	U	300	1	4	Catch rate reduced (unfamiliar prey: drosophila)	McLeay et al. (1987)
57)	Grayling (Arctic)	U	1,000	1	4	Feeding rate reduced (unfamiliar prey: tubificids)	McLeay et al. (1987)
and Swenson (1982)							

TABLE A.1.—Continued.

TABLE A.1.—Co

Species	Life stage ^a	Sediment dose		Fish response		Reference	Species
		Exposure concentration (mg/L)	Exposure duration (h)	SEV ^b	Description ^c		
Grayling (Arctic)	U	1,000	1	4	Feeding rate reduced (unfamiliar prey: drosophila)	McLeay et al. (1987)	Salmon (coho)
Grayling (Arctic)	YY	3,810	144	4	Food intake severely limited	Simmons (1982)	Salmon (coho)
Grayling (Arctic)	U	100	12	6	Reduced ability to tolerate high temperatures	McLeay et al. (1987)	Salmon (coho)
Grayling (Arctic)	U	100	756	7	Fish moved out of the test channel	McLeay et al. (1987)	Salmon (coho)
Grayling (Arctic)	U	1,000	1,008	8	Fish had frequent misstrikes while feeding	McLeay et al. (1987)	Salmon (coho)
Grayling (Arctic)	U	1,000	1,008	8	Fish responded very slowly to prey	McLeay et al. (1987)	Salmon (coho)
Grayling (Arctic)	U	300	1,008	8	Rate of feeding reduced	McLeay et al. (1987)	Salmon (coho)
Grayling (Arctic)	U	1,000	840	8	Rate of feeding reduced	McLeay et al. (1987)	Salmon (coho)
Grayling (Arctic)	U	1,000	1,008	8	Fish failed to consume all prey	McLeay et al. (1987)	Salmon (coho)
Grayling (Arctic)	U	300	840	8	Serious impairment of feeding behavior	McLeay et al. (1987)	Salmon (coho)
Grayling (Arctic)	U	300	1,008	8	Respiration rate increased (FSS)	McLeay et al. (1987)	Salmon (coho)
Grayling (Arctic)	U	300	1,008	8	Fish less tolerant of pentachlorophenol	McLeay et al. (1987)	Salmon (coho)
Grayling (Arctic)	YY	3,810	144	8	Mucus and sediment accumulated in the gill lamellae	Simmons (1982)	Salmon (coho)
Grayling (Arctic)	YY	3,810	144	8	Fish displayed many signs of poor condition	Simmons (1982)	Salmon (coho)
Grayling (Arctic)	YY	1,250	48	8	Moderate damage to gill tissue	Simmons (1982)	Salmon (sockeye)
Grayling (Arctic)	YY	1,388	96	8	Hyperplasia and hypertrophy of gill tissue	Simmons (1982)	Salmon (sockeye)
Grayling (Arctic)	U	100	1,008	9	Growth rate reduced	McLeay et al. (1984)	Salmon (sockeye)
Grayling (Arctic)	U	100	840	9	Fish responded less rapidly to drifting food	McLeay et al. (1987)	Salmon (sockeye)
Grayling (Arctic)	U	300	1,008	9	Weight gain reduced	McLeay et al. (1987)	Salmon (sockeye)
Grayling (Arctic)	U	1,000	1,008	9	Weight gained reduced by 33%	McLeay et al. (1987)	Salmon (sockeye)
Grayling (Arctic)	U	300	756	10	Fish displaced from their habitat	McLeay et al. (1987)	Salmon (sockeye)
Grayling (Arctic)	U	100,000	168	5	No changes in gill histology (not an end point)	McLeay et al. (1983)	Salmon (sockeye)
Salmon (chinook)	S	94.5	72	8	Tolerance to stress reduced (VA)	Stober et al. (1981)	Salmon (sockeye)
Salmon (chinook)	J	6	1,440	9	Growth rate reduced (LNFH)	MacKinley et al. (1987)	Salmon (sockeye)
Salmon (chinook)	J	1,400	36	12	Mortality rate 50%	Newcomb and Flagg (1983)	Salmon (sockeye)
Salmon (chinook)	J	9,400	36	12	Mortality rate 50%	Newcomb and Flagg (1983)	Salmon (sockeye)
Salmon (chinook)	S	488	96	12	Mortality rate 50%	Stober et al. (1981)	Salmon (sockeye)
Salmon (chinook)	S	11,000	96	12	Mortality rate 50%	Stober et al. (1981)	Salmon (sockeye)
Salmon (chinook)	S	19,364	96	12	Mortality rate 50%	Stober et al. (1981)	Salmon (sockeye)
Salmon (chinook)	J	39,400	36	14	Mortality rate 90% (VA)	Newcomb and Flagg (1983)	Salmon (sockeye)
Salmon (chum)	J	28,000	96	12	Mortality rate 50%	Smith (1940)	Salmon (sockeye)
Salmon (chum)	J	55,000	96	12	Mortality rate 50% (winter)	Smith (1940)	Salmon (sockeye)
Salmon (coho)	J	53.5	0.02	1	Alarm reaction	Berg (1983)	Salmon (sockeye)
Salmon (coho)	J	88	0.02	1	Alarm reaction	Bisson and Bilby (1982)	Salmon (sockeye)
Salmon (coho)	U	20	0.05	1	Cough frequency not increased	Servizi and Martens (1992)	Salmon (sockeye)
Salmon (coho)	J	53.5	12	3	Changes in territorial behavior	Berg and Northcote (1985)	Salmon (sockeye)
Salmon (coho)	J	88	0.08	3	Avoidance behavior	Bisson and Bilby (1982)	Salmon (sockeye)
Salmon (coho)	J	6,000	1	3	Avoidance behavior	Noggle (1978)	Steelhead
Salmon (coho)	U	300	0.17	3	Avoidance behavior within minutes	Servizi and Martens (1992)	Trout (brook)
Salmon (coho)	J	25	1	4	Feeding rate decreased	Noggle (1978)	Trout (brook)
Salmon (coho)	J	100	1	4	Feeding rate decreased to 55% of maximum	Noggle (1978)	Trout (brook)
Salmon (coho)	J	250	1	4	Feeding rate decreased to 10% of maximum	Noggle (1978)	Trout (rainbow)
Salmon (coho)	J	300	1	4	Feeding ceased	Noggle (1978)	Trout (rainbow)
Salmon (coho)	U	2,460	0.05	5	Coughing behavior manifest within minutes	Servizi and Martens (1992)	Trout (rainbow)

TABLE A.1.—Continued.

Reference	Species	Life stage ^a	Sediment dose		Fish response		Reference
			Exposure concentration (mg/L)	Exposure duration (h)	SEV ^b	Description ^c	
et al. (1987)	Salmon (coho)	J	53.5	12	6	Increased physiological stress	Berg and Northcote (1985)
(1982)	Salmon (coho)	U	2,460	1	6	Cough frequency greatly increased	Servizi and Martens (1992)
et al. (1987)	Salmon (coho)	U	240	24	6	Cough frequency increased more than 5-fold	Servizi and Martens (1992)
et al. (1987)	Salmon (coho)	U	530	96	6	Blood glucose levels increased	Servizi and Martens (1992)
et al. (1987)	Salmon (coho)	J	1,547	96	8	Gill damage	Noggle (1978)
et al. (1987)	Salmon (coho)	U	2,460	24	8	Fatigue of the cough reflex	Servizi and Martens (1992)
et al. (1987)	Salmon (coho)	U	3,000	48	8	High level sublethal stress: avoidance	Servizi and Martens (1992)
et al. (1987)	Salmon (coho)	J	102	336	9	Growth rate reduced (FC, BC)	Sigler et al. (1984)
et al. (1987)	Salmon (coho)	U	8,000	96	10	Mortality rate 1%	Servizi and Martens (1991)
et al. (1987)	Salmon (coho)	J	1,200	96	12	Mortality rate 50%	Noggle (1978)
et al. (1987)	Salmon (coho)	J	35,000	96	12	Mortality rate 50%	Noggle (1978)
et al. (1987)	Salmon (coho)	U	22,700	96	12	Mortality rate 50%	Servizi and Martens (1991)
et al. (1987)	Salmon (coho)	F*	8,100	96	12	Mortality rate 50%	Servizi and Martens (1991)
et al. (1987)	Salmon (coho)	PS	18,672	96	12	Mortality rate 50%	Stober et al. (1981)
et al. (1987)	Salmon (coho)	S	509	96	12	Mortality rate 50%	Stober et al. (1981)
(1982)	Salmon (coho)	S	1,217	96	12	Mortality rate 50% (VA)	Stober et al. (1981)
(1982)	Salmon (coho)	S	28,184	96	12	Mortality rate 50% (VA)	Stober et al. (1981)
(1982)	Salmon (coho)	S	29,580	96	12	Mortality rate 50%	Stober et al. (1981)
(1982)	Salmon (sockeye)	S	1,261	96	8	Body moisture content reduced	Servizi and Martens (1987)
(1982)	Salmon (sockeye)	S	7,447	96	8	Plasma chloride levels increased slightly	Servizi and Martens (1987)
(1982)	Salmon (sockeye)	U	1,465	96	8	Hypertrophy and necrosis of gill tissue (CSS)	Servizi and Martens (1987)
et al. (1984)	Salmon (sockeye)	U	3,143	96	8	Hypertrophy and necrosis of gill tissue (FSS)	Servizi and Martens (1987)
et al. (1987)	Salmon (sockeye)	U	9,851	96	8	Hypertrophy and necrosis of gill tissue (MCSS)	Servizi and Martens (1987)
et al. (1987)	Salmon (sockeye)	U	17,560	96	8	Hypertrophy of gill tissue (FSS)	Servizi and Martens (1987)
et al. (1987)	Salmon (sockeye)	U	23,790	96	8	Hypertrophy and necrosis of gill tissue (FSS)	Servizi and Martens (1987)
et al. (1983)	Salmon (sockeye)	U	2,683	96	8	Hypertrophy and necrosis of gill tissue (MCSS)	Servizi and Martens (1987)
et al. (1981)	Salmon (sockeye)	U	2,100	96	10	No fish died (MFSS)	Servizi and Martens (1987)
ey et al. (1987)	Salmon (sockeye)	U	9,000	96	10	No mortality	Servizi and Martens (1987)
ib and Flagg (1983)	Salmon (sockeye)	U	13,900	96	10	Mortality rate 10% (FSS)	Servizi and Martens (1987)
ib and Flagg (1983)	Salmon (sockeye)	U	9,850	96	10	Gill hyperplasia, hypertrophy, separation, necrosis (MFSS)	Servizi and Martens (1987)
et al. (1981)	Salmon (sockeye)	J	1,400	36	12	Mortality rate 50%	Newcomb and Flagg (1983)
et al. (1981)	Salmon (sockeye)	J	9,400	36	12	Mortality rate 50%	Newcomb and Flagg (1983)
ib and Flagg (1983)	Salmon (sockeye)	U	1,700	96	12	Mortality rate 50% (CSS)	Servizi and Martens (1987)
940)	Salmon (sockeye)	U	4,850	96	12	Mortality rate 50% (MCSS)	Servizi and Martens (1987)
83)	Salmon (sockeye)	U	8,200	96	12	Mortality rate 50% (MFSS)	Servizi and Martens (1987)
ind Bilby (1982)	Salmon (sockeye)	U	17,560	96	12	Mortality rate 50% (FSS)	Servizi and Martens (1987)
and Martens (1992)	Salmon (sockeye)	J	39,400	36	14	Mortality rate 90% (VA)	Newcomb and Flagg (1983)
d Northcote (1985)	Salmon (sockeye)	U	13,000	96	14	Mortality rate 90% (MFSS)	Servizi and Martens (1987)
ind Bilby (1982)	Salmon (sockeye)	U	23,900	96	14	Mortality rate 90% (FSS)	Servizi and Martens (1987)
(1978)	Sieelhead	J	102	336	9	Growth rate reduced (FC, BC)	Sigler et al. (1984)
and Martens (1992)	Trout (brook)	FF	12	5,880	9	Growth rates declined	Sykora et al. (1972)
(1978)	Trout (brook)	FF	24	5,208	9	Growth rate reduced (LNFH)	Sykora et al. (1972)
(1978)	Trout (brook)	FF*	100	1,176	9	Test fish weighed 16% of controls (LNFH)	Sykora et al. (1972)
(1978)	Trout (brook)	FF	50	1,848	9	Growth rates declined (LNFH)	Sykora et al. (1972)
(1978)	Trout (rainbow)	FF	1,750	480	12	Mortality rate 57% (controls 5%)	Campbell (1954)
(1978)	Trout (rainbow)	J	4,887	384	8	Hyperplasia of gill tissue	Golde (1983)
(1978)	Trout (rainbow)	J	4,887	384	8	Parasitic infection of gill tissue	Golde (1983)
and Martens (1992)	Trout (rainbow)	J	171	96	8	Particles penetrated cells of branchial epithelium	Golde (1983)

TABLE A.1.—Continued.

Species	Life stage ^a	Sediment dose		Fish response		Reference	Species
		Exposure concentration (mg/L)	Exposure duration (h)	SEV ^b	Description ^c		
Trout (rainbow)	Y	90	456	10	Mortality rates 0–20% (DE)	Herbert and Merckens (1961)	Herring
Trout (rainbow)	Y	90	456	10	Mortality rates 0–15% (KC)	Herbert and Merckens (1961)	Herring (lake)
Trout (rainbow)	Y	270	456	11	Mortality rates 10–35% (KC)	Herbert and Merckens (1961)	Herring (Pacific)
Trout (rainbow)	Y	810	456	12	Mortality rates 35–85% (DE)	Herbert and Merckens (1961)	Herring (Pacific)
Trout (rainbow)	Y	810	456	12	Mortality rates 5–80% (KC)	Herbert and Merckens (1961)	Herring (Pacific)
Trout (rainbow)	Y	270	456	12	Mortality rates 25–80% (DE)	Herbert and Merckens (1961)	
Trout (rainbow)	Y	7,433	672	11	Mortality rate 40% (CS)	Herbert and Wakeford (1962)	Perch (white)
Trout (rainbow)	Y	4,250	672	12	Mortality rate 50%	Herbert and Wakeford (1962)	
Trout (rainbow)	Y	2,120	672	14	Mortality rate 100%	Herbert and Wakeford (1962)	
Trout (rainbow)	J	4,315	57	14	Mortality rate ~100% (CSS)	Newcombe et al. (1995)	Perch (white)
Salmonid eggs and larvae (freshwater, group 4)							
Grayling (Arctic)	SF	25	24	10	Mortality rate 5.7%	J. LaPerriere (personal communication)	Perch (white)
Grayling (Arctic)	SF	22.5	48	10	Mortality rate 14.0%	J. LaPerriere (personal communication)	Perch (white)
Grayling (Arctic)	SF	65	24	10	Mortality rate 15.0%	J. LaPerriere (personal communication)	Perch (white)
Grayling (Arctic)	SF	21.7	72	10	Mortality rate 14.7%	J. LaPerriere (personal communication)	Perch (white)
Grayling (Arctic)	SF	20	96	10	Mortality rate 13.4%	J. LaPerriere (personal communication)	Perch (white)
Grayling (Arctic)	SF	142.5	48	11	Mortality rate 26%	J. LaPerriere (personal communication)	Perch (yellow)
Grayling (Arctic)	SF	185	72	12	Mortality rate 41.3%	J. LaPerriere (personal communication)	Perch (yellow)
Grayling (Arctic)	SF	230	96	12	Mortality rate of 47%	J. LaPerriere (personal communication)	Shad (American)
Salmon	E	117	960	10	Mortality; deterioration of spawning gravel	Cederholm et al. (1981)	Shad (American)
Salmon (chum)	E	97	2,808	13	Mortality rate 77% (controls, 6%)	Langer (1980)	Anchovy (bay)
Salmon (coho)	E	157	1,728	14	Mortality rate 100% (controls, 16.2%)	Shaw and Maga (1943)	Anchovy (bay)
Steelhead	E	57	1,488	12	Hatching success 42% (controls, 63%)	Slaney et al. (1977b)	Anchovy (bay)
Trout	E	117	960	10	Mortality; deterioration of spawning gravel	Cederholm et al. (1981)	Bass (striped)
Trout (rainbow)	EE	1,750	144	10	Mortality rate greater than controls (controls, 6%)	Campbell (1954)	Bass (striped)
Trout (rainbow)	E	6.6	1,152	11	Mortality rate 40%	Slaney et al. (1977b)	Bass (striped)
Trout (rainbow)	E	57	1,488	12	Mortality rate 47% (controls, 32%)	Slaney et al. (1977b)	Bass (striped)
Trout (rainbow)	E	120	384	13	Mortality rates 60–70% (controls, 38.6%)	Erman and Lignon (1988)	Bass (striped)
Trout (rainbow)	E	20.8	1,152	13	Mortality rate 72%	Slaney et al. (1977a)	Menhaden (Atlantic)
Trout (rainbow)	E	46.6	1,152	14	Mortality rate 100%	Slaney et al. (1977b)	Menhaden (Atlantic)
Trout (rainbow)	E	101	1,440	14	Mortality rate 98% (controls, 14.6%)	Turnpenny and Williams (1980)	Minnow (sheeps)
Nonsalmonid eggs and larvae (estuarine^d, group 4)							
Bass (striped)	L	200	0.42	4	Feeding rate reduced 40%	Breitburg (1988)	Minnow (sheeps)
Bass (striped)	E	800	24	9	Development rate slowed significantly	Morgan et al. (1983)	Mummichog
Bass (striped)	E	100	24	9	Hatching delayed	Schubel and Wang (1973)	Mummichog
Bass (striped)	E	1,000	168	10	Reduced hatching success	Auld and Schubel (1978)	Mummichog
Bass (striped)	L	1,000	68	11	Mortality rate 35% (controls, 16%)	Auld and Schubel (1978)	Perch (white)
Bass (striped)	L	500	72	12	Mortality rate 42% (controls, 17%)	Auld and Schubel (1978)	Perch (white)
Bass (striped)	L	485	24	12	Mortality rate 50%	Morgan et al. (1973)	Perch (white)

TABLE A.1.—Continued.

Author(s)	Species	Life stage ^a	Sediment dose		Fish response		Reference
			Exposure concentration (mg/L)	Exposure duration (h)	SEV ^b	Description ^c	
ns (1961)	Herring	L	10	3	3	Depth preference changed	Johnson and Wildish (1982)
ns (1961)	Herring (lake)	L	16	24	3	Depth preference changed	Swenson and Matson (1976)
ns (1961)	Herring (Pacific)	L	2,000	2	4	Feeding rate reduced	Boehlert and Morgan (1985)
ns (1961)	Herring (Pacific)	L	1,000	24	8	Mechanical damage to epidermis	Boehlert (1984)
ns (1961)	Herring (Pacific)	L	4,000	24	8	Epidermis punctured; microridges less distinct	Boehlert (1984)
ford (1962)	Perch (white)	E	800	24	9	Egg development slowed significantly	Morgan et al. (1983)
ford (1962)	Perch (white)	E	100	24	9	Hatching delayed	Schubel and Wang (1973)
ford (1962)	Perch (white)	E	1,000	168	10	Reduced hatching success	Auld and Schubel (1978)
(1995)	Perch (white)	L	155	48	12	Mortality rate 50%	Morgan et al. (1973)
	Perch (white)	L	375	24	12	Mortality rate 50%	Morgan et al. (1973)
	Perch (white)	L	280	48	12	Mortality rate 50%	Morgan et al. (1973)
	Perch (yellow)	L	500	96	11	Mortality rate 37% (controls, 7%)	Auld and Schubel (1978)
	Perch (yellow)	L	1,000	96	11	Mortality rate 38% (controls, 7%)	Auld and Schubel (1978)
	Shad (American)	L	100	96	10	Mortality rate 18% (controls, 5%)	Auld and Schubel (1978)
	Shad (American)	L	500	96	11	Mortality rate 36% (controls, 4%)	Auld and Schubel (1978)
	Shad (American)	L	1,000	96	11	Mortality rate 34% (controls, 5%)	Auld and Schubel (1978)
Adult nonsalmonids (estuarine or riverine-estuarine, group 5)							
	Anchovy (bay)	A	231	24	10	Mortality rate 10% (FE)	Sherk et al. (1975)
	Anchovy (bay)	A	471	24	12	Mortality rate 50% (FE)	Sherk et al. (1975)
	Anchovy (bay)	A	966	24	14	Mortality rate 90%	Sherk et al. (1975)
	Bass (striped)	A	1,500	336	8	Haematocrit increased (FE)	Sherk et al. (1975)
	Bass (striped)	A	1,500	336	8	Plasma osmolality increased (FE)	Sherk et al. (1975)
	Cunner	A	28,000	24	12	Mortality rate 50% (20.0–25.0°C)	Rogers (1969)
	Cunner	A	133,000	12	12	Mortality rate 50% (15°C)	Rogers (1969)
	Cunner	A	100,000	24	12	Mortality rate 50% (15°C)	Rogers (1969)
	Cunner	A	72,000	48	12	Mortality rate 50% (15°C)	Rogers (1969)
	Fish	A	3,000	240	10	Fish died	Kemp (1949)
(1943)	Herring (Atlantic)	A	20	3	4	Reduced feeding rate	Johnson and Wildish (1982)
	Hogchoker	A	1,240	24	8	Energy utilization increased	Sherk et al. (1975)
77b)	Hogchoker	A	1,240	120	8	Erythrocyte count increased	Sherk et al. (1975)
	Hogchoker	A	1,240	120	8	Haematocrit increased	Sherk et al. (1975)
(1981)	Killifish (striped)	A	960	120	8	Haematocrit increased	Sherk et al. (1975)
	Killifish (striped)	A	3,277	24	10	Mortality rate 10% (FE)	Sherk et al. (1975)
	Killifish (striped)	A	9,720	24	10	Mortality rate 10%	Sherk et al. (1975)
	Killifish (striped)	A	3,819	24	12	Mortality rate 50%	Sherk et al. (1975)
	Killifish (striped)	A	12,820	24	12	Mortality rate 50%	Sherk et al. (1975)
77b)	Killifish (striped)	A	16,930	24	13	Mortality rate 90%	Sherk et al. (1975)
77b)	Killifish (striped)	A	6,136	24	14	Mortality rate 90%	Sherk et al. (1975)
on (1988)	Menhaden (Atlantic)	A	154	24	10	Mortality rate 10% (FE)	Sherk et al. (1975)
	Menhaden (Atlantic)	A	247	24	12	Mortality rate 50% (FE)	Sherk et al. (1975)
	Menhaden (Atlantic)	A	396	24	14	Mortality rate 90% (FE)	Sherk et al. (1975)
77a)	Minnnow (sheepshead)	A	200,000	24	10	Mortality rate 10% (15°C)	Rogers (1969)
77b)	Minnnow (sheepshead)	A	300,000	24	11	Mortality rate 30% (10°C)	Rogers (1969)
Williams	Minnnow (sheepshead)	A	100,000	24	14	Mortality rate 90% (19°C)	Rogers (1969)
	Mummichog	A	300,000	24	10	No mortality (15°C)	Rogers (1969)
	Mummichog	A	2,447	24	10	Mortality rate 10% (FE)	Sherk et al. (1975)
	Mummichog	A	3,900	24	12	Mortality rate 50% (FE)	Sherk et al. (1975)
	Mummichog	A	6,217	24	14	Mortality rate 90%	Sherk et al. (1975)
1983)	Perch (white)	A	650	120	6	Haematocrit increased	Sherk et al. (1975)
	Perch (white)	A	650	120	6	Erythrocyte count increased	Sherk et al. (1975)
	Perch (white)	A	650	120	6	Haemoglobin concentration increased	Sherk et al. (1975)
ang (1973)	Perch (white)	A	305	120	8	Gill tissue may have been damaged	Sherk et al. (1975)
del (1978)	Perch (white)	A	650	120	8	Histological damage to gill tissue	Sherk et al. (1975)
del (1978)	Perch (white)	A	305	24	10	Mortality rate 10% (FE)	Sherk et al. (1975)
1973)	Perch (white)	A	985	24	12	Mortality rate 50%	Sherk et al. (1975)

TABLE A.1.—Continued.

TABLE A.1.—C

Species	Life stage ^a	Sediment dose		Fish response		Reference	Species
		Exposure concentration (mg/L)	Exposure duration (h)	SEV ^b	Description ^c		
Perch (white)	A	3.181	24	14	Mortality rate 90% (FE)	Sherk et al. (1975)	Sunfish (green)
Rasbora (harlequin)	A	40.000	24	10	Fish died (BC)	Alabaster and Lloyd (1980)	Sunfish (redear)
Rasbora (harlequin)	A	6.000	168	10	No mortality	Alabaster and Lloyd (1980)	
Shad (American)	A	150	0.25	3	Change in preferred swimming depth	Dadswell et al. (1983)	Sunfish (redear)
Silverside (Atlantic)	A	58	24	10	Mortality rate 10% (FE)	Sherk et al. (1975)	Sunfish (redear)
Silverside (Atlantic)	A	250	24	12	Mortality rate 50% (FE)	Sherk et al. (1975)	
Silverside (Atlantic)	A	1.000	24	14	Mortality rate 90% (FE)	Sherk et al. (1975)	
Spot	A	114	48	10	Mortality rate 10% (FE)	Sherk et al. (1975)	
Spot	A	1.309	24	10	Mortality rate 10% (FE)	Sherk et al. (1975)	
Spot	A	6.875	24	10	Mortality rate 10%	Sherk et al. (1975)	
Spot	A	189	48	12	Mortality rate 50% (FE)	Sherk et al. (1975)	
Spot	A	2.034	24	12	Mortality rate 50%	Sherk et al. (1975)	
Spot	A	8.800	24	12	Mortality rate 50%	Sherk et al. (1975)	
Spot	A	317	48	14	Mortality rate 90% (FE)	Sherk et al. (1975)	
Spot	A	11.263	24	14	Mortality rate 90%	Sherk et al. (1975)	
Stickleback (fourspine)	A	100	24	10	Mortality rate <1% (IA)	Rogers (1969)	
Stickleback (fourspine)	A	10.000	24	10	No mortality (KS; 10–12°C)	Rogers (1969)	
Stickleback (fourspine)	A	300	24	12	Mortality rate ~50% (IA)	Rogers (1969)	
Stickleback (fourspine)	A	18.000	24	12	Mortality rate 50% (15.0–16.0°C)	Rogers (1969)	
Stickleback (fourspine)	A	50.000	24	12	Mortality rate 50% (KS)	Rogers (1969)	
Stickleback (fourspine)	A	53.000	24	12	Mortality rate 50% (10–12°C)	Rogers (1969)	
Stickleback (fourspine)	A	330.000	24	12	Mortality rate 50% (9.0–9.5°C)	Rogers (1969)	
Stickleback (fourspine)	A	500	24	14	Mortality rate 100%	Rogers (1969)	
Stickleback (fourspine)	A	200.000	24	14	Mortality rate 95% (KS)	Rogers (1969)	
Stickleback (threespine)	A	28.000	96	10	No mortality in test designed to identify lethal threshold	LeGore and DesVoigne (1973)	
Toadfish (oyster)	A	3.360	1	6	Oxygen consumption more variable in prestressed fish	Neumann et al. (1975)	
Toadfish (oyster)	A	14.600	72	8	Fish largely unaffected, but developed latent ill effects	Neumann et al. (1975)	
Toadfish (oyster)	A	11.090	72	9	Latent ill effects manifested in subsequent test at low SS	Neumann et al. (1975)	
Adult nonsalmonids (freshwater, group 6)							
Bass (largemouth)	A	62.5	720	9	Weight gain reduced ~50%	Buck (1956)	
Bass (largemouth)	A	144.5	720	9	Growth retarded	Buck (1956)	
Bass (largemouth)	A	144.5	720	12	Fish unable to reproduce	Buck (1956)	
Bluegill	A	423	0.05	4	Rate of feeding reduced	Gardner (1981)	
Bluegill	A	15	1	4	Reduced capacity to locate prey	Vinyard and O'Brien (1976)	
Bluegill	A	144.5	720	9	Growth retarded	Buck (1956)	
Bluegill	A	62.5	720	9	Weight gain reduced ~50%	Buck (1956)	
Bluegill	A	144.5	720	12	Fish unable to reproduce	Buck (1956)	
Carp (common)	A	25.000	336	10	Some mortality (MC)	Wallen (1951)	
Darters	A	2.045	8,760	14	Darters absent	Vaughan (1979); Vaughan et al. (1982)	
Fish	A	120	384	10	Density of fish reduced	Erman and Lignon (1988)	
Fish	A	620	48	10	Fish kills downstream from sediment source	Hesse and Newcomb (1982)	
Fish	A	900	720	12	Fish absent or markedly reduced in abundance	Herbert and Richards (1963)	
Fish	A	2.045	8,760	12	Habitat destruction; fish populations smaller than expected	Vaughan (1979); Vaughan et al. (1982)	
Fish (warmwater)	A	100.000	252	10	Some fish died; most survived	Wallen (1951)	
Fish (warmwater)	A	200.000	1.125	10	Fish died; opercular cavities and gill filaments clogged	Wallen (1951)	
Fish (warmwater)	A	22	8,760	12	Fish populations destroyed	Menzel et al. (1984)	
Goldfish	A	25.000	336	10	Some mortality (MC)	Wallen (1951)	

^a A = adult; E = egg; J = juvenile; L = larva; of the year.

^b Severity-of-ill-effect.

^c Full response annotation.

source documents.

MCSS = medium calcium sulfate; CW = earth; IA = incinerator; VA = volcanic clay; VA = volcanic ash.

^d Lake herring larvae.

TABLE A.1.—Continued.

	Species	Life stage ^a	Sediment dose		Fish response		Reference
			Exposure concentration (mg/L)	Exposure duration (h)	SEV ^b	Description ^c	
1 (1980)	Sunfish (green)	A	9,600	1	5	Rate of ventilation increased	Horkel and Pearson (1976)
1 (1980)	Sunfish (reardear)	A	62.5	720	9	Weight gain reduced ~50% compared to controls	Buck (1956)
33)	Sunfish (reardear)	A	144.5	720	9	Growth retarded	Buck (1956)
	Sunfish (reardear)	A	144.5	720	12	Fish unable to reproduce	Buck (1956)

^a A = adult; E = egg; EE = eyed egg; F = fry; F* = swim-up fry; FF = young fry (<30 weeks old); FF* = older fry (>30 weeks old); J = juvenile; L = larva; PS = presmolt; S = smolt; SF = sac fry; U = underyearling; Y = approximate yearling; YY = young of the year.

^b Severity-of-ill-effect ranging from 0 (no detectible effect) to 14 (maximum effect; see Table 1).

^c Full response annotations are in Newcombe (1994). Particle sizes of suspended sediment (SS) sometimes were given categorically in source documents. As abbreviated here, VFSS = very fine (<15 μm); FSS = fine (15–74 μm); MFSS = medium to fine (75–149 μm); MCSS = medium to coarse (150–290 μm); and CSS = coarse (180–740 μm). Usual "sediments" used: BC = bentonite clay; CS = calcium sulfate; CWS = coal washery solids; DE = diatomaceous earth; DM = drilling mud (nontoxic); FC = fire clay; FE = fuller's earth; IA = incinerator ash; KC = kaolin clay; KS = Kingston silt; LNFH = lime-neutralized ferric hydroxide; MC = montmorillonite clay; VA = volcanic ash; WF = wood fibers. Other abbreviation: NTU = nephelometric turbidity units.

^d Lake herring larvae were tested in freshwater.

igne

75)

75)

75)

:n (1976)

aughan et

(1988)

ib (1982)

ds (1963)

aughan et

!)