

1

APPENDIX 2.C

2

CLIMATE CHANGE IMPLICATIONS AND ASSUMPTIONS

3

4

REVISED ADMINISTRATIVE DRAFT

5

BAY DELTA CONSERVATION PLAN

6

7

8

9

March 2013

10

11



1 ICF International. 2013. *Appendix 2.C: Climate Change Implications and*
2 *Assumptions. Bay Delta Conservation Plan*. Revised Administrative Draft.
3 March. (ICF 00343.12). Sacramento, CA. Prepared for: California
4 Department of Water Resources, Sacramento, CA.

Climate Change Implications and Assumptions

Contents

	Page
Appendix 2.C Climate Change Implications and Assumptions	2.C-1
2.C.1 Global Climate Change	2.C-1
2.C.1.1 What Is Climate Change?	2.C-1
2.C.1.2 What Causes Climate Change?	2.C-2
2.C.1.3 What Global Climate Changes Have Been Observed?	2.C-3
2.C.1.4 How is Global Climate Expected to Change?	2.C-3
2.C.2 Climate Change in California and the Plan Area	2.C-4
2.C.2.1 Current Climate	2.C-4
2.C.2.2 Surface Temperature	2.C-4
2.C.2.2.1 Observed	2.C-4
2.C.2.2.2 Projected	2.C-5
2.C.2.3 Water Temperature	2.C-6
2.C.2.4 Precipitation	2.C-7
2.C.2.4.1 Observed	2.C-7
2.C.2.4.2 Projected	2.C-7
2.C.2.5 Snowpack and Runoff	2.C-9
2.C.2.5.1 Observed	2.C-9
2.C.2.5.2 Projected	2.C-10
2.C.2.6 Sea Level Rise	2.C-10
2.C.2.6.1 Observed	2.C-10
2.C.2.6.2 Projected	2.C-10
2.C.2.7 Ocean Conditions	2.C-13
2.C.2.7.1 Observed	2.C-13
2.C.2.7.2 Projected	2.C-14
2.C.2.8 Extreme Weather Events	2.C-15
2.C.2.8.1 Observed	2.C-15
2.C.2.8.2 Projected	2.C-15
2.C.3 References Cited	2.C-16

1 List of Figures

2	Page
3	
4	Figure 2.C-1. The Greenhouse Effect 2.C-2
5	Figure 2.C-2. Historical Observed California Statewide Mean Annual Temperature Departure..... 2.C-5
6	Figure 2.C-3. Simulated Historical and Future Annual Temperature Projections for the
7	Sacramento Region..... 2.C-6
8	Figure 2.C-4. Historical Observed California Statewide Water Year Precipitation 2.C-8
9	Figure 2.C-5. Simulated Future Water Year Change in Precipitation for the
10	Intergovernmental Panel on Climate Change A2 and B1 Scenarios for the
11	Sacramento Region..... 2.C-9
12	Figure 2.C-6. Observed Mean Sea Level Trend for the San Francisco Tide Gage near the
13	Golden Gate 2.C-11
14	Figure 2.C-7. Past Global Mean Sea Level and Future Mean Sea Level Based on Global Mean
15	Temperature Projections 2.C-12
16	Figure 2.C-8. DWR-Generated Future Sea Level Rise Projections for the Bay Delta Using the
17	Rahmstorf Method and Regionally Downscaled Data 2.C-13
18	

1 Acronyms and Abbreviations

°F	degrees Fahrenheit
Bay-Delta	San Francisco Bay/Sacramento–San Joaquin River Delta
CALFED	CALFED Bay-Delta Program
CH ₄	methane
CO ₂	carbon dioxide
Delta	Sacramento–San Joaquin Delta
DWR	California Department of Water Resources
ENSO	El Niño Southern Oscillation
GCM	Global Circulation Model
GHGs	greenhouse gases
H ₂ O	water vapor
IPCC	Intergovernmental Panel on Climate Change
N ₂ O	nitrous oxide
NRC	National Research Council
O ₃	ozone
PDO	Pacific Decadal Oscillation
SRES	Special Report on Emissions Scenarios

2

Climate Change Implications and Assumptions

Observations from around the world demonstrate that the Earth is undergoing climate change much more quickly than would be expected from natural variation. In its most recent assessment of climate change, the Intergovernmental Panel on Climate Change, an international body made up of scientists from around the world, stated that, “Warming of the climate system is unequivocal, as is now evident from observations of increases in global average air and ocean temperatures, widespread melting of snow and ice, and rising global average sea level” (Intergovernmental Panel on Climate Change 2007). The Intergovernmental Panel on Climate Change concluded that, “At continental, regional and ocean basin scales, numerous long-term changes in climate have been observed. These include changes in arctic temperatures and ice, widespread changes in precipitation amounts, ocean salinity, wind patterns and aspects of extreme weather including droughts, heavy precipitation, heat waves and the intensity of tropical cyclones” (Intergovernmental Panel on Climate Change 2007). In a 2010 report, the National Research Council (NRC) concluded that, “There is a strong, credible body of evidence, based on multiple lines of research, documenting that climate is changing and these changes are in large part caused by human activities” (National Research Council 2010).

This appendix provides an overview of scientific understanding of climate change and observed and projected changes in California and the Plan Area. The focus is on the physical basis of climate change. Subsequent appendices discuss the ecological implications of these changes.

2.C.1 Global Climate Change

2.C.1.1 What Is Climate Change?


Climate is the average weather over a specific region over many years, measured most often in terms of temperature and precipitation. *Climate change* refers to a statistically significant change in the state of the climate or its variability that persists for an extended period (typically decades or longer). An individual year that is drier or hotter than average would indicate *climate variability* but might not indicate climate change, which is a large-scale shift or trend in the average weather that a region experiences.

The climate changes over many temporal and spatial scales as a result of meteorological processes such as variations in atmospheric circulation patterns. These changes may be due to natural processes or to anthropogenic factors that affect the composition of the atmosphere. Although the climate has changed in the past in response to natural drivers, recent climate change has been linked unequivocally to increasing concentrations of greenhouse gases (GHGs) in the Earth’s lower atmosphere, largely as a result of the burning of fossil fuels since the beginning of the industrial revolution (Intergovernmental Panel on Climate Change 2007).

1 2.C.1.2 What Causes Climate Change?

2 Higher concentrations of heat-trapping GHGs in the atmosphere, as well as modifications to the land
 3 surface, alter the energy balance of the climate system, increasing the Earth's temperature and thus
 4 influencing climate. The so-called "greenhouse effect" is a natural phenomenon in which GHGs—
 5 primarily water vapor (H₂O), carbon dioxide (CO₂), methane (CH₄), nitrous oxide (N₂O), and ozone
 6 (O₃)—allow solar radiation to pass through the atmosphere and warm the Earth's surface (Figure
 7 2.C-1). As the Earth's surface warms, infrared radiation is emitted back to the atmosphere, where
 8 GHGs absorb some of the radiation and re-emit it back to Earth, causing the surface to gain more
 9 heat (National Academy of Sciences 2006).

10 This warming of the Earth's surface has been increasing rapidly as levels of atmospheric CO₂ have
 11 increased since the onset of the Industrial Revolution, rising by about 30%, since the late 1800s. The
 12 atmospheric concentration of CO₂ is now higher than it has been for many millennia. Observations
 13 show that the global average surface temperature is rising at an increasing rate, with the decades
 14 from 1970 to 2009 being progressively warmer than prior decades (Arndt et al. 2010). Of the 10
 15 warmest years on record, 9 have occurred since 2001 (National Climatic Data Center 2011). Based
 16 on the current trajectory, the Intergovernmental Panel on Climate Change projects that atmospheric
 17 CO₂ concentrations could rise to more than three times preindustrial levels by the end of this
 18 century (Intergovernmental Panel on Climate Change 2007).



19 Source: Intergovernmental Panel on Climate Change 2007:115

20 **Figure 0.C-1. The Greenhouse Effect**

21

2.C.1.3 What Global Climate Changes Have Been Observed?

Key evidence of long-term changes in climate over the twentieth century includes the following documented changes (Intergovernmental Panel on Climate Change 2007, 2012; National Research Council 2010).

- An increase of 1.3 degrees Fahrenheit (°F) in the Earth's global average surface temperature.
- An increase of 6.7 inches in the global average sea level.
- A decrease in arctic sea-ice cover at a rate of approximately 4.1% per decade since 1979, with faster decreases of 7.4% per decade in summer.
- Decreases in the extent and volume of mountain glaciers and snow cover.
- A shift of cold-dependent habitats to higher altitudes and latitudes.
- Longer growing seasons.
- A decrease in the number of unusually cold days and nights and in periods of extreme cold weather.
- More frequent weather extremes such as droughts, floods, severe storms, and heat waves.

2.C.1.4 How is Global Climate Expected to Change?

Global circulation models (GCMs) are used to project future climate change based on assumptions about future emissions of GHGs. The Intergovernmental Panel on Climate Change Special Report on Emissions Scenarios (SRES) presented a range of possible future GHG emission scenarios based on assumptions about potential future fossil fuel use, regional political and social conditions, technologies, population, and governance (Intergovernmental Panel on Climate Change 2000). Based on a number of the SRES emissions scenarios, the Intergovernmental Panel on Climate Change projected an average increase in surface temperatures of 3.2 to 7.2°F by 2100 compared to 1980 through 1999 levels, with a likely increase in the range of 2.0 to 11.5°F when accounting for the uncertainty in climate science (Intergovernmental Panel on Climate Change 2007). Moreover, temperatures are rising at an increasing rate. The average rate of increase over the past century was 0.13°F per decade. Over the past 50 years, temperatures have been rising at nearly twice that average rate, reaching 0.23°F per decade (Intergovernmental Panel on Climate Change 2007). Over the past 30 years, the average global temperature has risen even faster, at an average of 0.29°F per decade (National Oceanic and Atmospheric Administration 2009).

Similar to the global trend, the United States average temperature is now 1.25°F warmer than at the beginning of the 20th century, with an average warming of 0.13°F per decade from 1895 through 2008. This rate of warming is increasing, and over the past decade, the average temperature for the contiguous United States has increased at a rate of 0.58°F per decade (National Oceanic and Atmospheric Administration 2009).

It is noteworthy that even if atmospheric CO₂ concentrations were stabilized or decreased immediately, the current elevated global average temperatures could persist for hundreds of years. This is because temperatures will not drop until the ocean has reached thermal equilibrium with the atmosphere, which would likely take centuries due to the heat capacity of the ocean (Matthews and Caldeira 2008). One study found that up to two-thirds of the maximum increase in global average temperature could persist for centuries (Eby et al. 2009).

1 **2.C.2 Climate Change in California and the Plan Area**

2 **2.C.2.1 Current Climate**


3 The climate in the Sacramento–San Joaquin Delta (Delta) region is generally characterized as
4 Mediterranean (Köppen climate classification), meaning that it has hot, dry summers and cool, rainy
5 winters (Bureau of Reclamation 2011). From 1981 to 2010, average monthly temperatures in
6 Sacramento ranged from 41.0°F in December and January to 94.1°F in July, with average monthly
7 rainfall ranging from a low of 0.02 inches in July to a high of 3.90 inches in February (Western
8 Regional Climate Center 2012).

9 Heat waves are common in summer months, during which temperatures can reach triple digits on
10 consecutive days. Average air temperatures in the mountainous regions of the watershed are
11 typically 5 to 10 degrees lower than temperatures on the valley floor. Periodically, a “Delta breeze”
12 of cool and humid air from the ocean moves onshore and cools the Central Valley in the vicinity of
13 the Delta by up to 7°F (Pierce and Gaushell 2005).

14 **2.C.2.2 Surface Temperature**

15 **2.C.2.2.1 Observed**

16 Figure 2.C-2 shows historical California statewide mean annual temperature departures during the
17 20th century, indicating the warming that has occurred in recent decades. During the 20th century,
18 warming increased by about 3°F over the Sacramento and San Joaquin River basins. (Bureau of
19 Reclamation 2011).




Source: Western Regional Climate Center 2009

Figure 2.C-2. Historical Observed California Statewide Mean Annual Temperature Departure

2.C.2.2.2 Projected

Six GCMs and two of the Intergovernmental Panel on Climate Change SRES emissions scenarios were selected by the California Climate Action Team for their Scenarios Project, resulting in a total of 12 climate change model simulations (Cayan et al. 2009). The scenarios included a high emissions scenario (SRES A2) and a low emissions scenario (SRES B1), Model simulations indicated a midcentury temperature increase in California of about 1.8 to 5.4°F and an end-of-century increase from 3.6 to 9°F, although the warming will not be uniform across California. All climate models projected a greater amount of warming during summer months, during nighttime, and in the interior regions of California (Cayan et al. 2009).

The average mean annual temperature in the Sacramento–San Joaquin basin is projected to increase by 5 to 6°F during this century, though with substantial variability in warming in the Central Valley (Bureau of Reclamation 2011). In addition, the duration of extreme warm temperatures is expected to increase from 2 months (July and August) to four months (June through September) (Climate Action Team 2010). In the early part of the 21st century, the amount of warming in the Sacramento region produced by the SRES A2 high emissions scenario is not very different from the SRES B1 low emissions scenario, but warming becomes increasingly greater through the middle and especially the latter part of the century (Figure 2.C-3).



Source: Cayan et al. 2009

Figure 2.C-3. Simulated Historical and Future Annual Temperature Projections for the Sacramento Region

2.C.2.3 Water Temperature

Cloern et al. (2011) linked a series of models to evaluate the entire San Francisco estuary-watershed system under two contrasting emissions scenarios. Projections for the period 2010 to 2099 indicated significant increases in water temperatures in both the Sacramento River and the Delta under both the B1 (low) and A2 (high) emissions scenarios. Temperatures in the Sacramento River responded to both increasing air temperature and decreasing snowmelt runoff, which reduce the amount of cold water in the upstream reservoirs that is available to cool waters downstream. By contrast, the temperature of Delta waters responded primarily to air temperature. The authors observed that the projected increases in water temperatures, if realized, would have serious implications for the Delta's native fishes. The frequency of projected water temperatures in the Delta above the thermal threshold for delta smelt (25°C) rose gradually under the B1 scenario but rapidly under the A2 scenario. Water temperatures above the threshold for Chinook salmon (16°C) increased at a modest rate under the B1 scenario, but river temperatures above the threshold were common after 2080 under the A2 scenario.

1 **2.C.2.4 Precipitation**

2 **2.C.2.4.1 Observed**


3 On average, total annual precipitation across the United States increased approximately 6% from
4 1901 to 2005 (Intergovernmental Panel on Climate Change 2007). In California, precipitation is
5 extremely variable at seasonal, annual, and decadal time scales (Figure 2.C-4).

6 In the Sacramento River watershed, average annual precipitation ranges from 80 to 90 inches,
7 primarily as snow in the mountainous regions, to 41 and 19 inches of rain in Redding and in
8 Sacramento, respectively. Average annual precipitation for the entire watershed is approximately 36
9 inches. Most precipitation occurs between November and April, with little or no precipitation falling
10 between May and October. The primary origin of precipitation is the seasonal arrival of low-
11 pressure systems from the Pacific Ocean (CALFED Bay-Delta Program 2000).

12 The warming trend in the Sacramento River basin has been accompanied by a gradual trend toward
13 increasing precipitation, starting in the 1930s. However, a similar precipitation trend is not evident
14 in the San Joaquin River basin (Bureau of Reclamation 2011).

15 **2.C.2.4.2 Projected**


16 Projections of future precipitation are more uncertain than those of temperature (Chung et al.
17 2009). While it is difficult to discern strong trends from the full range of climate projections, one
18 recent analysis generally indicated a drying trend in California during the 21st century (Cayan et al.
19 2009). According to this analysis, some areas in northern California may experience higher annual
20 rainfall amounts and potentially larger storm events, but California as a whole, particularly southern
21 California, will be 15 to 35% drier by 2100 (Cayan et al. 2009). Simulated future changes in
22 precipitation for the Sacramento region under the Intergovernmental Panel on Climate Change A2
23 (high) and B1 (low) emissions scenarios indicate declines in the region's precipitation during the
24 21st century (Figure 2.C-5), particularly during the second half of the century (Cayan et al. 2009;
25 Bureau of Reclamation 2011).



1
2
3

Source: Western Regional Climate Center 2009.

Figure 2.C-4. Historical Observed California Statewide Water Year Precipitation



Source: Cayan et al. 2009

Figure 2.C-5. Simulated Future Water Year Change in Precipitation for the Intergovernmental Panel on Climate Change A2 and B1 Scenarios for the Sacramento Region

2.C.2.5 Snowpack and Runoff

2.C.2.5.1 Observed

Historically, most of California’s precipitation fell during winter as snow along the Sierra Nevada mountain range, which serves as a natural reservoir holding the fallen snow. However, temperatures over the Sierra Nevada have increased during the past 100 years, resulting in less snowfall (and more rainfall) and an earlier snowmelt. As a result, there is less snowmelt to sustain runoff during the warm, dry conditions of summer and fall (Knowles et al. 2006; Cayan et al. 2009; Moser et al. 2009).

1 **2.C.2.5.2 Projected**

2 Climate models project that snowpack will continue to decrease in California, with more
3 precipitation falling as rain (Moser et al. 2009). As a result, cool-season runoff will increase during
4 the 21st century in the Sacramento River and San Joaquin River basins, while warm-season runoff is
5 projected to show significant declines (Bureau of Reclamation 2011).

6 Hydrologic model simulations indicate that even if mean precipitation remains unchanged, there
7 will be large impacts on snowpack accumulation, runoff, and soil moisture, with a significant change
8 in the timing and magnitude of flows in the tributary rivers of the San Francisco Bay and
9 Sacramento–San Joaquin River Delta (Bay-Delta) (Bureau of Reclamation 2011). Modeling based on
10 a high emissions scenario indicated a 22% decline in flows to the Sacramento Valley during this
11 century as a result of reduced snow accumulation (Medellín-Azuara et al. 2008).

12 An increase in flows in winter is likely to result in increased flooding, while the decline in spring
13 snowmelt will reduce late summer flows (Cayan et al. 2009). These changes have important
14 implications for California’s water supply system, which is dependent on snowpack storage in the
15 Sierra Nevada. The average early spring snowpack decreased by about 10% during the last century,
16 resulting in a loss of 1.5 million acre-feet of snowpack storage (California Department of Water
17 Resources 2008). Snowpack volumes are expected to decline 25% by 2050 (California Department
18 of Water Resources 2010). Results of modeling by Rauscher et al. (2008) indicate that runoff could
19 occur as much as 2 months earlier than at present.

20 **2.C.2.6 Sea Level Rise**

21 **2.C.2.6.1 Observed**


22 Global average sea level has been gradually rising, increasing by about 6.7 inches during the 20th
23 century (Intergovernmental Panel on Climate Change 2007). Consistent with the global trend, sea
24 level measured at tide gages along the California coast over the past several decades has risen at rate
25 of about 6.7 to 7.9 inches per century (Cayan et al. 2009). While there is considerable variability
26 among the gages, primarily reflecting local differences in vertical movement of the land and length
27 of gage record, this observed rate of change in mean sea level is similar to the global mean trend
28 (National Oceanic and Atmospheric Administration 2009).

29 Sea level has been measured at a tide gage near the Golden Gate since 1897. Based on monthly mean
30 sea-level data from this gage, the mean sea-level increase for San Francisco between 1897 and 2006
31 was 8.0 inches per century (National Oceanic and Atmospheric Administration 2009).

32 **2.C.2.6.2 Projected**

33 Global and regional sea levels have been increasing steadily over the past century, and are projected
34 to increase at a more rapid rate in the future because of the increased thermal expansion of water in
35 the oceans caused by global warming, changes in freshwater inputs to the oceans from melting
36 glaciers and ice sheets, and changes in water storage on land. In 2007, the Intergovernmental Panel
37 on Climate Change estimated a rise in sea level of 0.6 to 1.9 feet by 2100 (Intergovernmental Panel
38 on Climate Change 2007). However, more recent estimates suggest an even greater rise, particularly
39 if melting of the Greenland and Antarctic ice sheets accelerates, as suggested by recent satellite
40 observations. Rahmstorf (2007) used a semi-empirical approach to project future sea level rise,


1 yielding a projected sea level rise of 1.6 to 4.6 feet above 1990 levels by 2100 when applying the
 2 Third Assessment Report warming scenarios. Other recent estimates indicate global increases by
 3 2100 of 1.6 to 3.3 feet (National Research Council 2010), 2.6 to 6.6 feet (Pfeffer et al. 2008) and 3.2
 4 to 5.1 feet (Vermeer and Rahmstorf 2009) (Figure 2.C-6 and Figure 2.C-7).



Source: National Oceanic and Atmospheric Administration 2009

Figure 2.C-6. Observed Mean Sea Level Trend for the San Francisco Tide Gage near the Golden Gate

5
6
7
8
9




Source: Rahmstorf 2007

Figure 2.C-7. Past Global Mean Sea Level and Future Mean Sea Level Based on Global Mean Temperature Projections

Using the Rahmstorf (2007) method, the CALFED Bay-Delta Program (CALFED) Independent Science Board estimated ranges of sea level rise of 2.3 to 3.3 feet at midcentury and of 1.6 to 4.6 feet by the end of the century (CALFED Independent Science Board 2007). Some tidal gage and satellite data indicate that rates of sea level rise are increasing (Church and White 2006; Beckley et al. 2007). Scenarios modeled by the California Climate Action Team projected sea level rise increases along the California coast of 1.0 to 1.5 feet above 2000 levels by 2050 and 1.8 to 4.6 feet by 2100 (Cayan et al. 2009). However, if California's sea level continues to mirror global trends, increases in sea level during this century could be considerably greater. Increasing sea levels will seriously threaten the integrity of the Delta's levees and conveyance of water supplies through the Delta (Florsheim and Dettinger 2007).

For water planning purposes, the California Department of Water Resources (DWR) estimated sea level rise over the 21st century using the method of Rahmstorf (2007) and 12 climate projections selected by the California Climate Action Team (Chung et al. 2009). The historical 95% confidence interval was extrapolated to estimate the uncertainties in the future projections (Figure 2.C-8). Midcentury sea level rise projections ranged from 0.8 to 1.0 foot, with an uncertainty range spanning 0.5 to 1.2 feet. End-of-century projections ranged from 1.8 to 3.1 feet, with an uncertainty range of 1.0 to 3.9 feet. These estimates are slightly lower than those of Rahmstorf (2007) because DWR used a more limited ensemble of climate projections that did not include the highest projections of temperature increases (Chung et al. 2009).

1 Parker et al. (2011) observed that, in the Bay-Delta, other factors complicate sea level rise
 2 projections, including the Pacific Decadal Oscillation (PDO) and El Niño Southern Oscillation (ENSO)
 3 events. The PDO is characterized by cool or warm phase shifts in North Pacific sea surface
 4 temperatures that commonly persist for 20 to 30 years. Superimposed on the PDO cycles are
 5 smaller-scaled El Niño and La Niña events that persist for about a year. Climatic impacts associated
 6 with La Niña events are similar to those tied to the cool PDO phases, and climate conditions related
 7 to El Niño episodes parallel those of warm PDO phases. Parker et al. (2011) observed that rates of
 8 sea level rise slow during the negative (cool) phase and increase during the positive (warm) phase.
 9 They also noted that fluctuations in sea level rise, when combined with processes such as ENSO
 10 events, may have a greater effect on wetlands than a steady increase.



Source: Chung et al. 2009.

Figure 2.C-8. DWR-Generated Future Sea Level Rise Projections for the Bay Delta Using the Rahmstorf Method and Regionally Downscaled Data

11
12
13
14
15

16 2.C.2.7 Ocean Conditions

17 2.C.2.7.1 Observed

18 Global ocean temperatures are also increasing. Between 1961 and 2003, global ocean temperatures
 19 increased about 0.18°F from the surface to a depth of about 2,270 feet (Intergovernmental Panel on
 20 Climate Change 2007). In summer 2009, sea surface temperatures were 1.04°F above the average

1 global temperature recorded for the entire 20th century (Hoegh-Guldberg and Bruno 2010), and the
2 global ocean surface temperature for January 2010 was the second warmest January on record.

3 Another consequence of rising atmospheric concentrations of CO₂ is an increase in the acidity of the
4 oceans as a result of increasing absorption of CO₂ by ocean waters. Recent evidence suggests that
5 oceans have become 30% more acidic since the Industrial Revolution, with severe consequences for
6 marine life (Doney et al. 2009).

7 In the past, ocean productivity has generally adjusted to natural variations in ocean climate.
8 However, present climate trends are outside the bounds of historical variability, and changes in
9 productivity are increasingly apparent. Three factors are likely to affect ocean productivity in
10 response to climate change: temperature, light (as determined by ice cover, cloudiness, and mixed
11 layer thickness), and nutrient availability, with warming temperatures potentially the largest single
12 factor affecting productivity. In the northern high latitudes, reductions in productivity correspond in
13 part to increases in sea surface temperatures. The low latitudes have generally experienced an
14 increase in ocean primary productivity (Brander 2010).

15 **2.C.2.7.2 Projected**

16 Osgood (2008) identified five major climate change effects on the California system of currents.

- 17 • Increased variability in climate forcing.
- 18 • Changes to the magnitude and timing of freshwater inputs.
- 19 • Changes in the timing and strength of the spring transition.
- 20 • Ocean warming and increased stratification.
- 21 • Changes in ocean circulation.

22 More extreme precipitation events and years will affect coastal circulation and stratification. Climate
23 models project greater annual precipitation for northern California but lower streamflow in the
24 21st century, with a more rapid spring melt leading to a shorter, more intense spring period of river
25 flow and freshwater discharge. This will greatly alter coastal stratification and mixing, and riverine
26 plume formation and evolution.

27 Although upwelling along the California coast has increased over the past 30 years and these
28 increases are expected to continue (Snyder et al. 2003), greater thermal stratification and a
29 deepening of the thermocline could reduce upwelling, decreasing nearshore productivity
30 (Roemmich and McGowan 1995). The vertical gradient in ocean temperature off California has
31 intensified over the past several decades (Palacios et al. 2004). Regional climate models suggest that
32 winds favorable for upwelling will be stronger in summer, but the peak in seasonal upwelling will
33 occur later. If upwelling strengthens as a result of global climate change, cold-water species will be
34 favored in the coastal upwelling zones (Snyder et al. 2003).

35 The frequency of large coastal storms and heavy precipitation events is not projected to change
36 significantly over this century (Dettinger et al. 2009). However, the storms that do occur are likely to
37 have significant effects. Increases in the duration of high storm-forced sea levels increases the
38 likelihood that storms will occur during high tides, and higher sea levels combined with severe
39 winter storms and high tides will result in higher storm surges and more severe coastal erosion
40 (Cayan et al. 2009).

1 **2.C.2.8 Extreme Weather Events**

2 **2.C.2.8.1 Observed**

3 Extreme temperatures have changed significantly throughout the world over the past 50 years. Hot
4 days, hot nights, and heat waves have become more frequent; cold days, cold nights, and frost have
5 become less frequent (Intergovernmental Panel on Climate Change 2007). Since 1950, the frequency
6 of heat waves experienced in the United States has increased, although in many regions the heat
7 waves recorded in the 1930s remain the most severe on record. Also, fewer unusually cold days
8 occurred globally in the past few decades, with fewer severe cold waves for the most recent 10-year
9 period in the record (Intergovernmental Panel on Climate Change 2012). Daytime and nighttime
10 heat wave events throughout California have increased in intensity, particularly nighttime events
11 (Moser et al. 2009).

12 Heavy precipitation events have increased in the United States, primarily during the last 3 decades
13 of the 20th century, and mainly over eastern regions (U.S. Global Change Research Program 2009). A
14 recent analysis found that 8 of the top 10 years with extreme 1-day precipitation events occurred
15 between 1990 and 2010 (U.S. Environmental Protection Agency 2010).

16 Globally, longer, more intense droughts have occurred since the 1970s in some regions as a result of
17 higher temperatures and decreased precipitation, particularly in the tropics and subtropics. Changes
18 in sea surface temperatures, wind patterns, and decreased snowpack and snow cover have also been
19 linked to droughts (Intergovernmental Panel on Climate Change 2007). It is estimated that from
20 2001 through 2009, 30 to 60% of the land area in the United States experienced drought conditions
21 at any given time (U.S. Environmental Protection Agency 2010).

22 **2.C.2.8.2 Projected**

23 Significant increases in the frequency and magnitude of high temperature extremes are expected in
24 many parts of California. Under the B1 (low) emissions scenario, there is a projected ten-fold
25 increase in extreme temperatures that currently occur only once every 100 years. Under the A2
26 (high) emissions scenario, these extremes could occur every year (Mastrandrea et al. 2009).

27 Projections of precipitation extremes are highly variable across the state, depending on the model
28 and downscaling method used. However, in general, projections indicate that longer dry spells will
29 become more common, punctuated by occasional intense rainfall events (Mastrandrea et al. 2009).

30 Using a regional climate model, Bell et al. (2004) projected that there will be important changes in in
31 some extreme events in the Delta region. The frequency of extremely hot days (exceeding the long-
32 term 95th percentile value and prolonged (7-day) hot spells are projected to increase in both the
33 San Joaquin River Basin and the Sacramento River Basin.

Change Per Year	San Joaquin River Basin ^a	Sacramento River Basin ^a
Increased No. Hot Days	11.2	32.1
Increased No. Hot Events	1.6	1.5
Increased Days Frost-Free Season	40.1	30.8
Decreased Days Extreme Cold	- 34.9	- 35.5
Decreased Days Below Freezing	- 36	- 47.4
Decreased No. of Prolonged (7-day) Cold Spells	-1.2	-1.8
Decreased Days of Cold Spells	-3.2	-0.5

^a as defined by DWR

1

2 The frost-free growing season was projected to begin earlier and last longer. The models projected
3 fewer days of extreme cold and fewer days below freezing. Prolonged (7-day) cold spells and the
4 duration of cold spells also were projected to decrease.

5 2.C.3 References Cited

- 6 Arndt, D. S., M. O. Baringer, and M. R. Johnson (eds.). 2010. State of the Climate in 2009. *Bulletin of*
7 *the American Meteorological Society* 91:s1–s224.
- 8 Beckley, B. D., F. G. Lemoine, S. B. Luthcke, R. D. Ray, and N. P. Zelensky. 2007. A Reassessment of
9 Global and Regional Mean Sea Level Trends from TOPEX and Jason-1 Altimetry Based on
10 Revised Reference Frame and Orbits. *Geophysical Research Letters* 34:L14608.
- 11 Bell, J. L., L. C. Sloan, and M. A. Snyder. 2004. Regional Changes in Extreme Climatic Events: A Future
12 Climate Scenario. *Journal of Climate* 17:81–87.
- 13 Brander, K. 2010. Impacts of Climate Change on Fisheries. *Journal of Marine Systems* 79:389–402.
- 14 Bureau of Reclamation. 2011. SECURE Water Act Section 9503(c) – Reclamation Climate Change and
15 Water 2011. Available:
16 <<http://www.usbr.gov/climate/SECURE/docs/SECUREWaterReport.pdf>>.
- 17 CALFED Bay-Delta Program. 2000. *Final Programmatic Environmental Impact Statement/*
18 *Environmental Impact Report*. Prepared for the Bureau of Reclamation, U.S. Fish and Wildlife
19 Service, National Marine Fisheries Service, U.S. Environmental Protection Agency, Natural
20 Resources Conservation Service, U.S. Army Corps of Engineers, and California Resources Agency.
21 8 July. Sacramento, CA. State Clearinghouse # 96032083.
- 22 CALFED Independent Science Board. 2007. *Projections of Sea Level Rise for the Delta*. A memo from
23 Mike Healey, CALFED lead scientist, to John Kirlin, Executive Director of the Delta Blue Ribbon
24 Task Force, September 6, 2007. Available:
25 <http://deltavision.ca.gov/BlueRibbonTaskForce/Sept2007/Handouts/Item_9.pdf>.
- 26 California Department of Water Resources. 2008. *Managing an Uncertain Future. Climate Change*
27 *Adaptation Strategies for California's Water*. October.
- 28 California Department of Water Resources. 2010. *Climate Change Characterization and Analysis in*
29 *California Water Resources Planning Studies*. Final Report.

- 1 Cayan, D., M. Tyree, M. Dettinger, H. Hidalgo, T. Das, E. Maurer, P. Bromirski, N. Graham, and R. Flick.
2 2009. *Climate Change Scenarios and Sea Level Rise Estimates for the California 2008 Climate*
3 *Change Scenarios Assessment*. Prepared by the California Climate Change Center for the
4 California Energy Commission. CEC-500-2009-014-D. Sacramento, CA.
- 5 Chung, F., J. Anderson, S. Arora, M. Ejeta, J. Galef, T. Kadir, K. Kao, A. Olson, C. Quan, E. Reyes, M. Roos,
6 S. Seneviratne, J. Wang, and H. Yin. 2009. *Using Future Climate Projections to Support Water*
7 *Resources Decision Making in California*. California Energy Commission. CEC-500-2009-052-F.
- 8 Church, J., and N. White. 2006. A 20th Century Acceleration in Global Sea-Level Rise. *Geophysical*
9 *Research Letter* 33:L01602.
- 10 Climate Action Team. 2010. *Climate Action Team Report to Governor Schwarzenegger and the*
11 *California Legislature*. California Environmental Protection Agency. April 2010. Available:
12 <http://www.climatechange.ca.gov/climate_action_team/reports/index.html>.
- 13 Cloern, J. E., N. Knowles, L. R. Brown, D. Cayan, M. D. Dettinger, T. L. Morgan, D. H. Schoellhamer, M. T.
14 Stacey, M. van der Wegen, R. W. Wagner, and A. D. Jassby. 2011. *Projected Evolution of*
15 *California's San Francisco Bay-Delta-River System in a Century of Climate Change*. Available:
16 <<http://www.plosone.org/article/info%3Adoi%2F10.1371%2Fjournal.pone.0024465>>.
- 17 Dettinger, M. D., H. G. Hidalgo, T. Das., D. R. Cayan, and N. Knowles. 2009. *Projections of Potential*
18 *flood Regime Changes in California*. California Energy Commission, PIER Energy-Related
19 Environmental Research. CEC-500-2009-050.
- 20 Doney, S. C., V. J. Fabry, R. A. Feely, and J. A. Kleypas. 2009. Ocean Acidification: The Other CO₂
21 Problem. *Annual Review of Marine Science* 1:169–192.
- 22 Eby, M., K. Zickfeld, A. Montenegro, D. Archer, K. J. Meissner, and A. J. Weaver. 2009. Lifetime of
23 Anthropogenic Climate Change: Millennial Time Scales of Potential CO₂ and Surface
24 Temperature Perturbations. *Journal of Climate* 22(10):2501–2511.
- 25 Florsheim, J. L. and M. D. Dettinger. 2007. Climate and Floods Still Govern California Levee Breaks.
26 *Geophysical Research Letters* 34:L22403.
- 27 Hoegh-Guldberg, O. and J. F. Bruno. 2010. The Impact of Climate Change on the World's Marine
28 Ecosystems. *Science* 328:1523–1528.
- 29 Intergovernmental Panel on Climate Change. 2000. *IPCC Special Report on Emissions Scenarios*. N.
30 Nakicenovic and R. Swart (eds.). Cambridge University Press, UK. Available:
31 <<http://www.grida.no/climate/ipcc/emission/>>.
- 32 Intergovernmental Panel on Climate Change. 2007. *Climate Change 2007: The Physical Science Basis.*
33 *Contribution of Working Group I to the Fourth Assessment Report of the Intergovernmental Panel*
34 *on Climate Change*. S. Solomon S., D. Qin, M. Manning, Z. Chen, M. Marquis, K. B. Averyt, M. Tignor
35 and H. L. Miller (eds.). Cambridge University Press: Cambridge, UK, and New York, NY. 996 pgs.
36 Available: <http://www.ipcc.ch/publications_and_data/ar4/wg1/en/contents.html>.

- 1 Intergovernmental Panel on Climate Change. 2012. *Managing the Risks of Extreme Events and*
2 *Disasters to Advance Climate Change Adaptation. A Special Report of Working Groups I and II of*
3 *the Intergovernmental Panel on Climate Change* [Field, C. B., V. Barros, T. F. Stocker, D. Qin, D. J.
4 Dokken, K. L. Ebi, M. D. Mastrandrea, K. J. Mach, G.-K. Plattner, S. K. Allen, M. Tignor, and P.M.
5 Midgley (eds.)]. Cambridge University Press, Cambridge, UK, and New York, NY, USA, 582 pp.
- 6 Knowles, N., M. Dettinger, and D. Cayan. 2006. Trends in Snowfall versus Rainfall for the Western
7 United States. *Journal of Climate* 19 (18):4545–4559.
- 8 Mastrandrea, M. D., C. Tebaldi, C. P. Snyder, and S. H. Schneider. 2009. *Current and Future Impacts of*
9 *Extreme Events in California*. A paper from California Climate Change Center. California Energy
10 Commission Public Interest Energy Research Program. CEC-500-2009-026-F. Sacramento, CA.
- 11 Matthews, H. D., and K. Caldeira. 2008. Stabilizing Climate Requires Near-Zero Emissions.
12 *Geophysical Research Letters* 35(4):L04705.
- 13 Medellín-Azuara, J., J. J. Harou, M. A. Olivares, K. Madani, J. R. Lund, R. E. Howitt, S. K. Tanaka, M. W.
14 Jenkins, and T. Zhu. 2008. Adaptability and Adaptations of California’s Water Supply System to
15 Dry Climate Warming. *Climatic Change* 87(Suppl 1): S75–S90.
- 16 Moser, S., G. Franco, S. Pittiglio, W. Chou, and D. Cayan. 2009. *The Future Is Now: An Update on*
17 *Climate Change Science Impacts and Response Options for California* (Special Report California
18 Climate Change Center). Prepared for the California Energy Commission. CEC-500-2008-071.
19 Sacramento, CA.
- 20 National Climatic Data Center. 2011. *Global Surface Temperature Anomalies. Annual Global (land and*
21 *ocean combined) Anomalies*. Available: <[http://www.ncdc.noaa.gov/cmb-](http://www.ncdc.noaa.gov/cmb-faq/anomalies.php#anomalies)
22 [faq/anomalies.php#anomalies](http://www.ncdc.noaa.gov/cmb-faq/anomalies.php#anomalies)>.
- 23 National Oceanic and Atmospheric Administration. 2009. *State of the Climate: Global Analysis for*
24 *Annual 2008*. NOAA National Climatic Data Center. Available:
25 <<http://www.ncdc.noaa.gov/sotc/global/2008/13>>.
- 26 National Research Council. 2010. *Advancing the Science of Climate Change. America’s Climate Choices:*
27 *Panel on Advancing the Science of Climate Change*. Board on Atmospheric Sciences and Climate,
28 Division of Earth and Life Sciences. National Academies Press, Washington, DC. 504 pp.
29 Available: <http://www.nap.edu/catalog.php?record_id=12782>.
- 30 Osgood, K. E. (ed.) 2008. *Climate Impacts on U.S. Living Marine Resources: National Marine Fisheries*
31 *Service Concerns, Activities and Needs*. U.S. Department of Commerce, National Oceanic and
32 Atmospheric Administration, National Marine Fisheries Service, NOAA Technical Memorandum
33 NMFS-F/SPO-89. August 2008.
- 34 Palacios, D. M, S. J. Bograd, R. Mendelssohn, and F. B. Schwing. 2004. Long-Term and Seasonal
35 Trends in Stratification in the California Current, 1950–1993. *Journal of Geophysical Research –*
36 *Oceans* 109 (C10):C10016.
- 37 Pfeffer, W. T., J. T. Harper, and S. O’Neel. 2008. Kinematic Constraints on Glacier Contributions to
38 21st Century Sea-level Rise. *Science* 321:1340–1343.

- 1 Pierce, D. W. and D. Gaushell. 2005. *The California Delta Breeze: Characteristics and Sub-daily*
2 *Forecasting*. Abstract from the Sixth Conference on Coastal Atmospheric and Oceanic Prediction
3 and Processes, San Diego, CA.
- 4 Rahmstorf, S. 2007. A Semi-Empirical Approach to Projecting Future Sea-Level Rise. *Science*
5 315:368–370.
- 6 Rauscher, S. A., Pal, J. S., Diffenbaugh, N. S., and Benedetti, M. M. 2008. Future Changes in Snowmelt-
7 Driven Runoff Timing over the Western US. *Geophysical Research Letters* 35: Article No. L16703.
- 8 Roemmich, D., and J. McGowan. 1995. Climatic Warming and the Decline of Zooplankton in the
9 California Current. *Science* 267:1324–1326.
- 10 Snyder, M. A., L. C. Sloan, N. S. Diffenbaugh, and J. L. Bell. 2003. Future Climate Change and Upwelling
11 in the California Current. *Geophysical Research Letters* 30: Article No. 1823.
- 12 U.S. Environmental Protection Agency. 2010. *Climate Change Indicators in the United States*.
13 Washington, DC. Available:
14 <http://www.epa.gov/climatechange/indicators/pdfs/ClimateIndicators_full.pdf>.
- 15 U.S. Global Change Research Program. 2009. *Global Climate Change Impacts in the United States*. T. R.
16 Karl, J. M. Melillo and T. C. Peterson (eds.). Cambridge University Press: Cambridge, UK and New
17 York, NY. 196 pgs. Available: <[http://downloads.globalchange.gov/usimpacts/pdfs/climate-](http://downloads.globalchange.gov/usimpacts/pdfs/climate-impacts-report.pdf)
18 [impacts-report.pdf](http://downloads.globalchange.gov/usimpacts/pdfs/climate-impacts-report.pdf)>.
- 19 Vermeer M. and S. Rahmstorf. 2009. Global Sea Level Linked to Global Temperature. *Proceedings of*
20 *the National Academy of Sciences of the United States of America* 106:21527–21532.
- 21 Western Regional Climate Center. 2009. California Statewide Mean Temperature Departure (Jan-
22 Dec) to Date. Available: <http://www.wrcc.dri.edu/monitor/cal-mon/frames_version.html>.
- 23 Western Regional Climate Center. 2012. Climate Summary. Available:
24 <<http://www.wrcc.dri.edu/cgi-bin/cliMAIN.pl?ca7633>>. Accessed: June 5, 2012.