

**Long-Term Trends in
Benthos Abundance and Persistence
in the
Upper Sacramento-San Joaquin Estuary**

Summary Report: 1980-1990

Zachary Hymanson
California Coastal Commission

David Mayer, Ph.D.
TENERA Environmental Services

John Steinbeck
TENERA Environmental Services

Technical Report 38

May 1994

Interagency Ecological Program
for the
San Francisco Bay/Delta Estuary

A Cooperative Program of:

California Department of Water Resources
State Water Resources Control Board
U.S. Bureau of Reclamation
U.S. Army Corps of Engineers

California Department of Fish and Game
U.S. Fish and Wildlife Service
U.S. Geological Survey
U.S. Environmental Protection Agency

Contents

	CONCLUSIONS	vii
	RECOMMENDATIONS	ix
	ACKNOWLEDGMENTS	xi
Chapter 1	INTRODUCTION	1
	Program Origin	1
	Program Description	2
	Environmental Setting	5
Chapter 2	SURVEY AND ANALYSIS METHODS	9
	Benthic Organisms	9
	Sediment	10
	Water Quality	11
	Phytoplankton	11
	Zooplankton	13
Chapter 3	TRENDS IN FACTORS INFLUENCING THE BENTHOS	15
	Freshwater Flow	15
	Water Quality	15
	Sediments	17
	Food Supply	22
	Volatile Suspended Solids	22
	Phytoplankton	23
	Zooplankton	26
Chapter 4	CHANGES IN THE BENTHOS	29
	Grizzly Bay Site, D7-C	30
	Sacramento River Sites, D4-R, D4-L, D4-C	33
	Eastern Sites, D11-C, D19-C, D28A-L, D28A-R	35
	Persistent and Dominant Species	37
	<i>Limnodrilus hoffmeisteri</i> and <i>Varichaetadrilus angustipenis</i>	37
	<i>Corophium stimpsoni</i> and <i>C. spinicorne</i>	37
	<i>Manayunkia speciosa</i>	38
	<i>Corbicula fluminea</i>	39
	<i>Potamocorbula amurensis</i>	40
	<i>Gammarus daiberi</i>	40
	<i>Hemileucon hinumensis</i>	41
	Trends in the Benthos and Water Project Operations	41
	Detectability and Sensitivity Analysis of the Benthic Monitoring Program	41
	What should the sampling frequency be?	41
	What should the sample replication be?	42
	What should the spatial distribution of stations be?	45
Bibliography	LITERATURE PERTAINING TO THE BENTHOS OF THE SACRAMENTO-SAN JOAQUIN ESTUARY	47

Appendixes

A	Taxonomic List of Benthic Macrofauna Identified from Monitoring Samples	51
B	Graphs of Monthly Abundance of All Species (Total) and the Four Numerically Dominant Species at Each Benthic Site	57

Tables

1	Sources of Potential Impact to the Integrity of the Upper Estuary and Resources Potentially Affected	1
2	Initial Fate of Benthic Monitoring Stations Originally Listed in Decision 1379	3
3	Stations and Sites of Benthos and Substrate Sampling, 1975–1981	3
4	Characteristics of Five Monthly Benthic Sampling Stations	5
5	Benthic and Substrate Sampling Stations and Sites	10
6	Summary of Changes in Suisun Bay/Delta Zooplankton Anomalies	26
7	Summary of Regional Changes in Abundance of Zooplankton Taxa that Declined between 1972 and 1987	27
8	Correlation Analysis Results of Various Environmental Constituents versus First Dimension Correspondence Analysis Scores for the Benthic Monitoring Sites	33
9	Coefficients of Variation for Total Community Abundance Values and for <i>Corbicula fluminea</i> Abundance Values Calculated for Different Sites and Time Intervals	42

Figures

1	Benthic and Water Quality Monitoring Stations	4
2	Concentrations of <i>P. amurensis</i> and Chlorophyll <i>a</i> in Grizzly Bay, Site D7-C, 1980–1990	8
3	Stations and Regions Used in Water Quality and Phytoplankton Analyses	12
4	Mean Annual Flow, Sacramento River at Sacramento	15
5	Trends in Water Temperature	16
6	Trend in Annual Salinity, Suisun Bay Region	16
7	Trend in Annual Salinity, Western Delta Region	16
8	Trend in Annual Salinity, Central Delta Region	16
9	Trend in Annual Percentage of Fine Inorganic and Organic Sediments, Grizzly Bay, Station D7	17
10	Trend in Annual Percentage of Fine Inorganic and Organic Sediments, Sacramento River, Station D4	18
11	Trend in Annual Percentage of Fine Inorganic and Organic Sediments, Sherman Lake, Station D11	19


12	Trend in Annual Percentage of Fine Inorganic and Organic Sediments, Franks Tract, Station D19	20
13	Trend in Annual Percentage of Fine Inorganic and Organic Sediments, Old River, Station D28A	21
14	Trend in Volatile Suspended Solids, Suisun Bay Region	23
15	Trend in Volatile Suspended Solids, Western Delta Region	23
16	Trend in Volatile Suspended Solids, Central Delta Region	23
17	Chlorophyll <i>a</i> Concentrations at Various Stations in the Upper Estuary	24
18	Chlorophyll <i>a</i> Concentrations and Anomalies Versus Time	25
19	First Dimension Correspondence Analysis, All Sites	29
20	Second Dimension Correspondence Analysis, All Sites	30
21	First Dimension Correspondence Analysis, Grizzly Bay, Site D7-C	30
22	Second Dimension Correspondence Analysis, Grizzly Bay, Site D7-C	30
23	Species Scores, Grizzly Bay, Site D7-C	31
24	First Dimension Correspondence Analysis, Sacramento River, Station D4	33
25	Second Dimension Correspondence Analysis, Sacramento River, Station D4	33
26	Species Scores, Sacramento River, Station D4	34
27	First Dimension Correspondence Analysis, Eastern Stations, D11, D19, and D28A	35
28	Second Dimension Correspondence Analysis, Eastern Stations, D11, D19, and D28A	35
29	Species Scores, Eastern Stations, D11, D19, and D28A	36
30	<i>Corophium stimpsoni</i> Concentrations and Monthly Specific Conductance at D11-C, 1987-1990	38
31	Power of Detection Curves for Total Community Abundance at D7-C	43
32	Power of Detection Curves for <i>Corbicula fluminea</i> at Site D19-C	44
33	Similarity in Benthic Species Composition Among Sites in Two Delta Regions	45

CONCLUSIONS

A variety of analyses were used to summarize long-term trends in the benthos and relevant physical, chemical, and biological variables of the upper Sacramento-San Joaquin estuary. Results of these analyses were compared and related to determine probable causes for the trends in the benthos. The main conclusions from this effort are:

- The exotic organisms *Potamocorbula amurensis*, *Hemileucon hinumensis*, and *Gammarus daiberi* all became established as numerically dominant organisms at one or more benthic sampling sites in the upper estuary between 1980 and 1990. Establishment and numerical dominance of these exotic species has altered the ecology of the upper Sacramento-San Joaquin estuary.
- The combination of extreme deviations in freshwater flows and salinity along with the invasion of three introduced species resulted in a substantial change in the benthic communities at stations D7 and D4. These communities are now dominated by one or more of the recently introduced species.
- The more eastern stations (D11, D19, D28A) were also affected by the physicochemical changes and establishment of exotic species, although to a lesser degree than D7 and D4. The benthic communities at these eastern stations were apparently able to absorb these changes, since the communities did not show a substantial change in persistence of resident species.
- From 1980 through 1990, there was a general increase in the amount of fine sediment at many of the sampling locations as a result of reduced streamflow. In general, however, there was no connection between trends in sediment composition and the abundance and persistence of benthic organisms.
- From 1978 through 1990, concentrations of volatile suspended solids showed no significant trend with time, although there was a substantial decline in Suisun Bay. Phytoplankton and zooplankton biomass showed significantly negative trends in many parts of the upper estuary. In Suisun Bay beginning in 1986, Alpine and Cloern (1992) found a sustained and substantial decline in phytoplankton biomass that could be at least partly explained by the invasion of the suspension-feeding Asian clam, *Potamocorbula amurensis*. A similar phenomenon may have occurred in the western Delta.

LOCATIONS OF RECOMMENDED NEW AND RETAINED BENTHIC MONITORING STATIONS



NUMBER, LOCATION, AND RATIONALE FOR FIVE NEW BENTHIC MONITORING STATIONS

Station Number	Station Location	Rationale for Selection
D41A	Light 2, Mouth of Petaluma River, San Pablo Bay	Only site in San Pablo Bay. USGS and DWR have sampled site since 1988. Routine sampling could provide information relating benthos in San Pablo Bay to benthos in Suisun Bay.
D6	Ship channel in Suisun Bay near Martinez	Provides better spatial coverage of Suisun Bay. Samples a habitat different from site D7 in the Grizzly Bay shoal area.
D24	Sacramento River below Rio Vista Bridge	Provides better characterization of lower Sacramento River area.
D16	San Joaquin River at Twitchell Island	Provides information on the benthos of the lower San Joaquin River.
C9	West Canal opposite Intake Channel to Clifton Court Forebay	Permits better spatial coverage for assessment of potential water project related impacts to the benthos.

RECOMMENDATIONS

The premise of these recommendations is that a benthic monitoring program is needed in the upper Sacramento-San Joaquin estuary with the following objectives:

- Monitor trends in the abundance and distribution of benthic fauna.
- Detect major changes in species composition, especially introductions.
- Provide baseline information for special studies.

Given the stated objectives and results from the various analyses, a benthic monitoring program with the following attributes is recommended:

- Benthic and sediment sampling should continue at three existing sites: D7-C, D4-L, and D28A-L. Sampling at the other five existing sites (D11-C, D4-R, D4-C, D19-C, D28AR) could be discontinued. Instead, five new sampling stations should be established. Sampling these stations provides better spatial coverage of the monitoring area and may permit a better understanding of SWP and CVP related impacts in some cases.
- Three replicate samples should be collected from each site on a monthly basis. This level of sampling effort would result in a monitoring program that is consistent with all program objectives. All other sample collection methods could remain the same.
- Organism biomass should be estimated bimonthly at all sites by measuring total wet weight of major taxonomic groups (*eg*, phylum or class) and dominant species. Existing curves relating tissue weight to total weight could be used for organisms, such as clams, with a substantial portion of their total weight arising from nonliving parts. All other sample analysis methods could remain the same. Routine measurements of both biomass and abundance would permit estimates of benthic production. In addition, biomass estimates provide information useful to understanding benthic trophic dynamics.
- A summary analysis and full re-evaluation of the benthic monitoring program should be completed every 5 years. Annual analyses should continue to determine if program adjustments are necessary.

ACKNOWLEDGMENTS

Many people contributed to the completion of this summary review. Benthos and water quality monitoring data were collected by Shig Katsumata, Katherine Triboli, John Baker, and Lloyd Brenn (DWR), and George Collins (USBR). Members of the Interagency Food Chain Group provided technical advice throughout the course of this review. The authors would especially like to thank Wayne Fields (Hydrozoology), Steve Ford (DWR), Bruce Herbold (EPA), Tim Hollibaugh (SFSU), Wim Kimmerer (BioSystems Analysis), Fredric Nichols (USGS), Steve Obrebski (SFSU), Jim Orsi (DFG), Harlan Proctor (DWR), Ted Sommers (DWR), and Bruce Thompson (AHI) for their review and comments on an earlier draft of this report. Richard Wee (TENERA) and Vera Tharp (DWR) completed all technical editing. Funding for data collection and this summary review was provided by DWR and USBR.

INTRODUCTION

The Department of Water Resources began monitoring the soft-bottom benthos of Suisun Bay and the Sacramento-San Joaquin Delta in 1975 as part of a comprehensive environmental monitoring program mandated by the State Water Resources Control Board. The resulting program has produced a comprehensive data set, which is used in a variety of ways including routine examination of benthos abundance and distribution, detection and tracking of introduced organisms, and as baseline information for applied research projects.

This report begins with a review of the benthic monitoring program — its origin, purpose, and design criteria. The study area is then described, including a discussion of trends in relevant physical, chemical, and biological variables. Next, the results of summary analyses for benthic monitoring data collected from 1980 through 1990 are presented. These results are also related to trends in other relevant variables to discover causal relationships. Results of analyses that test the detectability and sensitivity of the monitoring methods are also included.

Program Origin

The benthic monitoring program is one element of a comprehensive baseline monitoring effort required of DWR by the SWRCB through its regulatory authority over California's water rights. The program also includes monitoring of water quality, phytoplankton, and zooplankton. Monitoring and reporting requirements of the program are described in the current Water Right Decision 1485¹.

Water Right Decision 1379² (which preceded Decision 1485) was the first delta water right decision to provide terms and conditions for a comprehensive

monitoring program. As a result of testimony presented during hearings for Decision 1379 and testimony heard in earlier decisions (Decisions 990 and 1275), SWRCB decided a monitoring program was needed to routinely determine water quality conditions, pollutant loads and sources, and changes in environmental conditions within the estuary.

The environmental monitoring program described in Decision 1379 was developed by Stanford Research Institute through a contract with SWRCB³. SWRCB's stated objective was to:

“develop a monitoring program sensitive to important parameters that characterize the environment, and that can provide information necessary for effective management of the water resources of the region”.

The resulting program was truly comprehensive in scope, as it considered a wide variety of impacts to the estuary. The report's inventory of potentially affected resources, combined with the sources of impact (Table 1), formed both the rationale and the basis for the environmental monitoring program.

<u>Sources of Potential Impact</u>	<u>Resources Potentially Affected</u>
Sea Water Contamination	Municipal Water Supply
Pesticide Manufacture & Application	Industrial Water Supply
Irrigation Return Water	Agricultural Water Supply
Domestic & Industrial Waste Water	Fish and Wildlife Propagation & Sustenance
Breakdown Products	Commercial & Sport Fisheries
	Navigation
	Recreation
	Esthetic Values (including historic value)

SOURCE: Weisbecker *et al* 1970.

1 State Water Resources Control Board. 1978. *Water Right Decision 1485 for the Sacramento-San Joaquin Delta and Suisun Marsh*. 44 pp.

2 State Water Resources Control Board. 1969. *Water Right Decision 1379 for the Sacramento-San Joaquin Delta and Suisun Marsh*.

3 LW Weisbecker, JL Mackin, AW Knight, RW Brocksen. 1970. *An Environmental Monitoring Program for the Sacramento-San Joaquin Delta and Suisun Bay*. Stanford Research Institute. Contract 9-2-32. Prepared for State Water Resources Control Board. Publication 40. 106 pp plus appendixes.

In its report, SRI recommended full implementation of a comprehensive environmental monitoring program to ensure that collection and interpretation of environmental information was sufficient for effective management of the estuary. SRI found very few estuary monitoring programs existed at the time of the review. Although numerous applied research projects had been completed, these investigations were conducted by several agencies working without a common objective or plan to investigate or manage the estuary. SWRCB had committed itself to such a monitoring program and used its powers in the water right decision process to implement a program that went beyond measuring changes in the estuary that might be directly related to water project operations.

SRI's review of existing monitoring programs also disclosed a lack of routine monitoring for biological constituents. The authors concluded that:

“as the relationships between physical, chemical, and biological conditions, and environmental effects become better defined, many more resource management actions that are directly concerned with water quality will be based upon environmental parameters. Aquatic organisms do not have the capability of processing or preconditioning water to meet their biological requirements as does man. Therefore, these organisms can be sensitive indicators of environmental change”.

The SRI environmental monitoring program included monitoring benthic species abundance and diversity primarily to detect the effects of wastewater discharges on the estuary. Although such discharges are not part of water project activities, there may be secondary relationships between water project exports and wastewater discharges that could affect the estuary environment. Distinguishing the primary and secondary impacts of water project operations was an important objective of the SRI monitoring program.

Implementation of the program began in 1972, as SWRCB, DWR, and USBR met to define their individual responsibilities for various elements of the monitoring program⁴. Benthic monitoring began in

1975. Benthic sampling frequency of once a month, as recommended by SRI, was modified to quarterly in Decision 1379 and then biannually in Decision 1485.

Program Description

Implementation of the benthic monitoring program in 1975 coincided with numerous other changes in the comprehensive monitoring program. The most notable change was the transfer of overall responsibility for the monitoring program from USBR to DWR. Initially, the majority of DWR's resources were directed toward implementation of the water quality monitoring element, because it was agreed that a program of this magnitude would have to be phased in over a reasonable period⁵. Once the routine of the water quality monitoring element was established, the benthic monitoring element was implemented essentially as described in Decision 1379. However, it was agreed in discussions during program implementation that: (1) samples would be collected biannually rather than quarterly, and (2) the number and location of sampling stations detailed in Decision 1379 would be adjusted. Initially 11 of the 16 benthic stations listed in Decision 1379 were sampled (Table 2).

In 1978, SWRCB released Decision 1485, which (as in Decision 1379) described the benthic monitoring element requirements. Although the basic requirements remained unchanged from Decision 1379, several changes were made in the number and location of sampling stations, as summarized in Table 3. These changes were based on results of data analysis and field experience gained from sampling in previous years. From 1975 through 1979, between 11 and 16 stations were sampled biannually for benthic species composition and abundance and sediment composition. These data on species presence, abundance, and distribution were used to characterize the delta's benthic environment and assess its benthic populations. DWR reported and evaluated these monitoring results in annual summary reports⁶.

4 Harlan Proctor, DWR; personal communication.

5 H. Proctor; personal communication.

6 Department of Water Resources. Annually, 1976 to Present. *Water Quality Conditions in the Sacramento-San Joaquin Delta*. Report to the State Water Resources Control Board in accordance with Water Right Decision 1485 [Decision 1379 until the 1979 report], Order 4(f).

In June 1980, DWR began monthly collections of benthic and sediment samples at five stations in the upper estuary (Figure 1). This change in sampling design was made "to more accurately monitor and evaluate seasonal changes in the composition of the benthic fauna and associated physical factors"⁷. The five stations were selected primarily on the basis of salinity and substrate criteria (Table 4). Monitoring results from the revised program continued to be reported annually. In addition, a summary report was prepared by Markmann⁸, in which she analyzed the benthic data collected from 1975 through 1981.

**Table 2
INITIAL FATE OF BENTHIC MONITORING STATIONS
ORIGINALLY LISTED IN DECISION 1379***

Station	Fate
Big Break off Jersey Island	Sampled, Designated D14A
Carquinez Strait at Martinez	Sampled, Designated D6
Hog Slough	Sampled; Relocated to MD6, Sycamore Slough
Middle River at Victoria	Never Sampled
Mokelumne River, South Fork near Terminous	Sampled, Designated MD7
Old River at Palm Tract	Sampled, Designated D28A
Sacramento River upstream of Confluence of American River	Never Sampled
Sacramento River at Chipps Island	Sampled, Designated D10
Sacramento River at Greens Landing	Sampled, Designated C3
Sacramento River just below Sacramento	Never Sampled
Sacramento River at Threemile Slough	Sampled; Relocated to D24, Sacramento River below Rio Vista Bridge
San Joaquin River at Mossdale	Sampled, Designated C7
San Joaquin River below Stockton	Sampled, Designated P8
San Joaquin River at Threemile Slough	Never Sampled
San Pablo Bay off Hercules in Dredged Channel	Never Sampled
Suisun Bay at Port Chicago	Sampled; Relocated to D8, Suisun Bay off Middle Point near Nichols

* Adapted from Water Right Decision 1379. (SWRCB 1969)

**Table 3
STATIONS AND SITES OF
BENTHOS AND SUBSTRATE SAMPLING,
1975-1981**

Station	Site*	1975	1976	1977	1978	1979	1980	1981
C3	R	B	S	S	S/B	S/B		
	C	B	S/B	S/B	S/B	S/B		
	L	B	S	S	S/B	S/B		
C7	R	S	S/B	S/B	S			
	C	S/B	S/B	S/B	S/B			
	L	S	S/B	S/B	S			
D4	R	S	S	S/B	S/B	S/B	S/B	S/B
	C	S/B	S/B	S/B	S/B	S/B	S/B	S/B
	L	S	S	S/B	S/B	S/B	S/B	S/B
D6	R	S/B	S/B	S/B	S			
	C	S/B	S/B	S/B	S/B			
	L	S/B	S/B	B	S			
D7	R			S/B	S	S		
	C			S/B	S/B	S/B	S/B	S/B
	L			S/B	S	S	S	S
D8	R	S	S					
	C	S/B	S/B					
	L	S	S					
D9	R			B	S/B	S		
	C			S/B	S/B	S/B		
	L					S		
D10	R	S	S/B					
	C	S/B	S/B					
	L	S	S/B					
D11	R		S/B	S/B	S/B	S	S	S
	C		S/B	S/B	S/B	S/B	S/B	S/B
	L		S/B	S/B	S/B	S	S	S
D12	R	S	S					
	C	S/B	S/B	B				
	L	S	S					
D14A	R	S	S/B	S/B	S	S		
	C	S/B	S/B	S/B	S/B	S/B		
	L	S	S/B	S/B	S	S		
D19	R			S	S/B	S	S	S
	C			S/B	S/B	S/B	S/B	S/B
	L			S	S/B	S	S	S
D24	R	S	S	S				
	C	S/B	S/B	S/B				
	L	S	S	S				
D26	R	S	S/B	S				
	C	S/B	S/B	S/B				
	L	S	S/B					
D28A	R	S/B	S	S/B	S/B	S/B	S/B	S/B
	C	S/B	S/B	S/B	S/B	S/B	S/B	S/B
	L	S	S	S/B	S/B	S/B	S/B	S/B
MD6	R	S	S/B	S	S	S		
	C	S/B	S/B	S/B	S/B	S/B		
	L	S	S/B	S	S	S		
MD7	R	S	S	S/B	S/B	S/B		
	C	S/B	S/B	S/B	S/B	S/B		
	L	S	S	S/B	S/B	S/B		
P8	R	S	S	S	S/B	S/B		
	C	S/B	S/B	S/B	S/B	S/B		
	L	S/B	S	S	S/B	S/B		

S = Substrate Collected; B = Benthos Collected
* Facing downstream: R = Right Bank, C = Center, L = Left Bank

7 DWR, Annual Report for 1980, cited.

8 C Markmann. 1986. *Benthic Monitoring in the Sacramento-San Joaquin Delta; Results from 1975 through 1981*. Interagency Ecological Study Program Technical Report 12. Department of Water Resources.




Figure 2
 CONCENTRATIONS OF *P. AMURENSIS* AND CHLOROPHYLL *a* IN GRIZZLY BAY, SITE D7-C, 1980-1990

SURVEY AND ANALYSIS METHODS

The study area spans a variety of habitats from narrow, freshwater channels in the delta to broad, estuarine bays. The Sacramento-San Joaquin estuary is one of the largest estuaries in the United States. It is also one of the nation's most modified estuaries.²⁵ The many recorded changes have affected virtually every aspect of this estuary at one time or another. Changes such as urban development, wetland development, diversion of fresh water, alterations in sediment loadings, species introductions, and weather patterns all have the potential of affecting the benthos. Overall, the ecology of the estuary is primarily dictated by physico-chemical processes; however, biological events such as introduction of exotic organisms have also had pronounced effects on the estuary's ecosystem.

The upper Sacramento-San Joaquin estuary is classified as a partially mixed and tidally dominated estuary. The estuary's hydrology is complicated by regional differences in geography, which strongly influence the system's hydrology. In general, tidal flows greatly exceed freshwater inflows except during periods of high streamflow in wet winters. A mixing zone of fresh water and salt water is always present, although its location is transient. Thus, salinity and water current patterns, which directly affect the distribution and transport of numerous organisms, vary according to local conditions throughout the study area.

Sediment loads and their distribution are additional features of the estuary that can affect the benthos. Water storage and diversion from major tributaries of the estuary have reduced the seasonal magnitude of freshwater inflows and the supply of sediment. Water velocities, bathymetry, and wind and weather patterns also affect sediment resuspension and composition at a given location. Both Suisun and San Pablo bays have extensive shoals. Sediments in

these areas are often resuspended as a result of the winds common to the region and then transported by prevailing water currents.

During the monitoring period discussed in this report (1980–1990), monthly benthic samples were collected consistently from five stations in the upper estuary (Figure 1). These stations were chosen for more intensive sampling from a larger set of stations sampled biannually between 1975 and 1979. The five stations chosen were thought to represent major aquatic environments within the upper estuary.²⁶ Key characteristics and the criteria used to select each station are shown in Table 4 (page 5).

Benthic Organisms

The benthos of the delta and western bays includes a diverse assemblage of organisms that range from single-cell bacteria and ciliates to large crabs and clams. Changes in the benthic macrofauna (those organisms larger than 0.5 mm²⁷) were documented in this monitoring program. For sampling purposes, each station was divided into a maximum of three sectors: right bank (R), left bank (L), and center (C). Thus, a sampling site is identified by the station and sector designations.

All samples were collected using a hydraulic winch and Ponar dredge. The dredge was fitted with screens that allow water to pass through on descent to minimize a bow wave effect on epifauna prior to impact. The Ponar dredge samples a bottom area of about 0.053 m² to a depth that varies with the type of sediment and the ability of the dredge to penetrate it. The number of organisms per square meter was determined by multiplying the count of organisms collected in each sample by 19 (*ie*, 1.0 m²/0.053 m² \cong 19). Three replicate grab samples were collected from eight sites each month (Table 5).

25 FH Nichols, JE Cloern, SN Luoma, DH Peterson. 1986. The modification of an estuary. *Science* 231:525-628.

26 Markmann, 1986; cited.

27 Nichols and Pamatmat, 1988; cited.

Table 5
BENTHIC AND SUBSTRATE
SAMPLING STATIONS AND SITES

Station	Site*	Type of Sample**	Habitat
D4	R	Substrate/Benthos	River Channel
	C	Substrate/Benthos	
	L	Substrate/Benthos	
D7	R	Substrate	Shallow Bay
	C	Substrate/Benthos	
D11	R	Substrate	Flooded Tract
	C	Substrate/Benthos	
	L	Substrate	
D19	R	Substrate	Flooded Tract
	C	Substrate/Benthos	
	L	Substrate	
D28A	R	Substrate/Benthos	River Channel
	L	Substrate/Benthos	

* Sites are determined while facing downstream (Right, Center, Left).

** Substrate samples consist of one random grab.
Benthic samples consist of three grabs.

After collection, each sample was rinsed through a screen with 0.6-mm openings. All material remaining after washing was preserved with 25% formalin for laboratory analysis.

Laboratory analysis of the preserved samples consisted of sorting, identifying, and enumerating all whole organisms. Identifications were made to the lowest taxonomic level possible, usually species. A taxonomic list of all organisms identified from the monitoring samples (Appendix A) was continuously maintained. Scientific names are updated annually, as new organisms are identified or existing organisms are reclassified.

Hydrozoology, a private laboratory under contract with the State of California, analyzed all benthic samples. All organisms collected are preserved in ethyl alcohol and archived after identification and enumeration. Identification and enumeration data are entered into an electronic data base maintained by DWR as an SAS data set.

One of the goals in analyzing the benthic monitoring data was to arrange the monitored variables, represented by the sampling sites and collected species, in an ecologically meaningful order. The distribution of most species in a community is

presumed to reflect the influence and variation of major environmental factors. Although such factors, and the species reflecting them, may not display a simple, continuous trend from one extreme to the other, the actual range of conditions in time or space can be viewed as a gradient. Environmental mosaics and interspecific relationships can make the interpretation of data from a community complex. Ordination techniques are recognized as methods that permit identification of major factors controlling the distribution of species²⁸. The benthic monitoring data were analyzed using the ordination method of correspondence analysis available in the SAS package.

Correspondence analysis and its application are described in detail by Greenacre²⁹. CA is a preferred method of ordination because the data transformation does not assume a linear relationship among the variables, which rarely occurs in ecological data. Also, there is a direct relationship between the species and sampling site scores because CA scales both the rows (species) and columns (sites) of the data matrix in the same manner. This second characteristic of CA allows the plotting and interpretation of both species and sampling site scores on the same axes. The CA scores from an individual axis can also be treated as normal random variates, as the scores on each axis are independent of one another. For this reason, CA scores can be used in further statistical tests to examine relationships between environmental variables and the benthic community. In this report, the CA sites scores were used in simple linear regression analyses to test for significant relationships between the CA scores and a variety of biotic and abiotic environmental variables measured at the benthic monitoring stations.

Sediment

Sediment composition was also measured as part of the benthic monitoring program. A single sediment sample was collected each month from 13 sites (Table 5). General trends in sediment composition are described for all sites where sediment samples were collected. Trends are depicted as the mean annual percentage of fines (silt and clay) and the mean annual percentage of organic material through time.

28 JJ Gonor and PF Kemp. 1984. *Procedures for Quantitative Ecological Assessments in Intertidal Environments*. US Environmental Protection Agency. Corvallis, OR.

29 MJ Greenacre. 1984. *Theory and Applications of Correspondence Analysis*. Academic Press. London.

For this report, however, the relationship between sediment composition and benthic species composition was investigated using data only from those eight locations where both benthic biota and sediment samples were collected.

Sediment samples were collected with the same winch and Ponar dredge set-up used in the infaunal sampling. A 1-liter subsample of sediment was haphazardly selected from a single dredge sample and stored. All sediment samples were analyzed at the DWR Soils Laboratory. Routine analysis of the sediment sample included determining the percent size fractions with the use of a mechanical sieve and hydrometer. Using the size fractionation data, the sample was categorized (on a percentage basis) as fines (silt and clay particles less than or equal to 0.08 mm in diameter), sand (particles greater than 0.08 mm in diameter), or gravel (particles greater than 2.5 mm in diameter). (During 1980 through 1990, no gravel was detected at any of the routinely monitored sites.) The percent organic content of each sediment sample was also routinely determined from the loss in weight of an oven-dried sample burned at 404°C for 8 hours. All laboratory analysis procedures follow the American Society for Testing Methods³⁰. Data analyses included sediment data collected from 1981 through 1990, which were stored in a personal computer data base. Data for 1980 were not available.

Water Quality

DWR collects water quality data at 26 stations throughout the upper estuary (Figure 1) as part of the environmental monitoring stipulated in Decisions 1379 and 1485, which started in 1975 and continues to the present. Stations are generally sampled monthly between November and February and bimonthly the rest of the year. All samples are collected from a depth of 1 meter by submersible pump or Van Dorn water sampler at or near high slack tide. Field measurements included water temperature, specific conductance, pH, dissolved oxygen, turbidity, and Secchi disc depth. All other analyses were completed at the DWR Chemistry Laboratory

using standard analytical methods.³¹ Data are stored on the Environmental Protection Agency's STORET system.

For this report, trends in surface water temperature, specific conductance, and volatile suspended solids were characterized for three regions of the upper estuary (Figure 4). Specific conductance measurements were converted to salinity values using the formula:

$$\text{Salinity (parts per thousand)} = -100(\ln(1-EC/178.5))$$

Where: EC = specific conductance, in milliSiemens per centimeter.

Water temperature and specific conductance were measured on-site using electronic sensing equipment. The concentration of volatile suspended solids was determined from the loss in weight of an oven-dried total suspended solids sample burned at 550°C for 24 hours.³² For all variables, annual means and 95% confidence intervals were calculated on a regional basis by pooling monthly data from all stations in a region and then averaging them over the calendar year.

Phytoplankton

DWR routinely sampled the composition and biomass of phytoplankton at numerous locations in the upper estuary as part of its environmental monitoring program. Taxonomic composition was assessed through microscopic analysis of water samples. Biomass measurements, used primarily to document the occurrence of abrupt increases in phytoplankton concentration (phytoplankton blooms), were estimated from measurements of chlorophyll *a* concentration of water samples routinely collected from 26 stations in the upper estuary (Figure 1). Changes in phytoplankton composition and biomass are summarized here using data from 16 stations and three regions (Figure 3).

Trends in chlorophyll *a* concentration anomalies were used to determine if total phytoplankton biomass changed over time. In this analysis, an anomaly value represents the mean annual concentration

30 American Society for Testing Materials. 1992. Annual book of ASTM standards, Section Four, Volume 4.08. *Soil, Rock Building Stones, and Geotextiles*. American Society for Testing Materials, Philadelphia.

31 LS Clesceri, AE Greenberg, RR Trussell (editors). 1989. *Standard Methods for the Examination of Water and Wastewater*. 17th edition. American Public Health Association, Washington, DC.

32 Clesceri *et al*, 1989; cited.




Figure 3
STATIONS AND REGIONS USED IN WATER QUALITY AND PHYTOPLANKTON ANALYSES

after subtraction of the long-term average. This transformation tends to dampen the influence of short-term changes such as those due to season or salinity. Anomalies greater than zero indicate the annual mean concentration was greater than the long-term average; anomalies less than zero indicate the annual mean concentration was less than the long-term average. More information on calculation of anomalies and a discussion of long-term trends in chlorophyll *a* for this estuary are available in Kimmerer's 1992 report³³.

To summarize anomaly data, annual mean anomaly values and 95 percent confidence intervals were calculated from a core data set after pooling data from stations within three geographically defined regions (Figure 3). Results are only presented for the regions from which benthic samples were collected, and only for 1978 through 1990. Linear regressions of trends in chlorophyll *a* anomalies were tested to determine if the slope of a regression line differed significantly from zero; non-linear relationships were not tested.

33 WJ Kimmerer. 1992. *An Evaluation of Existing Data in the Entrapment Zone of the San Francisco Estuary*. Interagency Ecological Studies Program, Technical Report 33. Department of Water Resources.

Zooplankton

Zooplankton abundance and distribution were monitored by the Department of Fish and Game. Zooplankton were sampled from a boat by towing a collection net from bottom to surface in a step-wise oblique 10-minute tow. Sampling surveys were conducted once in March, once in November, and

twice each month in April through October. Laboratory analyses included sorting, identification, and enumeration of all samples.

Data are presently stored as SAS data sets. Methods for the zooplankton field sampling and laboratory analyses are described in more detail by Obrebski and others³⁴.

³⁴ S Obrebski, JJ Orsi, W Kimmerer. 1992. *Long-Term Trends in Zooplankton Distribution and Abundance in the Sacramento-San Joaquin Estuary*. Interagency Ecological Studies Program for the Sacramento-San Joaquin Estuary, Technical Report 32. Department of Water Resources.

TRENDS IN FACTORS INFLUENCING THE BENTHOS

In their community profile of the soft-bottom benthos of San Francisco Bay, Nichols and Pamatmat³⁵ concluded that many of the most dramatic inter-annual community changes may be attributable to extreme deviations in the physicochemical environment. These deviations may influence both the timing and success of recruitment and the survival of existing individuals. In this chapter we describe the trends in various physicochemical and biological variables that may affect the benthos of the upper estuary.

Freshwater Flow

Freshwater flow strongly affects the physicochemical environment of the upper estuary. In addition to the direct effects on sediment composition and stability, freshwater flows affect salinity, water clarity, water temperature, and several other water quality variables. About 40% of California's watershed drains into the Sacramento-San Joaquin estuary, with the largest segment of this fresh water (about 72%) entering from the Sacramento River.³⁶

From 1980 through 1990, the amount of fresh water entering the estuary has ranged over wide extremes, as indicated by mean annual Sacramento River flows at Sacramento (Figure 4). During this period, mean annual flows to the estuary were highest in 1983. They generally declined through 1985 and then increased sharply in 1986 because of extremely heavy precipitation during February. Freshwater flows have been persistently low since 1987, as a result of one of the most severe droughts in recent history.

The variability in freshwater flows within a year may be as important to the composition, abundance, and distribution of the benthos as the annual amount of freshwater entering the system. As indi-




Figure 4
MEAN ANNUAL FLOW,
SACRAMENTO RIVER AT SACRAMENTO
Annual Means \pm 95% C.I.

cated by Sacramento River flows, the monthly variability generally increases with the annual average. Thus, the low freshwater flows that prevailed between 1987 and 1990 were accompanied by reduced intra-annual variability. This variability, which is strongly related to seasonal changes, may be important in determining the recruitment success and distribution of benthic organisms with planktonic life stages.

Water Quality

Although many water quality variables are measured throughout the upper estuary, only a few could have directly affected benthic species composition and abundance. Many of the variables, such as nutrient concentrations or total dissolved solids, have little direct effect on the benthos over the range of values measured in this estuary. Other water quality variables, such as temperature, have been very stable over the years (Figure 5) and show little

35 FH Nichols and MM Pamatmat. 1988. *The Ecology of the Soft-Bottom Benthos of San Francisco Bay: A Community Profile*. US Fish & Wildlife Service Biological Report 85(7.19). 73 pp.

36 Comprehensive Region Framework Study Committee. 1971. *Comprehensive Framework Study, California Region; Appendix V, Water Resources*. US Bureau of Reclamation. 339 pp. plus tables and maps.




Figure 5
TRENDS IN WATER TEMPERATURE
Annual Means +/- 95% C.I.

connection to the variability in abundance and distribution of benthic organisms.

Freshwater inflow to the estuary is a principal determinant of the estuary's physical and chemical environment. The extreme fluctuations in freshwater flows observed between 1980 and 1990 produced similar, but inversely related, fluctuations in salinity levels. Salinity patterns are described on a regional basis because of the wide range in salinity between Suisun Bay and the central delta.

Fluctuations in salinity have been most extreme in the Suisun Bay region (Figure 6). Between 1980 and 1990, annual mean salinity in Suisun Bay ranged from 0.18 to 10.6 ppt. Salinity patterns in the western and central delta regions were similar to the pattern in the Suisun Bay region, but the range in values was smaller. In the western delta region, annual mean salinity ranged from 0.08 to 2.27 ppt (Figure 7). In the central delta region, annual mean salinity ranged from 0.06 to 0.25 ppt (Figure 8).




Figure 6
TRENDS IN ANNUAL SALINITY, SUISUN BAY REGION
Annual Means +/- 95% C.I.




Figure 7
TRENDS IN ANNUAL SALINITY, WESTERN DELTA REGION
Annual Means +/- 95% C.I.




Figure 8
TRENDS IN ANNUAL SALINITY, CENTRAL DELTA REGION
Annual Means +/- 95% C.I.

Sediments

The substrate throughout the study area is entirely soft-bottom. The composition of the substrate is largely determined by the sediments present and the physical processes (wind and water motion) that move these materials. Changes in sediment composition can also occur as a result of bioturbation and biogeochemical processes, but in this estuary these effects are thought to be relatively minor compared to the physical processes. It is important to understand the trends in sediment composition, which can directly affect the benthos in terms of both the community composition and species abundance.

Both the inorganic and organic sediment fractions at sites D7-C and D7-R were extremely stable from 1981 to 1990 (Figure 9). The inorganic fraction was consistently dominated by fine material (silt and clay) and showed little variability. The organic fraction, which was mainly particulate organic matter, ranged from 7 to 10% at both sites.

The inorganic sediment fraction at Station D4 varied considerably at all sites (Figure 10). Sediment composition was most consistent in the center channel, where sand (% sand = 100 - % fines) was the dominant substrate type. From 1981 to 1990, the mean annual percentage of fines was consistently below 25%. The




Figure 9

TREND IN ANNUAL PERCENTAGE OF FINE INORGANIC AND ORGANIC SEDIMENTS, GRIZZLY BAY, STATION D7

inorganic fraction was much more variable at D4-R. The mean annual percentage of fines ranged from 54% in 1984 to 94% in 1987 and was inversely related to Sacramento River flow (Figure 4). The percentage of fines at D4-R was generally higher after 1985, but declined sharply in 1990. Unlike D4-R, the mean annual percentage of fines at D4-L was not clearly related to Sacramento River flow. At the left bank,

the mean annual percentage of fines ranged from 23% in 1987, to 61% in 1983.

The organic sediment fraction was much more stable than the inorganic fraction at station D4 (Figure 10). Organic content was lowest at D4-C, where the mean annual percentage ranged from 1 to 3%. The organic fraction was slightly higher at D4-R,




Figure 10
TREND IN ANNUAL PERCENTAGE OF FINE INORGANIC AND ORGANIC SEDIMENTS, SACRAMENTO RIVER, STATION D4

where mean annual percentages ranged from 4 to 6%. Organic content was highest but most variable at D4-L, where the mean annual percentage ranged from 8 to 13%.

The percentage of fines at Station D11 increased significantly ($P < 0.05$) at all sites from 1981 through 1990 (Figure 11). The inorganic fraction was most stable at D11-C, where mean annual percentage of fines ranged from 76 to 98%. The inorganic fraction varied most at

D11-R, where mean annual percentage of fines ranged from 29 to 91%. By comparison, the percentage of fines was generally higher and less variable at D11-L, where mean annual percentage ranged from 37 to 93%.

The organic fraction at station D11 showed no significant trend over time at any of the sites (Figure 11). Organic content was generally lowest and most variable at D11-R (mean annual percentage 4-10%) and highest at D11-L (mean annual percent-




Figure 11
TREND IN ANNUAL PERCENTAGE OF FINE INORGANIC AND ORGANIC SEDIMENTS, SHERMAN LAKE, STATION D11

at the channel stations (D4, D28A), but many locations had increased amounts of fines during the drought (1987–1990). Organic content showed no significant trend through time at any station. Qualitative laboratory observations found peat to be the dominant organic material at all sites except at D7.

Food Supply

Abundance and distribution of benthic organisms can be affected by numerous biological, physical, and chemical processes. Food supply is a biological factor that affects growth rates, survivorship, and fecundity of benthic organisms. Thus, changes in food supply influence several life stages, leading to direct population effects.

The relationship between food supply and the abundance of a benthic organism may not be linear. If sufficient food is available to consumers, their abundance will not be affected by any further increase in food supply. However, food concentrations that remain chronically below the level required for growth and/or reproduction will have deleterious effects on the consumer's abundance. Determining the effects of changes in the quality and quantity of food, whether through inferences or specific studies, is difficult. Trends in food abundance and benthos abundance can be compared, using correlation analysis of monitoring data, to determine if relationships exist; however, the cause and effect of these relationships are inferred and cannot be proven by the analysis. The major use of the correlation test is to identify relationships that warrant further investigation.

In this section, trends in three potential food sources, volatile suspended solids, phytoplankton, and zooplankton, are described based on analyses of routine monitoring data. Volatile suspended solids (which include phytoplankton and other particulate organic matter) are thought to be primary food sources for many benthic invertebrates. Although

zooplankton are a minor food source for some benthic organisms,³⁷ a description of their trends in relation to other food sources provides a more complete picture of possible secondary links between the lower food chain levels and the effects that zooplankton grazing on phytoplankton may have on the benthos. Other items, such as benthic microalgae and bacterioplankton that may also be important benthic food sources, were not measured during this study.

Volatile Suspended Solids

Volatile suspended solids are the organic portion of total suspended solids. This suspended organic material may represent a food source for both benthic and pelagic organisms, but studies to determine the importance of VSS to the benthic food supply have not been completed for this estuary.

Mean annual concentration of volatile suspended solids was generally highest in Suisun Bay and lowest in the central delta (Figures 14–16). In Suisun Bay, mean annual concentration ranged from 4.3 to 10.1 mg/L. The concentration decreased significantly ($P < 0.05$) between 1978 and 1983 but showed no significant trend thereafter. VSS levels in 1983 were about 40% lower than in 1978. In the western delta, mean annual concentration ranged from 3.2 to 6.7 mg/L. Concentrations were highest in the western delta during 1978, were lower but stable from 1979 through 1985, and then increased somewhat but remained variable thereafter. In the central delta, mean annual VSS concentration ranged from 2.4 to 4.7 mg/L from 1978 through 1990. The concentration declined between 1978 and 1979 but remained fairly stable thereafter. General trends among the three regions suggest the concentration of VSS declined early on over much of the study area. Although concentrations were relatively stable in both the central and western delta from 1980 through 1990, VSS concentration did decline in the Suisun Bay region.

37 W Kimmerer, personal communication.




Figure 14
TREND IN VOLATILE SUSPENDED SOLIDS,
SUISUN BAY REGION
 Annual Means +/- 95% C.I.




Figure 15
TREND IN VOLATILE SUSPENDED SOLIDS,
WESTERN DELTA REGION
 Annual Means +/- 95% C.I.




Figure 16
TREND IN VOLATILE SUSPENDED SOLIDS,
CENTRAL DELTA REGION
 Annual Means +/- 95% C.I.

Phytoplankton

Between 1978 and 1990, seasonal peaks in phytoplankton biomass (blooms) occurred in all regions of the upper estuary examined (Figure 17). Phytoplankton blooms typically occur between spring and fall and are most often dominated by one of four diatom genera: *Skeletonema* sp., *Thalassiosira* sp., *Cyclotella* sp., or *Melosira* sp. From 1980 through 1990, *Melosira* sp. was the dominant bloom organism in the delta, and *Thalassiosira* sp. dominated in Suisun Bay.

In the central delta region, mean annual chlorophyll *a* concentrations were moderate (4–12 µg/L) and variable between 1978 and 1990 (Figure 18). Annual anomalies of chlorophyll *a* concentration, which showed no statistically significant linear trend, were associated with relatively large confidence limits. In these cases, a high seasonal variability may be masking the lower variability in annual changes of phytoplankton biomass. Anomalies of chlorophyll *a* did exhibit a convex curve-shaped trend, with negative values between 1978 and 1981, positive values between 1982 and 1986, and negative values between 1987 and 1990.

The western delta is a convergence zone between the northern and central delta regions and Suisun Bay (Figure 4). As a result, physical, chemical, and biological processes in this region are often driven by events that originate in the surrounding areas. Annual variations in the mean chlorophyll *a* concentrations were generally moderate (4–12 µg/L) (Figure 18). Annual anomalies of chlorophyll *a* show phytoplankton biomass has declined significantly ($P < 0.05$) in this region, particularly during the last 4 years. Anomaly values increased between 1978 and 1982, declined sharply in 1983, increased through 1986, and declined steadily thereafter. The sharp decline in 1983 may reflect a downstream shift in the position of the entrapment zone and associated phytoplankton as a result of the extremely high outflows that occurred during winter and spring.

In the Suisun Bay region, mean annual chlorophyll *a* concentrations remained below 5 µg/L from 1978 through 1990 (Figure 18). Annual anomalies of chlorophyll *a* show a significant ($P < 0.05$) linear decrease in phytoplankton biomass over the last 13 years. The 1990 anomaly value was somewhat higher than the 1989 value; however, the average phytoplankton biomass generally remained at extremely low levels in this region.

Overall, phytoplankton biomass has declined significantly since 1986 throughout much of the upper estuary. In addition, a decrease in the frequency and

intensity of phytoplankton blooms in many regions of the upper estuary has been noted since 1987.




Figure 17
CHLOROPHYLL a CONCENTRATIONS AT VARIOUS STATIONS IN THE UPPER ESTUARY

Letters above peaks denote dominant bloom organisms:
 C = *Cyclotella* sp., S = *Skeletonema* sp., T = *Thalassiosira* sp., M = *Melosira* sp.
 Site locations are shown in Figure 1.




Figure 18
CHLOROPHYLL *a* CONCENTRATION AND ANOMALIES VERSUS TIME
 Graphs on left are mean annual concentration, in µg/L.
 Graphs on right are mean annual anomalies and 95% confidence intervals (vertical bars).

Zooplankton

Zooplankton occupy an intermediate level in many estuarine food chains, because most feed on primary carbon sources and because they are a major food source for various life stages of several estuarine fishes. In this estuary, salinity and season are the major factors related to between-year and within-year fluctuations in zooplankton stocks.³⁸

The analysis of DFG zooplankton compliance monitoring data completed by Obrebski and others in 1992 used methods similar to those described above for the analysis of phytoplankton data to determine long-term trends in zooplankton abundance. Results show 12 of the 20 zooplankton taxa routinely sampled have undergone significant declines in abundance between 1972 and 1988. Seven taxa exhibited no abundance trend and one introduced copepod, *Oithona davisae*, increased in abundance (Table 6). Obrebski *et al* also examined regional and seasonal trends in zooplankton abundance. Results showed that declines in zooplankton abundance were scattered throughout the upper estuary but were more prevalent in the Sacramento and San Joaquin rivers than in Suisun Bay (Table 7). Zooplankton abundance trends exhibited no clear seasonal pattern (Tables 6 and 7).

Table 6 SUMMARY OF CHANGES IN SUISUN BAY/DELTA ZOOPLANKTON ANOMALIES Results of Regression Analysis of Annual Mean Anomalies				
	Pooled Data (All Months)	Spring (Mar-May)	Summer (Jun-Aug)	Fall (Sep-Nov)
COPEPODS				
<i>Acartia</i>	O	O	O	O
<i>Diaptomus</i>	D**	O	D**	D***
<i>Eurytemora</i>	D***	D**	D***	D**
<i>Harpacticoids</i>	D**	D**	D*	D*
<i>Cyclopoids</i>	D*	O	O	D*
<i>Sinocalanus</i>	O	O	O	O
<i>Limnithona</i>	O	O	O	O
<i>Oithona davisae</i>	I*	O	I*	I*
CLADOCERA				
<i>Bosmina</i>	O	O	O	O
<i>Daphnia</i>	D*	O	D*	D*
<i>Diaphanosoma</i>	D*	U*	D*	D***
ROTIFERA				
<i>Asplanchna</i>	D**	D*	D**	D**
<i>Keratella</i>	D***	D**	D**	D***
<i>Polyarthra</i>	D***	D***	D***	D***
<i>Synchaeta</i> spp.	O	O	O	O
<i>Synchaeta bicornis</i>	D***	D**	D***	D***
<i>Trichocerca</i>	D***	D**	D**	D**
OTHER				
<i>Neomysis</i>	D*	O	O	D**
Barnacle Nauplii	O	O	O	O
Crab Zoea	O	O	O	O
O = No Change, D = Decline, I = Increase, U = U-Shaped Trend				
* 0.01 <P <0.05				
** 0.001 <P <0.01				
***P <0.001				
SOURCE: S. Obrebski, J.J. Orsi, W. Kimmerer. 1992. Long-Term Trends in Zooplankton Distribution and Abundance in the Sacramento-San Joaquin Estuary. Interagency Ecological Studies Program, Technical Report 32. Department of Water Resources.				

38 Obrebski *et al*, 1992; cited.

Table 7
SUMMARY OF REGIONAL CHANGES IN ABUNDANCE OF ZOOPLANKTON TAXA THAT DECLINED BETWEEN 1972 AND 1987
 Numbers are adjusted R² for either a linear or quadratic model, whichever yielded the highest R².
 AL = Data pooled for all months, SP = Spring, SU = Summer, FA = Fall

	Suisun Bay				Sacramento River				Lower San Joaquin River				Western Delta				Entrapment Zone				Upper San Joaquin River			
	AL	SP	SU	FA	AL	SP	SU	FA	AL	SP	SU	FA	AL	SP	SU	FA	AL	SP	SU	FA	AL	SP	SU	FA
<i>Diaptomus</i>	.33 *	NS	NS	.55 **	.29 *	NS	.24 *	.36 *	.23 *	NS	NS	.41 *	.62 **	.25 *	.57 ***	.60 **	.52 **	NS	.31 *	.67 ***	.28 *	NS	NS	.57 **
<i>Eurytemora</i>	.26 *	NS	.20 *	.22 *	.57 **	.50 **	.42 *	NS	.87 ***	.50 **	.44 **	.61 **	.40 *	.36 *	.42 *	NS	.68 ***	NS	.52 **	.63 **	.39 **	NS	.61 ***	.21 **
Harpacticoids	NS	NS	NS	NS	.20 *	NS	.29 *	NS	.73 ***	.58 ***	.58 **	.61 **	NS	NS	.23 *	NS	.57 **	NS	NS	.59 **	NS	NS	NS	NS
Cyclopoids	NS	NS	NS	NS	NS	NS	NS	NS	.22 *	NS	NS	.23 *	.40 *	NS	.31 *	.34 *	.37 *	NS	NS	NS	NS	NS	NS	NS
<i>Daphnia</i>	NS	NS	NS	NS	NS	NS	NS	NS	.32 *	NS	.31 *	.36 *	.48 *	NS	.38 *	.42 **	.59 **	NS	.41 *	.41 **	NS	NS	NS	NS
<i>Diaphanosoma</i>	NS	NS	NS	NS	.72 ***	.60 **	.63 **	.41 **	.78 ***	NS	.45 *	.74 ***	.44 **	NS	.35 *	.64 *	.37 *	NS	NS	.60 **	NS	NS	NS	NS
<i>Neomysis</i>	.45 **	NS	.45 *	.70 ***	.62 **	.46 **	NS	.62 **	.39 *	NS	NS	.57 **	.65 ***	.62 **	NS	.47 **	.55 **	.58 **	NS	.57 **	.36 **	NS	.23 *	.61 **
<i>Trichocerca</i>	.61 **	.54 **	.59 **	.58 **	NS	NS	.29 *	NS	.30 *	.21 *	.51 **	NS	NS	NS	NS	NS	.58 **	.46 **	.52 **	NS	.59 **	.47 **	.59 **	.55 **
<i>Polyarthra</i>	.69 ***	.74 ***	.47 **	.64 **	.87 ***	.58 **	.86 ***	.69 ***	.93 ***	.72 ***	.91 ***	.73 **	.89 ***	.73 **	.86 ***	.88 ***	.93 ***	.78 ***	.87 ***	.80 ***	.73 **	.73 **	.64 **	.70 **
<i>Synchaeta bicornis</i>	.46 **	.30 *	.68 ***	.47 **	.62 ***	.51 **	.49 **	.35 *	.53 **	.34 *	.50 **	.31 *	.59 ***	NS	.49 **	.34 *	.58 **	.38 *	.54 **	.54 **	.50 **	.30 *	.45 *	NS
<i>Asplanchna</i>	.39 *	NS	NS	NS	.84 ***	.77 ***	.79 ***	.75 ***	.82 ***	.70 ***	.81 ***	.74 ***	.50 **	NS	.59 **	.46 *	.76 ***	.62 **	.72 ***	.60 **	.53 **	.38 *	.23 *	NS
<i>Keratella</i>	NS	NS	NS	NS	.89 ***	.74 ***	.83 ***	.75 **	.90 ***	.60 **	.87 ***	.71 **	.78 ***	.57 **	.74 ***	.85 ***	.91 ***	.74 ***	.70 **	.88 ***	.77 **	.64 **	.51 *	.71 **

NS Not Significant
 * 0.01 < P < 0.05
 ** 0.001 < P < 0.009
 *** P < 0.001

SOURCE: S. Obrebski, J.J. Orsi, W. Kimmerer. 1992. *Long-Term Trends in Zooplankton Distribution and Abundance in the Sacramento-San Joaquin Estuary*. Interagency Ecological Studies Program, Technical Report 32. Department of Water Resources.

CHANGES IN THE BENTHOS

The large number of benthic species identified through the monitoring program and the extreme variability in their abundance present a major challenge to efforts to detect long-term changes in the benthos and to identify probable causes for those changes. From 1980 through 1990, a total of 196 species were identified from all stations. On the average, 12 species were identified at any one location each month. The majority of these organisms generally occurred in low (<100 individuals/m²) abundance or were found only sporadically. Typically, the four numerically dominant organisms at each location accounted for at least 80% of the total abundance at any one time.

There was also substantial temporal variability within the benthos of the upper estuary. It was not uncommon for monthly abundance to vary by an order of magnitude. An examination of the graphs in Appendix B gives some indication of the temporal variability in community abundance. Also, the graphs of the mean monthly abundance of the four numerically dominant organisms illustrate the temporal and spatial variability of individual species.

Correspondence analysis (CA) was used to investigate long-term changes in benthic species abundance and persistence, given the large number of species collected and the temporal and spatial variability common in the benthos. (CA methods are described in more detail in Chapter 2.) It is important to remember, however, that results for each site are relative responses to changes in benthic species abundance and persistence based on the suite of sites compared. Thus, the choice of sites compared directly affects the results.

Data from all sites were analyzed initially to determine if any differences among sampling sites existed. Annual mean abundances of the species were used to reduce seasonal variation within the data set, which could obscure long-term patterns of species abundance differences among sampling sites. CA results from the first CA dimension, which explains 33.7% of the total variation in the data set, show several distinct responses among the sites (Figure 19). Between 1980 and 1984, benthic species

abundance and persistence was remarkably stable, with little deviation among sites or years. Beginning in 1985, however, the patterns of response among sites diverged. Site D7-C (Grizzly Bay) showed the largest response; followed by the three Sacramento River sites (D4-R, D4-L, D4-C), which showed an intermediate response. The remaining upper estuary sites (D11-C, D19-C, D28A-L, D28A-R) showed little response, which suggests little change in benthic species abundance and persistence during the sampling period. Using these response patterns, sites were grouped as: Grizzly Bay (D7-C); Sacramento River (D4-R, D4-L, D4-C); and eastern sites (D11-C, D19-C, D28A-L, D28A-R). Using these groupings, additional correspondence analyses were performed to further examine temporal changes in benthic species abundance and persistence and their potential causes.

Although first ordination dimension results explain the largest portion of the total deviation, examination of the second and third dimensions of ordination can reveal other meaningful patterns of response. In theory, response patterns for all ordination dimensions that explain some portion of the total variation can be examined; however, ordination




Figure 19
FIRST DIMENSION CORRESPONDENCE ANALYSIS,
ALL SITES

results explaining successively smaller portions of the variation may not be meaningful. Only results from the first and second dimensions of ordination are presented here. The third dimension results typically explained less than 10% of the total deviation, probably representing random variation.

In the CA, which compared all sites, second ordination dimension results explained 14.6% of the total variation (Figure 20). With the exception of Site D7-C, responses for all sites show relatively small changes from year to year. These response patterns are thought to represent the ongoing underlying variability in benthic habitat. Constituents such as water temperature, sediment composition, and food supply and continuous processes such as tidal action all contain inherent variability that contributes to this ongoing habitat variability. This variability is thought to be the source of these species and community changes.

In contrast, a definite response in the pattern of species abundance and persistence at D7-C was detected in the second ordination dimension. The pattern and timing of this response is similar to the first dimension response suggesting similar processes are responsible.




Figure 20
SECOND DIMENSION CORRESPONDENCE ANALYSIS,
ALL SITES

Grizzly Bay Site, D7-C

The CA analysis for Site D7-C (Figures 21 and 22) used monthly mean abundances instead of annual means. Monthly values were used to show the patterns of variation at an individual sampling site that are a function of seasonal fluctuations in species abundance and persistence. Overall, the pattern of change shown in the first dimension results for D7-C in Figure 21 is similar to the pattern for that site seen in the analysis of all sites (Figure 19). These results show that annual mean abundance is useful for station comparisons and can also be used to show long-term community changes.




Figure 21
FIRST DIMENSION CORRESPONDENCE ANALYSIS,
GRIZZLY BAY, SITE D7-C




Figure 22
SECOND DIMENSION CORRESPONDENCE ANALYSIS,
GRIZZLY BAY, SITE D7-C

First ordination dimension results for D7-C show species abundance and persistence oscillated between periods of rapid change and transition (1980-1982 and 1985-1988) and periods of relative stability (1982-1985 and 1988-1990) (Figure 21). With the exception of 1989, the pattern of response tracks fluctuations in mean annual salinity for the Suisun Bay region. This pattern shows little change between 1988 and 1990, even though mean annual salinity declined in 1989. However, the salinity decline was relatively small and short-lived compared to salinity fluctuations in other years. Inspection of CA and salinity curves (Figure 21) indicates a 6- to 12-month lag between changes in annual average salinity and CA score. This lag may reflect the benthos dampening the effect of short-term salinity changes. We interpret the overall pattern to reflect a response of species abundance and persistence to abiotic (salinity) and biotic (invasion of exotic species) changes.

Second dimension results for D7-C show continual oscillations in species abundance and persistence

with a periodicity of 3 to 4 years (Figure 22). Patterns in the first and second dimension ordination values mirror each other from 1980 to 1986 and then become inversely related. Anomalous shifts in both first and second dimension response values occurred in 1981 and 1985. The second dimension pattern may reflect a community-level response to establishment of exotic species combined with irregular physical disturbance of the habitat such as occurred in the floods of 1983 and 1986 and the drought that persisted from 1987 through 1990.

Another way to examine the CA results is to graph the individual species scores in the first and second dimensions of ordination (Figure 23). In these graphs, with the x = zero / y = zero point as the centroid, the individual species scores indicate the influence of the species on the observed CA values through time and the relative contribution (loading) of each species to the CA value. The farther an organism is from the centroid, the greater its influence (loading) on the CA value.




Figure 23
SPECIES SCORES, GRIZZLY BAY, SITE D7-C

Individual species scores for the Sacramento River sites show that three groups of species are responsible for the CA site score patterns of response (Figure 26). The first group consists of organisms with only negative scores in the first ordination dimension and both positive and negative scores in the second ordination dimension. This is the largest group and consists mostly of brackish and freshwater species such as *Corophium stimpsoni*, *Limnodrilus hoffmeisteri*, and *Manayunkia speciosa*. Various members of this group are always present at one or more of the sites.

The second group includes organisms with positive scores in the first ordination dimension and negative scores in the second ordination dimension (Figure 26). This group includes estuarine species such as *Balanus improvisus* and *Boccardia ligERICA*. These estuarine species were only found at appreciable concentrations after 1986, when salinity increased

and remained at higher levels. However, only *B. ligERICA* has been numerically dominant since 1987.⁴⁴

The third group includes two introduced organisms, *P. amurensis* and *H. hinumensis*, which have positive species scores in both the first and second dimensions of ordination and contribute the largest loadings (ie, their positions are farthest from the centroid). These organisms became numerically dominant at one or more of the Sacramento River sites after 1986.⁴⁵

CA site scores in the first ordination dimension and the individual species scores for the Sacramento River sites show a pattern in the benthos that is very similar to the one observed for D7-C. Between 1980 and 1986, patterns varied in relation to changes in salinity. The floodflows in 1986 substantially reduced population abundances, resulting in relatively large amounts of open space.




Figure 26
SPECIES SCORES, SACRAMENTO RIVER, STATION D4

44 DWR, 1992; cited.

45 DWR, 1992; cited.

Drought-associated increases in salinity, beginning in 1987, limited the recolonization of resident brackish water species. Meanwhile, *P. amurensis* and *H. hinumensis* were rapidly colonizing Suisun Bay. In relatively stable environmental conditions, these exotic organisms, along with other resident species (particularly *Corbicula fluminea*), were able to colonize the Sacramento River sites. Patterns of response for CA scores from all three sites showed little change after 1988. The persistent dominance of *P. amurensis*, *H. hinumensis*, and *C. fluminea* appears to have resulted in a new and stable benthic community at the Sacramento River sites. The species composition of this new community, however, is substantially different from the community observed at D4 in the early 1980s.

Correlation analyses show a significantly positive relationship between salinity and CA scores at all D4 sites (Table 8). Sacramento River flow is negatively related to species abundance and persistence at all D4 sites, but the relationship is not significant. The percentages of silt and clay were significantly related to benthic species abundance and persistence only at D4-C. A significantly negative relationship between chlorophyll *a* concentration and benthic species abundance and persistence at D4-C may be due to the presence of both *P. amurensis* and *C. fluminea*.

In general, the channel environment of D4 is more variable than the shoal region of Grizzly Bay or the lacustrine environment of Sherman Lake (D11) and Franks Tract (D19). This is particularly true for sediment composition. Nevertheless, fluctuations in sedi-

ment composition appear to have played a primary role in determining the pattern of benthic species abundance and persistence only at the center of D4 but not at the banks. Overall, results suggest that salinity had the broadest and most significant influence on benthic species abundance and persistence at D4 (Table 8). Thus, as with D7-C, both abiotic (salinity and sediment composition) and biotic (invasion of exotic species) processes have acted to alter the benthos at D4.

Eastern Sites, D11-C, D19-C, D28A-L, D28A-R

CA results for the first dimension from the third group of sites (eastern sites D11-C, D19-C, D28A-L, D28A-R) show response patterns that differ from those for the other two site groups (Figure 27). First dimension CA results explained 25.4% of the total variation. In general, response patterns in the first dimension, with the exception of values for D11-C, showed little change over time. The pattern for D11-C showed little change between 1980 and 1985 but moved steadily downward thereafter. None of these response patterns track the trend in average annual salinity for this region.

Response patterns for CA results in the second dimension were more evident, although only 17.6% of the total sample variation was explained (Figure 28). CA site scores for all sites showed a consistent pattern of change from 1980 through 1990. Between 1980 and 1982, CA scores were stable, with little fluctuation among years or sites. However, a




Figure 27
FIRST DIMENSION CORRESPONDENCE ANALYSIS,
EASTERN STATIONS, D11, D19, AND D28A




Figure 28
SECOND DIMENSION CORRESPONDENCE ANALYSIS,
EASTERN STATIONS, D11, D19, AND D28A

major shift occurred in CA scores for all of the eastern sites from 1982 to 1984. This shift occurred over a relatively short time and led to establishment of a new and stable benthic community. These response patterns may be due to habitat changes that were not measured by the monitoring program.

Individual species scores for the eastern sites show that, with the exception of *Hemileucon hinumensis*, no one species overwhelmingly influenced the patterns of CA scores in the first or second dimension of ordination (Figure 29). Within this group of sites, *H. hinumensis* has only been collected from D11-C and only since 1987. This suggests the presence of *H. hinumensis* at D11-C is at least partly responsible for the negative trend in the first dimension CA scores for this site.

Results of correlation analyses show freshwater flow, salinity, and chlorophyll *a* concentration were all significantly related to benthic species abundance

and persistence at one or more of the eastern sites (Table 8). Water temperature showed a positive relationship ($P < 0.05$) with benthic species abundance and persistence, but only for D28A-R. The significantly positive relationship between first ordination dimension CA results and chlorophyll *a* concentrations may be related to temporally matched seasonal increases in benthos abundance and phytoplankton biomass. Although the first dimension ordination patterns for the eastern sites differed from those for the other site groups, the same factors, namely salinity and freshwater flow, are thought to have had the largest effect on benthic species abundance and persistence at the eastern sites. Based on the limited change in CA scores for the eastern sites and the limited distribution among individual species scores, changes in salinity and freshwater flow appear to have influenced species abundance more than species persistence.




Figure 29
SPECIES SCORES, EASTERN STATIONS, D11, D19, AND D28A

Persistent and Dominant Species

A major finding from the correspondence analysis is that a relatively small number of numerically dominant and persistent species have a large influence over the composition of the benthos in the upper Sacramento-San Joaquin estuary. In her report, Markmann⁴⁶ discusses the life history and population patterns of the most numerous benthic organisms. She found that of the 140 benthic species identified from 1975 to 1981, only about 13 species typically comprised 10% or more of the community density at any one site. The suite of species that dominated the sampling area from 1980 to 1990 changed little from the suite of species that dominated from 1975 to 1981.

This section begins with a status review and update of the persistent and dominant species discussed by Markmann. A discussion of three exotic benthic organisms (*P. amurensis*, *H. hinumensis*, *Gammarus daiberi*) detected from 1980 to 1990 follows this review.

Limnodrilus hoffmeisteri and *Varichaetadrilus angustipenis*

These oligochaete worms are both in the family Tubificidae. In fact, until 1989, both species were classified as members of the genus *Limnodrilus*. These worms are able to withstand extreme environmental changes and can tolerate polluted conditions and hypoxic sediments.⁴⁷ Temperature (primarily) and substrate composition (secondarily) have been shown to regulate reproduction and recruitment.^{48,49} Above temperatures of 15°C, breeding is continuous. Recruitment success is optimal in organically rich mud.⁵⁰ Crumb⁵¹ found a relationship between the annual temperature and population abundance

of *L. hoffmeisteri* in the Delaware River. Abundances were highest during the spring, when temperatures ranged from 20–25°C at sites with high concentrations of organic mud.

In this estuary, either or both *L. hoffmeisteri* and *V. angustipenis* were among the four numerically dominant species at every monitoring site from 1980 to 1990 (Appendix B). Abundances were variable within and among sites, but were often highest at D11-C. Markmann reported *Limnodrilus* spp. as the most numerous organism at D7-C from 1975 to 1981. While *L. hoffmeisteri* maintained numerical dominance at D7-C through June 1983, it declined in abundance in 1984 and remained at lower levels thereafter (Appendix B).

The broad distribution of both *L. hoffmeisteri* and *V. angustipenis* within the sampling area is evidence of the robust nature of these species. Salinity levels often differ by an order of magnitude between Suisun Bay and the central delta. *L. hoffmeisteri* and *V. angustipenis* are among the few native benthic organisms that have maintained their numerical dominance and broad distribution throughout the existence of this monitoring program.

Corophium stimpsoni and *C. spinicorne*

Corophium spp. are native tube-building detritivorous amphipods most prevalent in areas with moderate levels of fine sediments and organic material and slightly brackish to fresh water.⁵² These amphipods are reported to be a food source for other arthropods, such as *Crangon franciscorum*, and several estuarine fishes such as striped bass, *Morone saxatilis*, and catfish, *Ictalurus* spp.⁵³

46 Markmann, 1986; cited.

47 RO Brinkhurst. 1972. *The Role of Sludge Worms in Eutrophication*. US Environmental Protection Agency, Ecol Res Serv EPA-R3-72-004. 68 pp.

48 Brinkhurst, 1972; cited.

49 CR Kennedy. 1966. The life history of *Limnodrilus hoffmeisteri* Clap. (Oligochaeta: Tubificidae) and its adaptive significance. *Oikos* 17:158-168.

50 Brinkhurst, 1972, and Kennedy, 1966; cited.

51 SE Crumb. 1977. Macroenthos of the tidal Delaware River between Trenton and Burlington, New Jersey. *Chesapeake Sci* 18:253-265.

52 Nichols and Pamatmat, 1988; cited.

53 Markmann, 1986; cited.

Historically these amphipods have been the numerically dominant benthic organism in many parts of the delta, often exceeding concentrations of 20,000 individuals per square meter.^{54,55,56} From 1980 to 1990, abundances appear to have varied seasonally, with peak concentrations occurring between summer and fall (Appendix B). However, in 1987 abundance of both *C. stimpsoni* and *C. spinicorne* declined sharply at all Sacramento River sites (D4-L, D4-R, D4-C) and remained at extremely low levels through 1990. Markmann suggested that specific conductance above 5,000 $\mu\text{S}/\text{cm}$ (2.8 ppt) may limit the occurrence of *C. stimpsoni*. An examination of the relationship between *C. stimpsoni* abundance and specific conductance at D11-C supports this hypothesis (Figure 30). In general, the abundance of *C. stimpsoni* was depressed when specific conductance exceeded 4,000 $\mu\text{S}/\text{cm}$ (2.3 ppt). Between 1987 and 1990, specific conductance at D4 exceeded 4,000 $\mu\text{S}/\text{cm}$ 68% of the time. These drought-associated increases in specific conductance (salinity) appear to have limited the occurrence of at least *C. stimpsoni* in the western delta and illustrate the effect physico-chemical changes can have on native benthic organisms.

Manayunkia speciosa

Manayunkia speciosa is a colonial tube-building polychaete worm commonly found in fresh water.⁵⁷ *M. speciosa* is hermaphroditic and reproduces sexually or asexually within its tube.⁵⁸ The tube is constructed of fine particles cemented together by a mucoid secretion.⁵⁹ The young mature in the parental tube and crawl out as small adults to form their own tube within the colony.⁶⁰




Figure 30
COROPHIUM STIMPSONI CONCENTRATIONS AND
MONTHLY SPECIFIC CONDUCTANCE AT D11-C,
1987-1990

Hazel and Kelley⁶¹ first reported the presence of *M. speciosa* along the West Coast from samples taken in the San Joaquin River and one location in Oregon. DWR benthic monitoring samples collected from 1975 to 1979 showed *M. speciosa* to exist at several locations in the interior delta at concentrations from 2,000 to 50,000 individuals per square meter.⁶² From 1975 to 1979, the greatest number of *M. speciosa* were found in the lower Mokelumne and San Joaquin rivers and at D28A on Old River.⁶³

- 54 CR Hazel and DW Kelley. 1966. Zoobenthos of the Sacramento-San Joaquin Delta. Pages 113-132 in *Ecological Studies of the Sacramento-San Joaquin Estuary, Part I, Zooplankton, Zoobenthos, and Fishes of San Pablo and Suisun Bays, Zooplankton and Zoobenthos of the Delta*. DW Kelley, editor. Department of Fish and Game, Fish Bulletin 136.
- 55 L Eng. 1975. *Biological Studies of the Delta-Mendota Canal, Central Valley Project, California II*. California Academy of Science, Contract 14-06-200-7762A. 178 pp.
- 56 Markmann, 1986; cited.
- 57 TP Poe and DC Stefan. 1974. Several environmental factors influencing the distribution of the freshwater polychaete, *Manayunkia speciosa* Leidy. *Chesapeake Sci* 15:235-237.
- 58 P Croskery. 1978. The freshwater co-occurrence of *Eurytemora affinis* (Copepoda: Calanoida) and *Manayunkia speciosa* (Annelida: Polychaeta): possible relics of a marine incursion. *Hydrobio* 59:237-241.
- 59 Poe and Stefan, 1974; cited.
- 60 Croskery, 1978, and Poe and Stefan, 1974; cited.
- 61 Hazel and Kelley, 1966; cited.
- 62 Markmann, 1986; cited.
- 63 Markmann, 1986; cited.

From 1980 to 1990, *M. speciosa* was numerically dominant at D19-C and D28A-R. Abundance at D19-C fluctuated between zero and 10,000 individuals per square meter from 1980 to 1985 but remained below 2,000 thereafter (Appendix B). Abundance at D28A-R was much more variable, ranging from zero to 36,000 individuals per square meter from 1980 to 1990. Abundance was highest during the extremely wet years of 1983 and 1986. Numerical dominance of *M. speciosa* at D19-C and D28A-R suggests this polychaete prefers freshwater habitats where the substrate is dominated by fine material.

Corbicula fluminea

The freshwater clam *Corbicula fluminea* was the most common benthic organism collected in the sampling area. This clam was introduced into California in the late 1940s and quickly became a dominant member of the benthos in the upper estuary.⁶⁴ From 1980 to 1990, *C. fluminea* was among the four numerically dominant organisms at all sampling sites except D7-C (Appendix B).

The ecology and biology of *C. fluminea* have been thoroughly studied by researchers throughout the world. Only a brief summary is presented here; refer to Mattice *et al*⁶⁵ for additional information. *C. fluminea* is a suspension-feeding clam that filters phytoplankton and organic detritus from the water column.⁶⁶ More recent studies suggest that, like *Potamocorbula amurensis*, high concentrations of *C. fluminea* are able to filter a significant portion of the phytoplankton from the water column.⁶⁷ Reproduction in the Sacramento-San Joaquin estuary is

thought to occur twice annually between spring and fall.^{68,69} Adult clams brood their larvae in a marsupium for about one month.⁷⁰ Larvae are released from the marsupium when temperatures exceed 15°C.⁷¹ Released larvae settle out within 48 hours.⁷²

Immature clams are readily dispersed to other parts of the estuary by flowing water.⁷³ *C. fluminea* have been collected at D7-C, but it is thought these clams are brought in during times of increased outflows.⁷⁴ Salinity levels in Suisun Bay prevent establishment of permanent populations.⁷⁵ Markmann⁷⁶ suggested *C. fluminea* populations in the central delta serve as recruitment pools for the western delta, where immature clams are transported downstream during high outflows in the spring. She believed higher salinity levels in the fall, followed by increased water velocities in winter and spring, prevent establishment of large, permanent populations of *C. fluminea* in the western delta. However, benthic monitoring data from 1980 to 1990 suggest established populations of *C. fluminea* do exist in the western delta. Although abundance in the western delta was generally lower than in the central delta, clams were continually collected at both D4 and D11 during years of extremely high outflow (1983 and 1986) and during drought years (1987 to 1990) when salinity levels increased in the western delta. In addition to the lower abundance of clams in the western delta, Winternitz⁷⁷ found the productivity of *C. fluminea* was lower in the western delta than in the central delta. The western delta is probably marginal habitat for *C. fluminea*, primarily due to the higher salinity levels.

64 DS Cherry, J Cairns, RL Graney. 1980. Asiatic clam invasion causes and effects. *Water Spectrum* Fall:19-24.

65 JS Mattice, LL Eng, BN Collier. 1979. *Corbicula 1979: A Bibliography*. Environmental Sciences Division, Oak Ridge National Laboratory. Publication 1315.

66 Eng. 1975; cited.

67 R Cohen, PV Dresler, E Phillips, R Cory. 1984. The effect of the Asiatic clam *Corbicula fluminea* on phytoplankton of the Potomac River, Maryland. *Limnol Oceanogr* 29:170-180.

68 L Eng. 1977. Population dynamics of the Asiatic clam, *Corbicula fluminea* (Muller), in the concrete-lined Delta-Mendota Canal of central California. Pages 40-68 in *Proc First Intl Corbicula Symp, October 13-15, 1977*.

69 Hazel and Kelley, 1966; Siegfried *et al*, 1978; Eng 1975; cited.

70 Eng, 1977; cited.

71 Crumb, 1977; cited.

72 PV Dresler and RL Cory. 1980. The Asiatic clam, *Corbicula fluminea* (Muller), in the tidal Potomac River, Maryland. *Estuaries* 3:150-151.

73 Eng, 1977; cited.

74 Hazel and Kelley, 1966; cited.

75 Hazel and Kelley, 1966; cited.

76 Markmann, 1986; cited.

77 L Winternitz. 1992. *Estimating Secondary Production Level of Corbicula fluminea in the Sacramento-San Joaquin Estuary*. Masters Thesis. University of San Francisco.

Potamocorbula amurensis

The Asian clam, *Potamocorbula amurensis*, first detected in this estuary in late 1986, is thought to have been introduced into Suisun Bay as larvae from ship ballast water.⁷⁸ This clam is native to estuaries along the east coast of Asia.⁷⁹ The abundance and distribution of *P. amurensis* has increased dramatically in the upper Sacramento-San Joaquin estuary, since it was first detected.^{80,81} By 1990 (four years after first detection), *P. amurensis* was well established in a variety of habitats throughout San Pablo and Suisun Bays, and Suisun Marsh, often at concentrations exceeding 1,000 clams per square meter.⁸²

Results from the correspondence analyses clearly show *P. amurensis* has altered the benthos at both D7 and D4. This clam has been a numerically dominant organism at both stations since 1988.⁸³ However, the persistently low salinity in the central delta has probably prevented the establishment of *P. amurensis* in this region. Although this clam is reported to be euryhaline,⁸⁴ laboratory observations of *P. amurensis* confirm that they cease all activity when exposed to freshwater and exhibit a high rate of mortality after several weeks of such exposure.⁸⁵

Trophic dynamics within the upper estuary have been altered by the introduction of the Asian clam. In particular, *P. amurensis* is known to have contributed to the substantial and sustained reductions in surface chlorophyll *a* concentrations in Suisun Bay.⁸⁶ This clam is a suspension feeder capable of consuming phytoplankton, bacterioplankton, particulate organic matter, and zooplankton nauplii.^{87,88} Although the establishment of *P. amurensis* may

have increased the competition between other benthic organisms for space and food, it does provide a new and abundant food source for bottom feeding birds, fish, and crabs.⁸⁹

Gammarus daiberi

The amphipod *Gammarus daiberi* is endemic to much of the Atlantic coast, commonly occurring in estuaries and sounds from New York through South Carolina.⁹⁰ Populations reach highest concentrations during spring and summer in salinities of 1-5 ppt; however, individuals do occur seaward to salinities of 15 ppt. Gammarid species are typically macrophagous and free-swimming. *G. daiberi* is pelagic, occurring in mid- to near-bottom depths, but may also reside epibenthically. The species co-occurs with *G. fasciatus* and *G. tigrinus* in tidal areas of fresh and brackish water. The life history and habitat requirements of *G. daiberi* have not been studied in its Pacific coast setting, but they are presumed to be similar to its native ecology.

G. daiberi was first detected in this estuary in 1983 from benthic samples collected in the central delta. Between 1983 and 1986 the amphipod was collected only rarely. Beginning in 1986, however, appreciable concentrations of *G. daiberi* were collected in both benthic and zooplankton monitoring samples. Since 1986, *G. daiberi* has been routinely collected in the central and western delta regions, and in Suisun Bay. Abundance fluctuates seasonally, with highest concentrations typically occurring in spring and early summer. Because of this amphipod's mobility, estimates of benthic concentrations are subject to

-
- 78 JT Carlton, JK Thompson, LE Schemel, FH Nichols. 1990. The remarkable invasion of San Francisco Bay (California, USA) by the Asian clam *Potamocorbula amurensis*. I. Introduction and dispersal. *Mar Ecol Prog Ser* 66:81-94.
- 79 Carlton *et al*, 1990; cited.
- 80 Carlton *et al*, 1990; cited.
- 81 ZP Hymanson. 1992. *Results of a Spatially Intensive Survey for Potamocorbula amurensis in the Upper San Francisco Bay Estuary*. Interagency Ecological Studies Program, Technical Report 30. Department of Water Resources.
- 82 Hymanson, 1992; cited.
- 83 DWR, 1992; cited.
- 84 Carlton *et al*, 1990; cited.
- 85 F Nichols, US Geological Survey, Palo Alto, CA; personal communication. 1993.
- 86 Alpine and Cloern, 1992; cited.
- 87 I Werner and JT Hollibaugh. *Potamocorbula amurensis* (Mollusca, Pelecypoda): Comparison of clearance rates and assimilation efficiencies for phytoplankton and bacterioplankton. *Limnology and Oceanography* 38:949-964. 1993.
- 88 Kimmerer, personal communication.
- 89 Alpine and Cloern 1992, Carlton *et al* 1990, Nichols *et al* 1990; all cited.
- 90 EL Bousfield. *Shallow-Water Gammaridean Amphipoda of New England*. Comstock Publisher Associates, Ithaca, NY. 1973.

considerable error. However, benthic and zooplankton monitoring show *G. daiberi* is well established throughout much of the upper estuary. This amphipod is known to serve as a food source for young striped bass.⁹¹

Hemileucon hinumensis

The cumacean crustacean *Hemileucon hinumensis* was first detected in Suisun Bay in 1986. Little is apparently known about the ecology of this organism. No information on this species' ecology or life history was found in a search of recent literature. The abundance and distribution of *H. hinumensis* increased between 1986 and 1990, a period coincident with increased abundance and distribution of other exotic species, such as *Potamocorbula amurensis* and *Gammurus daiberi*. The coincident appearance and establishment of these introduced species, suggests the presence of related ecological requirements that probably originated during the drought. Research efforts into the resulting interactions and effects of *H. hinumensis* on the resident benthic community may provide new clues to the species' ecology and role in the benthic community.

Trends in the Benthos and Water Project Operations

Markmann⁹² concluded that water project operations could affect the benthos of the upper estuary through changes in seasonal salinity patterns and localized changes in water velocity and sediment dynamics. It is clear that seasonal salinity patterns do affect the benthos of the upper estuary. However, this summary analysis showed that from 1980 through 1990 most of the substantial variability in the benthos was due to longer-term (drought and flood mediated) changes in salinity.

Determination of water project related impacts on the benthos was not included as a specific criterion in the design of this monitoring program. Such a criterion would require a substantially different program design. The hydrology of the upper estuary is very complex, while water project operations are both spatially and temporally variable. We think focused modeling and field studies are required to

determine if and what effects water project operations are having on the benthos of the upper estuary.

Detectability and Sensitivity Analysis of the Benthic Monitoring Program

Further analyses of the benthic data collected from 1980 to 1990 were completed to determine the ability of the existing monitoring program to detect changes in benthic community structure. Our approach in this section is to answer three questions that are key to the design and implementation of a benthic monitoring program with current objectives to:

- Meet the monitoring obligations described in Water Rights Decision 1485.
- Monitor trends in the abundance and distribution of benthic fauna.
- In conjunction with other monitoring data, determine what environmental factors (including water project operations) are responsible for the trends in abundance and distribution of benthic fauna.

What should the sampling frequency be?

Data collected from 1980 to 1990 were analyzed to determine the variance structure of organism abundance at the eight benthic sampling sites. Coefficients of variation

$$CV = [(standard\ deviation / mean)100]$$

were used as a standardized measure of variance. CVs were calculated using total community abundance values over three time intervals: month, season (3 months), and year. Results show total abundance was highly variable at all sites over all time periods (Table 9). Within each period, the CVs among sites were similar, however, demonstrating that the magnitude of variation is similar throughout the sampling region.

CV results for total community abundance were also used to generate power curves. These curves show the number of samples needed to detect various levels of change in community abundance (expressed as percentages) on a yearly, seasonal, and monthly basis. Currently, the monitoring program

91 L Miller, Department of Fish and Game, Stockton, CA; personal communication, 1993.

92 Markmann, 1986; cited.

Table 9
COEFFICIENTS OF VARIATION FOR
TOTAL COMMUNITY ABUNDANCE VALUES AND FOR
CORBICULA FLUMINEA ABUNDANCE VALUES
CALCULATED FOR DIFFERENT SITES AND TIME INTERVALS

Total Community Abundance			
Site	Year CV	Season CV	Month CV
D7-C	246	286	302
D11-C	283	286	304
D4-R	253	247	313
D4-L	291	286	306
D4-C	264	340	400
D19-C	264	226	238
D28A-R	267	279	289
D28A-L	347	272	331
Corbicula Fluminea Abundance			
Site	Year CV	Season CV	Month CV
D7-C	79.8	160	222
D11-C	78.4	103	113
D4-R	114	101	125
D4-L	115	139	142
D4-C	120	152	158
D19-C	69.1	93.5	108
D28A-R	82.1	138	140
D28A-L	129	135	173

collects three replicate samples at each site, each month. Thus, nine samples are collected during a season and 36 samples are collected at each site during a year. Results from the power curve analysis for D7-C show that at the current sampling frequency the monitoring program is able to detect a 55% change in total abundance between years but is not able to accurately detect quantitative changes between months or seasons (Figure 31). Results of power curve analyses for other sites (not shown) were similar because of the similarity in CV results.

CVs were also calculated for several prominent benthic organisms. Results for *Corbicula fluminea*, the most prominent organism throughout the sampling region from 1980 to 1990, are shown in Table 9 and exemplify results for other species tested. Although substantial variability in the abundance of a single species also exists, CVs for all sites and time periods were lower for *C. fluminea* than for the total community abundance values (Table 9). This suggests the monitoring program is better able to detect abundance changes in prominent species. However, power curve analyses show increased sensitivity in detecting the change in abundance of a single species

is limited to the yearly and seasonal time period. For example, results from Site D19-C, where variation in *C. fluminea* was lowest for all periods, show the current level of sampling is able to detect somewhat less than a 30% difference in abundance between years and about a 50% difference between seasons. No quantitative difference could be detected between months (Figure 32).

As an alternative to monitoring the abundance and distribution of all macrobenthic organisms, the program could be structured to monitor only the abundance and distribution of dominant species. Fluctuations in abundance of the more persistent organisms, which tend to be lower, would allow for a reduction in sampling effort. However, any reduction in sampling frequency would compromise the ability to detect seasonal or annual abundance changes and other basic life-history information and further reduce the ability to characterize abundance and distribution trends in less persistent but ecologically important species.


What should the sample replication be?

Altering the number of sample replicates is another way to change the level of sensitivity and detectable difference. Currently, three replicate samples are collected monthly at each site. This is the minimum required to obtain quantitative monthly abundance estimates, because of the high variability in abundance. As discussed, power of detection curves show the benthic monitoring program is at the lower limits of detection on a monthly or seasonal time scale and at the mid-detection level on an annual scale.

Any increase in the number of replicates could increase the detection levels at all time scales. However, a substantial increase in the number of replicates would be required if the sampling frequency were less than monthly, because sensitivity of the monitoring program is based on the total number of samples collected at a site. For example, suppose sampling frequency were reduced from monthly to quarterly and the number of replicates remained the same. This would reduce the total number of samples collected annually at each site from 36 to 12. From the results in Figure 31 it can be seen that this reduced sampling frequency would only provide the ability to detect a 90% difference in total community abundance between years. On the other hand, maintaining the same level of detectability


YEAR

Percent Detectable Difference	Number of Samples
10	1012
20	253
30	112
40	63
50	40
60	28
70	20
80	16
90	13
100	10



SEASON

Percent Detectable Difference	Number of Samples
10	1598
20	400
30	178
40	81
50	52
60	45
70	33
80	25
90	20
100	16



MONTH

Percent Detectable Difference	Number of Samples
10	1522
20	381
30	169
40	95
50	61
60	42
70	31
80	24
90	19
100	15





Figure 31
POWER OF DETECTION CURVES FOR TOTAL COMMUNITY ABUNDANCE AT D7-C


YEAR

Percent Detectable Difference	Number of Samples
10	97
30	11
50	4
70	3
100	1



SEASON

Percent Detectable Difference	Number of Samples
10	176
30	20
50	7
70	4
100	2



MONTH

Percent Detectable Difference	Number of Samples
10	217
30	24
50	9
70	6
100	3




Figure 32
POWER OF DETECTION CURVES FOR *CORBICULA FLUMINEA* AT D19-C

between years (*ie*, about 55%) would require collection of nine replicate samples from each site each quarter. Thus, while detecting differences in abundance between months or seasons is not a primary objective, collecting three replicate samples each month provides a level of effort consistent with all objectives. In addition, no amount of increase in sample replication could replace the loss of life history information if the sampling frequency were less than monthly. Collecting samples on at least a monthly basis is essential to monitoring recruitment events.

What should the spatial distribution of stations be?

A variety of procedures can be used to determine the location of benthic sampling stations. One simple and objective way is to choose locations where the maximum chance of collecting a species occurs. Presumably, the presence and persistence of a benthic organism at any one location is based on the characteristics and variability of the local environment. The presence or absence of a species and subsequent changes in abundance are a reflection of changes in the environment. By choosing to sample sites with maximum species diversity, we increase the chance of detecting the response of benthic organisms to changes in the environment. However, detection of new introductions is an exception to this line of reasoning, since we have no idea which organisms are going to be introduced or where they will be introduced.

Correspondence analyses of all sites together suggest the existing sites fall into one of three groups based on benthic species abundance and persistence. The results are fairly intuitive, given the spatial distribution of the sites, and they provide an objective basis for making comparisons among sites.

Since 1980, D7-C has been the only site sampled in the Suisun Bay region. Due to the large fluctuations in salinity and the recent establishment and dominance of two exotic organisms, *Potamocorbula amurensis* and *Hemileucon hinumensis*, this region is distinct from the two delta regions sampled. D7-C is, by default, the best site to sample in the future, because it is the only site in this region.

The second region consists of three sites in the lower Sacramento River: D4-R, D4-L, and D4-C. Together these sites form a transect across the river where samples are taken from the right and left banks and center channel. Comparisons of species occurrence at these sites show more species have been found on either bank than in the center of the channel (Figure 33). This is probably due to the high degree of scouring and substrate instability in the center of the channel. Results from the bank sites show sampling the left bank of the Sacramento River (D4L) provides the best opportunity for detection of a species in this region.

The third benthic sampling region includes one site in the western delta (D11-C) and three sites in the central delta (D19-C, D28A-R, D28A-L) (Figure 33). Comparisons of species occurrence among these sites show more species have been found in the banks of Old River (D28A-L, D28A-R) than in either of the lacustrine sites (D11-C, D19-C). Of the two sites on Old River, D28A-L provides a slightly better




Figure 33
SIMILARITY IN BENTHIC SPECIES COMPOSITION AMONG SITES IN TWO DELTA REGIONS

chance of collecting a higher diversity of benthic species.

These results are limited by the relatively few sites sampled in each region or, in many cases, the complete lack of sites in a region. Ideally, one or more sites should be sampled in each environmentally

distinct region within the delta and Suisun Bay. This distribution of sampling effort would yield more ecologically relevant information necessary in determining environmental factors that regulate the abundance and distribution of benthic organisms in the upper estuary.

LITERATURE PERTAINING TO THE BENTHOS OF THE SACRAMENTO-SAN JOAQUIN ESTUARY

- Aplin, J.A. 1967. Biological survey of San Francisco Bay: 1963-66. California Department of Fish and Game, Marine Resources Operations Laboratory. 131 pp.
- Anatec Laboratories, Inc. 1980a. Point Richmond outfall study, predischARGE monitoring phase. 78 pp. plus appendices.
- Anatec Laboratories, Inc. 1980b. San Pablo Bay field studies: Phase I, Novato Sanitary Component. Volume I. 78 pp. plus appendices.
- Anatec Laboratories, Inc. 1980c. San Pablo Bay field studies: Central Marin component. Volume I. 58 pp. plus appendices.
- Anatec Laboratories, Inc. 1980d. San Pablo Bay field studies: Petaluma River biological survey. 48 pp. plus appendices B and C.
- Anatec Laboratories, Inc. and Kinnetic Laboratories, Inc. 1982. East Bay Municipal Utility District local effects monitoring program: Final report, volume 3, Biology, Part I, benthic studies. 137 pp. plus appendixes.
- Brinkhurst, R.O., and Simmons, M.L. 1968. The aquatic Oligochaeta of the San Francisco Bay system. California Department of Fish and Game 54:180-194.
- California Department of Fish and Game. 1968. San Francisco Bay-Delta water quality control program; Task VII-1B (DFG). Fish and wildlife resources of San Francisco Bay and Delta-description, environmental requirements, problems, opportunities and the future. California State Water Quality Control Board. 338 pp.
- California Department of Public Health. 1954. Department of Fish and Game report, part 1 of Richmond shoreline investigation: California Department Public Health Report. Project number 54-2-3. 84 pp.
- CH₂M Hill, 1982. Equivalent protection study, final report to Chevron, USA.
- Daniel, D.A., and H.K. Chadwick. 1971. Effects of selected waste discharges on benthic invertebrate communities: Volume VII of a study of toxicity and biostimulation in San Francisco Bay-Delta waters. California Department of Fish and Game.
- Eng. L.L. 1975. Biological studies of the Delta-Mendota Canal, Central Valley Project, California, II. California Academy of Sciences. Contract No. 14-06-200-7762A. 178 pp.
- Ferreira, R.F., and D.B. Green. 1977. Distribution and abundance of benthic organisms in the Sacramento River, California. United States Geological Survey. USGS/WRI-77-60. 24 pp.
- Filice, F.P. 1954a. An ecological survey of the Castro Creek area in San Pablo Bay. Wasmann Jor. Biol. 12(1):1-24.
- Filice, F.P. 1954b. A study of some factors effecting the bottom fauna of a portion of the San Francisco Bay estuary. Wasmann Jor. Biology 12(3):257-292.
- Filice, F.P. 1958. Invertebrates from the estuarine portion of San Francisco Bay and some factors influencing their distribution. Wasmann Jor. Biology 16:159-211.
- Fisk, L.O., T.R. Doyle, G.E. Reiner, and D.C. Joseph. 1962. Sacramento River water pollution survey, Appendix D: Benthic biology. Department of Water Resources. Bulletin 111. 78 pp.

- Grieb, T.M. 1990. Use of ODES to evaluate San Francisco Bay data: 1986 USGS benthic infauna pilot survey data. TetraTech, Inc.
- Hazel, C.R., and D.W. Kelley. 1966. Zoobenthos of the Sacramento-San Joaquin Delta. In D.W. Kelley [Ed.]. Ecological studies of the Sacramento-San Joaquin Estuary, Part I. zooplankton, zoobenthos, and fishes of San Pablo and Suisun Bays, zooplankton and zoobenthos of the Delta. California Department of Fish and Game. Fish Bulletin 136:113-132.
- Herbold, B., and P.B. Moyle. 1989. The ecology of the Sacramento-San Joaquin Delta: a community profile. U.S. Fish Wildl. Serv. Biol. Rep. 85(7.22). 106 pp.
- Hopkins, D.R. 1986. Atlas of the distributions and abundances of common benthic species in San Francisco Bay, California. United States Geological Survey. Water Resources Investigations Report 86-4003. 228 pp.
- Hopkins, D.R. 1987. Temporal variations in the benthic communities at four intertidal sites in San Francisco Bay, California, 1983-85. United States Geological Survey. Open-File Report 81-132. 107 pp.
- Hymanson, Z.P. 1991. Results of a spatially intensive survey for *Potamocorbula amurensis* in the upper San Francisco Bay estuary. Interagency Ecological Studies Program. Technical report 30. 21 pp.
- Jones, M.L. 1961. A quantitative evaluation of the benthic fauna off Point Richmond, California. California University Pubs. Zoology 67:219-320.
- Kaiser Engineers. 1968. San Francisco Bay-Delta water quality control program; Task VII-1b (KE), biologic-ecologic study: California State Water Quality Control Board, p. I1 to VI-4, with appendices.
- Kaiser Engineers. 1969. San Francisco Bay-Delta water quality control program; final report, prelim. ed., to the state of California: California State Water Quality Control Board, p.I-1 to XXIII-17, with appendices.
- Kinney, P.J. 1981. Benthic community analysis. East Bay Dischargers Authority. Prepared by Kinetics Laboratories, Inc. KLI-81-11. 119 pp.
- Liu, D.H.W., K.D. Martin, and C.R. Norwood. 1975. Dredge disposal study, San Francisco Bay and estuary Appendix D biological community. United States Army Engineer District, San Francisco Corps of Engineers. DACW 07-74-C-005. 244 pp.
- Markmann, C. 1986. Benthic monitoring in the Sacramento-San Joaquin Delta: Results from 1975 through 1981. Interagency Ecological Studies Program. Technical report 12. 55 pp.
- McAllister, R.D., and Moore, T.O., Jr. 1982. Selected shellfish resources of San Francisco Bay: their distribution, abundance, use, public access, and recommended management alternatives: San Francisco Bay Regional Water Quality Control Board. 168 pp.
- Nichols, F.H. 1973. A review of benthic faunal surveys in San Francisco Bay. United States Geological Survey. Geological Survey Circular 677. 20 pp.
- Nichols, F.H. 1979. Natural and anthropogenic influences on benthic community structure in San Francisco Bay. In: Conomos, T.J. [Ed.]. San Francisco Bay: The Urbanized Estuary. Pacific Division AAAS, San Francisco. Pages 409-426.
- Nichols, F.H. 1984. Significance of long-term variations in estuarine benthos. United States Geological Survey. Circular 938:91-97.
- Nichols, F.H. 1985a. Abundance fluctuations among benthic invertebrates in two Pacific estuaries. Estuaries 8:136-144.
- Nichols, F.H. 1985b. Increased benthic grazing: an alternative explanation for low phytoplankton biomass in Northern San Francisco Bay during the 1976-77 drought. Estuarine Coastal and Shelf Science 21:379-388.
- Nichols, F.H., and J.K. Thompson. 1985. Time scales of change in the San Francisco Bay benthos. Hydrobiologia 129:121-138.

- Nichols, F.H. 1987. Benthic ecology and heavy metal accumulation. In: Goodrich, D.M. [Ed.]. San Francisco Bay issues, resources, status, and management. National Oceanic and Atmospheric Administration. Estuary of the Month Seminar Series 6:65-68.
- Nichols, F.H. and M.M. Pamatmat. 1988. The ecology of the soft-bottom benthos of San Francisco Bay: a community profile. U.S. Fish Wildl. Serv. Biol. Rep. 85(7.19). 73 pp.
- Pacific Environmental Laboratory. 1976a. Summary data sheets for macroinvertebrates: California Regional Water Quality Control board, order number 74-112, NPDES permit number CA0037729, waste discharge monitoring for the city of Richmond, California.
- Pacific Environmental Laboratory. 1976b. Summary data sheets for macroinvertebrates: California Regional Water Quality Control board, NPDES permit number CA0005789, waste discharge requirements for Shell Oil Company, Martinez, Contra Costa County, California.
- Packard, E.L. 1981a. Molluscan fauna from San Francisco Bay. California University Pubs. Zoology. 14:199-452.
- Packard, E.L. 1981b. A quantitative analysis of the molluscan fauna of San Francisco Bay. California University Pubs. Zoology. 18:299-336.
- Painter, R.E. 1966. Zoobenthos of San Pablo and Suisun Bays. In D.W. Kelley [Ed.]. Ecological studies of the Sacramento-San Joaquin Estuary, Part I. zooplankton, zoobenthos, and fishes of San Pablo and Suisun Bays, zooplankton and zoobenthos of the Delta. California Department of Fish and Game. Fish Bulletin 136:40-55.
- Pearson, E.A., P.N. Storrs, and R.E. Selleck. 1970. A comprehensive study of San Francisco Bay, final report; VIII, Summary, conclusions and recommendations. California University (Berkeley) Sanitary Engineering Research Laboratory Report. 67(5)1-85.
- Radtke, L.D. 1966. Distribution of smelt, juvenile sturgeon, and starry flounder in the Sacramento-San Joaquin Delta, with observations on food of sturgeon. In D.W. Kelley [Ed.]. Ecological studies of the Sacramento-San Joaquin Estuary, Part II. zooplankton, zoobenthos, and fishes of San Pablo and Suisun Bays, zooplankton and zoobenthos of the Delta. California Department of Fish and Game. Fish Bulletin 136:115-126.
- San Francisco Bay Marine Research Center, 1976. Summary data sheets for macroinvertebrates collected: California Regional Water Quality Control Board, order number 74-151, waste discharge requirements for Standard Oil Company of California, Richmond Refinery, Contra Costa County.
- San Francisco Bay Marine Research Center. 1976. Summary data sheets for macroinvertebrates collected: California Regional Water Quality Control Board, order number 74-152, NPDES permits number CA 0005053, waste discharge requirements for Union Oil Company of California, San Francisco Refinery, Rodeo, Contra Costa County.
- Schemel, L.E., A.Y. Ota, J.G. Harmon, J.M. Shay, and R.N. Adorador. 1988. Benthic macrofauna and ancillary data for San Francisco Bay, California, March to November 1987. United States Geological Survey. Open-File Report 88-192. 73 pp.
- Schemel, L.E., A.Y. Ota, J.G. Harmon, J.M. Shay, and R.N. Adorador. 1990. Benthic macrofauna and ancillary data for San Francisco Bay, California, January to November 1988. United States Geological Survey. Open-File Report 89-596. 65 pp.
- Siegfried, C.A., A.W. Knight, and M.E. Kopache. 1980. Ecological studies on the western Sacramento-San Joaquin Delta during a dry year. Dept. Water Sci. Eng. paper 4506. 121 pp.
- Skinner, J.E. 1962. An historical review of the fish and wildlife resources of the San Francisco Bay area. California Department of Fish and Game, Water Project Branch Report. 1:1-226.
- Storrs, P.N., R.E. Selleck, and E.A. Pearson. 1963. A comprehensive study of San Francisco Bay, 1961-62. University of California (Berkeley), Sanitary Engineering Research Laboratory Report. 63(3)1-221.

- Storrs, P.N., R.E. Selleck, and E.A. Pearson. 1964. A comprehensive study of San Francisco Bay, 1962-63. University of California (Berkeley), Sanitary Engineering Research Laboratory Report, Volume 63, number 4, pages I-1 to IX-9.
- Storrs, P.N., R.E. Selleck, and E.A. Pearson. 1965. A comprehensive study of San Francisco Bay, 1963-64. University of California (Berkeley), Sanitary Engineering Research Laboratory Report, Volume 65, number 1, pages I-1 to VII-7.
- Storrs, P.N., E.A. Pearson, and R.E. Selleck. 1966a. A comprehensive study of San Francisco Bay, final report; II, Biological sampling and analytical methods. California University (Berkeley), Sanitary Engineering Research Laboratory Report 65(8)1-75.
- Storrs, P.N., E.A. Pearson, and R.E. Selleck. 1966b. A comprehensive study of San Francisco Bay, final report; V, Summary of physical, chemical, and biological water and sediment data. California University (Berkeley), Sanitary Engineering Research Laboratory Report 67(2)1-140.
- Thompson, J.K., and F.H. Nichols. 1981. Benthic macrofaunal biomass of San Francisco Bay, California: January, February and August, 1973. United States Geological Survey. Open-File Report 81-1331. 39 pp.
- Underwater Biological Research. 1978. Survey of benthic macrofauna at the San Pablo bay dredge disposal site July 1977-April 1978, final report. In: U.S. Army Corps of Engineers. Dredge disposal study, San Francisco Bay and Estuary. Appendix N, Addendum. 66 pp. plus appendices.
- U.S. Army Corps of Engineers. 1975. Dredge disposal study, San Francisco Bay and Estuary, Appendix D, Biological community study. 244 pp.

Appendix A

**Taxonomic List of Benthic Macrofauna
Identified from Monitoring Samples**

Taxonomic List of Benthic Macrofauna Identified from Monitoring Samples

ORG NUM	PHYLUM	FAMILY	GENUS	SPECIES	FULL TAXONOMIC NAME
000	NONE	NONE	NONE	NONE	NOTHING IN SAMPLE
100	CNID	HYDR	HYDR	SP A	CNIDARIA HYDRIDAE HYDRA SPECIES A
101	PLAT	PLAN	DUGE	TIGR	PLATYHELMINTHES PLANARIIDAE DUGESIA TIGRINA
103	PLAT	UNKN	TRIC	SP A	PLATYHELMINTHES UNKNOWN UID TRICLAD SPECIES A
105	PLAT	UNKN	TRIC	SP B	PLATYHELMINTHES UNKNOWN UID TRICLAD SPECIES B
107	PLAT	UNKN	TRIC	SP C	PLATYHELMINTHES UNKNOWN UID TRICLAD SPECIES C
109	PLAT	UNKN	TURB	SP A	PLATYHELMINTHES UNKNOWN MICROTURBELLARIAN SPECIES A
111	NEME	UNKN	PALE	SP A	NEMERTEA UNKNOWN UID PALEONEMERTEAN SPECIES A
113	NEME	TERT	PROS	GRAE	NEMERTEA TERTASTEMATIDAE PROSTOMA GRAECENSE
115	NEME	UNKN	NEME	SP A	NEMERTEA UNKNOWN UID NEMERTEAN SPECIES A
121	NEMA	PLEC	TERA	SP A	NEMATODA PLECTIDAE TERATOCEPHALUS SPECIES A
123	NEMA	DORY	DORY	SP A	NEMATODA DORYLAIMIDAE DORYLAIMUS SPECIES A
125	NEMA	DORY	EUDO	SP A	NEMATODA DORYLAIMIDAE EUDORYLAIMUS SPECIES A
127	NEMA	DORY	ACTI	SP A	NEMATODA DORYLAIMIDAE UID ACTINOLAIMINAE SPECIES A
129	NEMA	UNKN	NEMA	SP A	NEMATODA UNKNOWN UID NEMATODA SPECIES A
201	ANNE	NAID	BRAT	BILO	ANNELIDA NAIDIDAE BRATISLAVIA BILONGATA
203	ANNE	NAID	CHAE	DIAP	ANNELIDA NAIDIDAE CHAETOGASTER DIAPHANUS
205	ANNE	NAID	CHAE	LIMI	ANNELIDA NAIDIDAE CHAETOGASTER LIMNAEI
207	ANNE	NAID	DERO	DIGI	ANNELIDA NAIDIDAE DERO DIGITATA
209	ANNE	NAID	DERO	TRIF	ANNELIDA NAIDIDAE DERO TRIFIDA
211	ANNE	NAID	NAIS	COMM	ANNELIDA NAIDIDAE NAIS COMMUNIS/VARIABILIS
213	ANNE	NAID	NAIS	PARD	ANNELIDA NAIDIDAE NAIS PARDALIS
215	ANNE	NAID	NAIS	PSEU	ANNELIDA NAIDIDAE NAIS PSEUDOBTUSA
217	ANNE	NAID	NAIS	SIMP	ANNELIDA NAIDIDAE NAIS SIMPLEX
219	ANNE	NAID	OPHI	SERP	ANNELIDA NAIDIDAE OPHIDONAIIS SERPENTINA
221	ANNE	NAID	PARN	FRIC	ANNELIDA NAIDIDAE PARANAIIS FRICI
223	ANNE	NAID	PRIS	BREV	ANNELIDA NAIDIDAE PRISTINA BREVISETA
225	ANNE	NAID	PRIS	LONS	ANNELIDA NAIDIDAE PRISTINA LONGISETA
227	ANNE	NAID	SLAV	APPE	ANNELIDA NAIDIDAE SLAVINA APPENDICULATA
229	ANNE	NAID	STYL	LACU	ANNELIDA NAIDIDAE STYLARIA LACUSTRIS
231	ANNE	NAID	VEJD	COMA	ANNELIDA NAIDIDAE VEJDOVSKYELLA COMATA
233	ANNE	NAID	VEJD	INTE	ANNELIDA NAIDIDAE VEJDOVSKYELLA INTERMEDIA
241	ANNE	TUBI	AULO	LIMM	ANNELIDA TUBIFICIDAE AULODRILUS LIMNOBIUS
243	ANNE	TUBI	AULO	PIGU	ANNELIDA TUBIFICIDAE AULODRILUS PIGUETI
245	ANNE	TUBI	AULO	PLUR	ANNELIDA TUBIFICIDAE AULODRILUS PLURISETA
247	ANNE	TUBI	BOTH	VEJO	ANNELIDA TUBIFICIDAE BOTHRIONEURUM VEJDOVSKYANUM
249	ANNE	TUBI	BRAN	SOWE	ANNELIDA TUBIFICIDAE BRANCHIURA SOWERBYI
251	ANNE	TUBI	ILYO	FRAT	ANNELIDA TUBIFICIDAE ILYODRILUS FRANTZI CAPILLATUS
253	ANNE	TUBI	ILYO	TEMP	ANNELIDA TUBIFICIDAE ILYODRILUS TEMPLETONI
255	ANNE	TUBI	LIMO	HOFF	ANNELIDA TUBIFICIDAE LIMNODRILUS HOFFMEISTERI
257	ANNE	TUBI	LIMO	UDEK	ANNELIDA TUBIFICIDAE LIMNODRILUS UDEKEMIANUS
259	ANNE	TUBI	POTX	BAVA	ANNELIDA TUBIFICIDAE POTAMOTHRIX BAVARICUS
261	ANNE	TUBI	PSAM	CALF	ANNELIDA TUBIFICIDAE PSAMMORYCTIDES CALIFORNIANUS
263	ANNE	TUBI	QUIS	MULT	ANNELIDA TUBIFICIDAE QUISTADRILUS MULTISETOSUS
265	ANNE	TUBI	SPIR	FERO	ANNELIDA TUBIFICIDAE SPIROSPERMA FEROX
267	ANNE	TUBI	TENE	MAST	ANNELIDA TUBIFICIDAE TENERIDRILUS MASTIX

ORG NUM	PHYLUM	FAMILY	GENUS	SPECIES	FULL TAXONOMIC NAME
269	ANNE	TUBI	TUBF	BROW	ANNELIDA TUBIFICIDAE TUBIFICOIDES BROWNAE
271	ANNE	TUBI	TUBF	FRAS	ANNELIDA TUBIFICIDAE TUBIFICOIDES FRASERI
272	ANNE	TUBI	TUBF	SP A	ANNELIDA TUBIFICIDAE TUBIFICOIDES SPECIES A
273	ANNE	TUBI	VARI	ANGU	ANNELIDA TUBIFICIDAE VARICHAETADRILUS ANGUSTIPENIS
275	ANNE	TUBI	TUBD	SP A	ANNELIDA TUBIFICIDAE UID TUBIFICID SPECIES A
277	ANNE	ENCH	ENCH	SP A	ANNELIDA ENCHYTRAEIDAE UID ENCHYTRAEID SPECIES A
279	ANNE	ENCH	ENCH	SP B	ANNELIDA ENCHYTRAEIDAE UID ENCHYTRAEID SPECIES B
281	ANNE	GLOS	SPAR	EISE	ANNELIDA GLOSSOSCOLECIDAE SPARGANOPHILUS EISENI
283	ANNE	MEGA	MEGA	SP A	ANNELIDA MEGASCOLECIDAE UID MEGASCOLECID SPECIES A
285	ANNE	LUMB	LUBR	VARE	ANNELIDA LUMBRICULIDAE LUMBRICULUS VARIEGATUS
287	ANNE	LUMB	LUBR	SP A	ANNELIDA LUMBRICULIDAE LUMBRICULUS SPECIES A
289	ANNE	BRAN	CAMB	SP A	ANNELIDA BRANCHIOBDELLIDAE CAMBARINCOLA SPECIES A
291	ANNE	GLOP	HELO	STAG	ANNELIDA GLOSSIPHONIIDAE HELOBDELLA STAGNALIS
293	ANNE	GLOP	HELO	TRIS	ANNELIDA GLOSSIPHONIIDAE HELOBDELLA TRISERIALIS
295	ANNE	GLOP	PLAC	MONT	ANNELIDA GLOSSIPHONIIDAE PLACOBDELLA MONTANA
297	ANNE	ERPO	MOOR	MICR	ANNELIDA ERPOBDELLIDAE MOOREOBDELLA MICROSTOMA
301	ANNE	LUMN	LUMN	SP A	ANNELIDA LUMBRINERIDAE LUMBRINERIS SPECIES A
303	ANNE	ORBI	HAPL	ELON	ANNELIDA ORBINIIDAE HAPLOSCOLOPLOS ELONGATUS
305	ANNE	SPIO	BOCC	LIGE	ANNELIDA SPIONIDAE BOCCARDIA LIGERICA
307	ANNE	SPIO	POLD	LIGN	ANNELIDA SPIONIDAE POLYDORA LIGNI
309	ANNE	SPIO	PSED	KEMP	ANNELIDA SPIONIDAE PSEUDOPOLYDORA KEMPI
311	ANNE	SPIO	STRE	BENE	ANNELIDA SPIONIDAE STREBLOSPIO BENEDICTI
313	ANNE	SPIO	SPIO	SP A	ANNELIDA SPIONIDAE UID SPIONID SPECIES A
315	ANNE	CIRR	CIRR	SPIR	ANNELIDA CIRRHATULIDAE CIRRHIFORMIA SPIRABRANCHA
317	ANNE	CAPI	HETE	FILI	ANNELIDA CAPITELLIDAE HETEROMASTUS FILIFORMIS
319	ANNE	MALD	ASYC	ELON	ANNELIDA MALDANIDAE ASYCHIS ELONGATA
321	ANNE	NEPH	NEPH	CAEC	ANNELIDA NEPHTYIDAE NEPHTYS CAECOIDES
323	ANNE	NEPH	NEPH	CORN	ANNELIDA NEPHTYIDAE NEPHTYS CORNUTA FRANCISCANA
325	ANNE	PHYL	ETEO	CALI	ANNELIDA PHYLLODOCIDAE ETEONE CALIFORNICA
327	ANNE	PHYL	ETEO	LIGH	ANNELIDA PHYLLODOCIDAE ETEONE LIGHTI
329	ANNE	SYLL	SPHR	CALI	ANNELIDA SYLLIDAE SPHAEROSYLLIS CALIFORNIENSIS
331	ANNE	POLY	HARM	IMBR	ANNELIDA POLYNOIDAE HARMOTHOE IMBRICATA
333	ANNE	NERE	NERI	LIMN	ANNELIDA NEREIDAE NEREIS LIMNICOLA
335	ANNE	NERE	NERI	PROC	ANNELIDA NEREIDAE NEREIS PROCERA
337	ANNE	NERE	NERI	SUCC	ANNELIDA NEREIDAE NEREIS SUCCINEA
339	ANNE	GONI	GLYC	ARMI	ANNELIDA GONIADIDAE GLYCIDAE ARMIGERA
341	ANNE	SABE	FABR	BERK	ANNELIDA SABELLIDAE FABRICIA BERKELEYI
343	ANNE	SABE	MANA	SPEC	ANNELIDA SABELLIDAE MANAYUNKIA SPECIOSA
345	ANNE	SABE	POTM	SP A	ANNELIDA SABELLIDAE POTAMILLA SPECIES A
401	ARTH	CYPR	EUCY	SP A	ARTHROPODA CYPRIDIDAE EUCYPRIS SPECIES A
403	ARTH	CYPR	ISOC	SP A	ARTHROPODA CYPRIDIDAE ISOCYPRIS SPECIES A
405	ARTH	CAND	CAND	SP A	ARTHROPODA CANDONIDAE CANDONA SPECIES A
407	ARTH	CYLI	SARS	ZOST	ARTHROPODA CYLINDROLEBERIDIDAE Sarsiella ZOSTERICOLA
409	ARTH	CYTH	CYPR	SP A	ARTHROPODA CYTHERIDAE CYPRIDEIS SPECIES A
411	ARTH	MYSI	NEOM	MERC	ARTHROPODA MYSIDAE NEOMYSIS MERCEDIS
415	ARTH	BALA	BALA	IMPR	ARTHROPODA BALANIDAE BALANUS IMPROVISUS
421	ARTH	NANN	CUME	VULG	ARTHROPODA NANNASTACIDAE CUMELLA VULGARIS
423	ARTH	LEUC	HEMI	HINU	ARTHROPODA LEUCONIDAE HEMILEUCON HINUMENSIS
427	ARTH	TANA	TANA	SP A	ARTHROPODA TANAIDAE TANAIIS SPECIES A
431	ARTH	IDOT	SYNI	LATI	ARTHROPODA IDOTEIDAE SYNIDOTEA LATICAUDA
433	ARTH	ASEL	ASEL	OCCI	ARTHROPODA ASELLIDAE ASELLUS OCCIDENTALIS


ORG NUM	PHYLUM	FAMILY	GENUS	SPECIES	FULL TAXONOMIC NAME
435	ARTH	SPHA	GNOR	INSU	ARTHROPODA SPHAEROMATIDAE GNORIMOSPHAEROMA INSULARE
437	ARTH	SPHA	GNOR	OREG	ARTHROPODA SPHAEROMATIDAE GNORIMOSPHAEROMA OREGONENSIS
439	ARTH	SPHA	SPRO	PENT	ARTHROPODA SPHAEROMATIDAE SPHAEROMA PENTODON
441	ARTH	MUNN	MUNN	SP A	ARTHROPODA MUNNIDAE UID MUNNID SPECIES A
451	ARTH	AMPE	AMPE	ABDI	ARTHROPODA AMPELISCIDAE AMPELISCA ABDITA
453	ARTH	CORO	CORO	ACHE	ARTHROPODA COROPHIIDAE COROPHIUM ACHERUSICUM
455	ARTH	CORO	CORO	ALIE	ARTHROPODA COROPHIIDAE COROPHIUM ALIENENSE
457	ARTH	CORO	CORO	INSI	ARTHROPODA COROPHIIDAE COROPHIUM INSIDIOSUM
459	ARTH	CORO	CORO	OAKL	ARTHROPODA COROPHIIDAE COROPHIUM OAKLANDENSE
461	ARTH	CORO	CORO	SPIN	ARTHROPODA COROPHIIDAE COROPHIUM SPINICORNE
463	ARTH	CORO	CORO	STIM	ARTHROPODA COROPHIIDAE COROPHIUM STIMPSONI
465	ARTH	CORO	CORO	HETE	ARTHROPODA COROPHIIDAE COROPHIUM HETERO CERATUM
467	ARTH	CORO	GRAN	JAPO	ARTHROPODA COROPHIIDAE GRANDIDIERELLA JAPONICA
471	ARTH	PHOX	PARX	MILL	ARTHROPODA PHOXOCEPHALIDAE PARAPHOXUS MILLERI
473	ARTH	PLEU	PARP	PUGE	ARTHROPODA PLEUSTIDAE PARAPLEUSTES PUGETTENSIS
475	ARTH	GAMM	GAMM	DAIB	ARTHROPODA GAMMARIDAE GAMMARUS DAIBERI
477	ARTH	GAMM	MELI	NITI	ARTHROPODA GAMMARIDAE MELITA NITIDA
479	ARTH	GAMM	ELAS	ANTE	ARTHROPODA GAMMARIDAE ELASMOPUS ANTENNATUS
481	ARTH	TALI	HYAL	AZTE	ARTHROPODA TALITRIDAE HYALELLA AZTECA
483	ARTH	CAPR	CAPR	SP A	ARTHROPODA CAPRELLIDAE CAPRELLA SPECIES A
487	ARTH	PALA	PALA	MACR	ARTHROPODA PALAEMONIDAE PALAEMON MACRODACTYLUS
489	ARTH	CRAN	CRAN	FRAN	ARTHROPODA CRANGONIDAE CRANGON FRANCISCORUM
491	ARTH	ASTA	PACI	LENI	ARTHROPODA ASTACIDAE PACIFASTACUS LENIUSCULUS
493	ARTH	CALL	UPOG	PUGE	ARTHROPODA CALLIANASSIDAE UPOGEBIA PUGETTENSIS
495	ARTH	XANT	RHIT	HARR	ARTHROPODA XANTHIDAE RHITHROPANOPEUS HARRISII
497	ARTH	GRAP	HEMG	NUDU	ARTHROPODA GRAPSIDAE HEMIGRAPUS NUDUS
499	ARTH	MAJI	PYRO	TUBE	ARTHROPODA MAJIDAE PYROMAIA TUBERCULATA
501	ARTH	BAET	BAET	BICA	ARTHROPODA BAETIDAE BAETIS BICAUDATUS
503	ARTH	HEPT	HEPT	ROSE	ARTHROPODA HEPTAGENIIDAE HEPTAGENIA ROSEA
505	ARTH	LEPT	PARE	SP A	ARTHROPODA LEPTOPHLEBIIDAE PARALEPTOPHLEBIA SPECIES A
507	ARTH	EPHE	HEXA	LIMB	ARTHROPODA EPHEMERIDAE HEXAGENIA LIMBATA CALIFORNICA
509	ARTH	TRIC	TRYC	SP A	ARTHROPODA TRICORYTHIDAE TRICORYTHODES SPECIES A
511	ARTH	CAEN	CAEN	SIMU	ARTHROPODA CAENIDAE CAENIS SIMULANS
513	ARTH	GOMP	GOMP	OLIV	ARTHROPODA GOMPHIDAE GOMPHUS OLIVACEUS
515	ARTH	COEN	ZONI	EXCL	ARTHROPODA COENAGRIONIDAE ZONIAGRION EXCLAMATIONIS
517	ARTH	NAUC	AMBR	SP A	ARTHROPODA NAUCORIDAE AMBRYBUS SPECIES A
519	ARTH	CORI	CORI	INSC	ARTHROPODA CORIXIDAE CORISELLA INSCRIPTA
521	ARTH	CORI	TRIK	VERT	ARTHROPODA CORIXIDAE TRICHOCORIXA VERTICALIS
523	ARTH	HYDO	HYDO	SP A	ARTHROPODA HYDROPSYCHIDAE HYDROPSYCHE SPECIES A
525	ARTH	HYOP	HYOP	SP A	ARTHROPODA HYDROPTILIDAE HYDROPTILA SPECIES A
527	ARTH	HYOP	OXYE	SP A	ARTHROPODA HYDROPTILIDAE OXYETHIRA SPECIES A
529	ARTH	LEPC	NECT	GRAC	ARTHROPODA LEPTOCERIDAE NECTOPSYCHE GRACILIS
531	ARTH	LEPC	OECE	SP A	ARTHROPODA LEPTOCERIDAE OECETIS SPECIES A
533	ARTH	CHAO	CHAO	ALBA	ARTHROPODA CHAOBORIDAE CHAOBORUS ALBATUS
541	ARTH	CHIR	PROC	SP A	ARTHROPODA CHIRONOMIDAE PROCLADIUS SPECIES A
543	ARTH	CHIR	TANY	STEL	ARTHROPODA CHIRONOMIDAE TANYPUS STELLATUS
545	ARTH	CHIR	ABLA	SP A	ARTHROPODA CHIRONOMIDAE ABLABESMYIA SPECIES A
547	ARTH	CHIR	CLAD	SP A	ARTHROPODA CHIRONOMIDAE CLADOTANYTARSUS SPECIES A
549	ARTH	CHIR	MICR	SP A	ARTHROPODA CHIRONOMIDAE MICROPSECTRA SPECIES A
551	ARTH	CHIR	PARY	SP A	ARTHROPODA CHIRONOMIDAE PARATANYTARSUS SPECIES A
553	ARTH	CHIR	CHIR	ATTE	ARTHROPODA CHIRONOMIDAE CHIRONOMUS ATTENUATUS

ORG NUM.	PHYLUM	FAMILY	GENUS	SPECIES	FULL TAXONOMIC NAME
555	ARTH	CHIR	CRYP	SP A	ARTHROPODA CHIRONOMIDAE CRYPTOCHIRONOMUS SPECIES A
557	ARTH	CHIR	CRYP	SP B	ARTHROPODA CHIRONOMIDAE CRYPTOCHIRONOMUS SPECIES B
559	ARTH	CHIR	DEMI	SP A	ARTHROPODA CHIRONOMIDAE DEMICRYPTOCHIRONOMUS SPECIES A
561	ARTH	CHIR	EINF	SP A	ARTHROPODA CHIRONOMIDAE EINFELDIA SPECIES A
563	ARTH	CHIR	ENDO	SP A	ARTHROPODA CHIRONOMIDAE ENDOCHIRONOMUS SPECIES A
565	ARTH	CHIR	ENDO	SP B	ARTHROPODA CHIRONOMIDAE ENDOCHIRONOMUS SPECIES B
567	ARTH	CHIR	HARN	CURT	ARTHROPODA CHIRONOMIDAE HARNISCHIA CURTILAMELLATA
569	ARTH	CHIR	PARA	SP A	ARTHROPODA CHIRONOMIDAE PARACHIRONOMUS SPECIES A
571	ARTH	CHIR	PARC	SP A	ARTHROPODA CHIRONOMIDAE PARACLADOPELMA SPECIES A
573	ARTH	CHIR	PARL	SP A	ARTHROPODA CHIRONOMIDAE PARALAUTERBORNIELLA SPECIES A
575	ARTH	CHIR	PART	SP A	ARTHROPODA CHIRONOMIDAE PARATENDIPES SPECIES A
577	ARTH	CHIR	PHAE	SP A	ARTHROPODA CHIRONOMIDAE PHAENOPSECTRA SPECIES A
579	ARTH	CHIR	POLY	SP A	ARTHROPODA CHIRONOMIDAE POLYPEDILUM SPECIES A
581	ARTH	CHIR	ROBA	CLAV	ARTHROPODA CHIRONOMIDAE ROBACKIA CLAVIGER
583	ARTH	CHIR	STEC	SP A	ARTHROPODA CHIRONOMIDAE STENOCHIRONOMUS SPECIES A
585	ARTH	CHIR	STIC	SP A	ARTHROPODA CHIRONOMIDAE STICTOCHIRONOMUS SPECIES A
587	ARTH	CHIR	MONO	SP A	ARTHROPODA CHIRONOMIDAE MONODIAMESA SPECIES A
589	ARTH	CHIR	CRIC	BICI	ARTHROPODA CHIRONOMIDAE CRICOTOPUS BICINCTUS
591	ARTH	CHIR	CRIC	SP A	ARTHROPODA CHIRONOMIDAE CRICOTOPUS SPECIES A
593	ARTH	CHIR	EPOI	SP A	ARTHROPODA CHIRONOMIDAE EPOICOCADIUS SPECIES A
595	ARTH	CHIR	NANO	DIST	ARTHROPODA CHIRONOMIDAE NANOCLADIUS DISTINCTUS
597	ARTH	CHIR	NANO	SP A	ARTHROPODA CHIRONOMIDAE NANOCLADIUS SPECIES A
599	ARTH	CHIR	PSEC	SP A	ARTHROPODA CHIRONOMIDAE PSECTROCLADIUS SPECIES A
601	ARTH	CERA	PALP	SP A	ARTHROPODA CERATOPOGONIDAE PALPOMYIA SPECIES A
603	ARTH	LIME	LIME	SP A	ARTHROPODA LIMNESIIDAE LIMNESIA SPECIES A
605	ARTH	PION	FORE	SP A	ARTHROPODA PIONIDAE FORELIA SPECIES A
607	ARTH	UNIO	UNIO	SP A	ARTHROPODA UNIONICOLIDAE UNIONICOLA SPECIES A
609	ARTH	UNIO	UNIO	SP B	ARTHROPODA UNIONICOLIDAE UNIONICOLA SPECIES B
651	MOLL	UNKN	NUDI	SP A	MOLLUSCA UNKNOWN NUDIBRANCH SPECIES A
653	MOLL	PLAO	GYRA	SP A	MOLLUSCA PLANORBIDAE GYRAULUS SPECIES A
655	MOLL	PLAO	GYRA	SP B	MOLLUSCA PLANORBIDAE GYRAULUS SPECIES B
657	MOLL	PHYS	PHYS	GRYI	MOLLUSCA PHYSIDAE PHYSIA GYRINA
659	MOLL	ANCY	FERR	RIVU	MOLLUSCA ANCYLIDAE FERRISSIA RIVULARIS
661	MOLL	ASSI	ASSI	CALI	MOLLUSCA ASSIMINEIDAE ASSIMINEA CALIFORNICA
663	MOLL	THIA	MELA	TUBE	MOLLUSCA THIARIDAE MELANOIDES TUBERCULATA
665	MOLL	PYRA	ODOS	FETE	MOLLUSCA PYRAMIDELLIDAE ODOSTOMIA FETELLA
667	MOLL	SEME	THEO	LUBR	MOLLUSCA SEMELIDAE THEORA LUBRICA
669	MOLL	MYTI	MUSC	SENH	MOLLUSCA MYTILIDAE MUSCULISTA SENHOUSIA
671	MOLL	MYTI	MYTI	EDUL	MOLLUSCA MYTILIDAE MYTILUS EDULIS
673	MOLL	CORB	CORB	FLUM	MOLLUSCA CORBICULIDAE CORBICULA FLUMINEA
675	MOLL	SPHE	PISI	CASE	MOLLUSCA SPHAERIIDAE PISIDIUM CASERTANUM
677	MOLL	SPHE	PISI	COMP	MOLLUSCA SPHAERIIDAE PISIDIUM COMPRESSUM
679	MOLL	SPHE	SPHE	SP A	MOLLUSCA SPHAERIIDAE SPHAERIUM SPECIES A
681	MOLL	TELL	MACO	BALT	MOLLUSCA TELLINIDAE MACOMA BALTHICA
683	MOLL	VENE	PROT	STAM	MOLLUSCA VENERIDAE PROTOTHACA STAMINEA
685	MOLL	VENE	GEMM	GEMM	MOLLUSCA VENERIDAE GEMMA GEMMA
687	MOLL	MYID	MYA	AREN	MOLLUSCA MYIDAE MYA ARENARIA
689	MOLL	CORL	POTA	AMUR	MOLLUSCA CORBULIDAE POTAMOCORBULA AMURENSIS
691	MOLL	UNIN	ANOD	NUTT	MOLLUSCA UNIONIDAE ANODONTA NUTTALLIANA
700	CHOR	MOLG	MOLG	MANH	CHORDATA MOLGULIDAE MOLGULA MANHATTENSIS


Appendix B

**Graphs of Mean Monthly Abundance of
All Species (Total)
and the
Four Numerically Dominant Species at Each Benthic Site**


MEAN TOTAL ABUNDANCE AND MEAN MONTHLY ABUNDANCE OF THE
FOUR NUMERICALLY DOMINANT ORGANISMS AT SITE D7-C




MEAN TOTAL ABUNDANCE AND MEAN MONTHLY ABUNDANCE OF THE FOUR NUMERICALLY DOMINANT ORGANISMS AT SITE D11-C




MEAN TOTAL ABUNDANCE AND MEAN MONTHLY ABUNDANCE OF THE
FOUR NUMERICALLY DOMINANT ORGANISMS AT SITE D4-R




MEAN TOTAL ABUNDANCE AND MEAN MONTHLY ABUNDANCE OF THE
FOUR NUMERICALLY DOMINANT ORGANISMS AT SITE D4-L




MEAN TOTAL ABUNDANCE AND MEAN MONTHLY ABUNDANCE OF THE
FOUR NUMERICALLY DOMINANT ORGANISMS AT SITE D4-C




MEAN TOTAL ABUNDANCE AND MEAN MONTHLY ABUNDANCE OF THE FOUR NUMERICALLY DOMINANT ORGANISMS AT SITE D19-C



MEAN TOTAL ABUNDANCE AND MEAN MONTHLY ABUNDANCE OF THE
FOUR NUMERICALLY DOMINANT ORGANISMS AT SITE D28A-L



MEAN TOTAL ABUNDANCE AND MEAN MONTHLY ABUNDANCE OF THE
FOUR NUMERICALLY DOMINANT ORGANISMS AT SITE D28A-R



COMMON ABBREVIATIONS AND METRIC CONVERSIONS

Area

km ²	square kilometers; to convert to square miles, multiply by 0.3861
m ²	square meters; to convert to square feet, multiply by 10.764

Length

cm	centimeters; to convert to inches, multiply by 0.3937
FL	fork length; length from the most anterior part of a fish to the median caudal fin rays (fork in the tail)
km	kilometers; to convert to miles, multiply by 0.62139
m	meters; to convert to feet, multiply by 3.2808
mm	millimeters; to convert to inches, multiply by 0.03937
SL	standard length; tip of upper jaw of a fish to crease formed when tail is bent sharply upward
TL	total length; length from the most anterior part of a fish to the end of the tail

Volume

AF	acre-foot; equal to 43,560 cubic feet
L	liters; to convert to quarts, multiply by 1.05668; to convert to gallons, multiply by 0.26417
mL	milliliters

Flow

cfs	cubic feet per second; to convert to acre-feet per day, multiply by 1.98
gpm	gallons per minute
mgd	million gallons per day

Velocity

fps	feet per second
m/s	meters per second; to convert to feet per second, multiply by 3.2808

Mass

kg	kilograms; to convert to pounds, multiply by 2.2046
----	---

Concentration

mg/L	milligrams per liter; equals parts per million (ppm)
µg/L	micrograms per liter; equals parts per billion (ppb)

Specific Conductance

µS	microsiemens; equivalent to micromhos
µS/cm	microsiemens per centimeter

Temperature

°C	degrees Celsius; to convert to °F, multiply by 1.8 then add 32 degrees
°F	degrees Fahrenheit; to convert to °C, subtract 32 degrees then divide by 1.8

Mathematics and Statistics

df	degrees of freedom
e	base of natural logarithm
E	expected value
log	logarithm
N	sample size
NS	not significant
%	percent
‰	per thousand
P	probability
r	correlation or regression coefficient (simple)
R	correlation or regression coefficient (multiple)
SD	standard deviation
SE	standard error
V	variance

Interagency Program Members

COE	U.S. Army Corps of Engineers
DFG	California Department of Fish and Game
DWR	California Department of Water Resources
EPA	U.S. Environmental Protection Agency
FWS	U.S. Fish and Wildlife Service
SWRCB	California State Water Resources Control Board
USBR	U.S. Bureau of Reclamation
USGS	U.S. Geological Survey

General

CPUE	catch per unit effort
YOY	young of the year

SCIENTIFIC NAMES OF FISH

American eel	<i>Anguilla rostrata</i>	pumpkin seed	<i>Lepomis gibbosus</i>
American shad	<i>Alosa sapidissima</i>	rainwater killifish	<i>Lucania parva</i>
bay goby	<i>Lepidogobius lepidus</i>	redeer sunfish	<i>Lepomis microlophus</i>
bigscale logperch	<i>Percina macrolepida</i>	red shiner	<i>Cyprinella lutrensis</i>
black bullhead	<i>Ameiurus melas</i>	rifle sculpin	<i>Cottus gulosus</i>
black crappie	<i>Pomoxis nigromaculatus</i>	river lamprey	<i>Lampetra ayresii</i>
blue catfish	<i>Ictalurus furcatus</i>	Sacramento blackfish	<i>Orthodon microlepidotus</i>
bluegill	<i>Lepomis macrochirus</i>	Sacramento perch	<i>Archoplites interruptus</i>
brown bullhead	<i>Ameiurus nebulosus</i>	Sacramento splittail	<i>Pogonichthys macrolepidotus</i>
brown trout	<i>Salmo trutta</i>	Sacramento squawfish	<i>Ptychocheilus grandis</i>
California halibut	<i>Paralichthys californicus</i>	Sacramento sucker	<i>Catostomus occidentalis</i>
California roach	<i>Hesperoleucus symmertricus</i>	shiner surfperch	<i>Cymatogaster aggregata</i>
chameleon goby	<i>Tridentiger trigonocephalus</i>	silver salmon	<i>Oncorhynchus kisutch</i>
channel catfish	<i>Ictalurus punctatus</i>	smallmouth bass	<i>Micropterus dolomieu</i>
Chinook salmon	<i>Oncorhynchus tshawytscha</i>	speckled dace	<i>Rhinichthys osculus</i>
common carp	<i>Cyprinus carpio</i>	speckled sanddab	<i>Citharichthys stigmaeus</i>
delta smelt	<i>Hypomesus transpacificus</i>	splittail	<i>Pogonichthys macrolepidotus</i>
English sole	<i>Pleuronectes vetulus</i>	staghorn sculpin	<i>Leptocottus armatus</i>
fathead minnow	<i>Pimephales promelas</i>	starry flounder	<i>Platichthys stellatus</i>
golden shiner	<i>Notemigonus crysoleucas</i>	steelhead trout	<i>Oncorhynchus mykiss</i>
goldfish	<i>Carassius auratus</i>	striped bass	<i>Morone saxatilis</i>
green sturgeon	<i>Acipenser medirostris</i>	striped mullet	<i>Mugil cephalus</i>
green sunfish	<i>Lepomis cyanellus</i>	surf smelt	<i>Hypomesus pretiosus</i>
hardhead	<i>Mylopharodon conocephalus</i>	threadfin shad	<i>Dorosoma petenense</i>
hitch	<i>Lavinia exilicauda</i>	threespine stickleback	<i>Gasterosteus aculeatus</i>
inland silverside	<i>Menidia beryllina</i>	tui chub	<i>Gila bicolor</i>
jacksmelt	<i>Atherinopsis californiensis</i>	tule perch	<i>Hysteroecarpus traski</i>
largemouth bass	<i>Micropterus salmoides</i>	wakasagi	<i>Hypomesus nipponensis</i>
longfin smelt	<i>Spirinchus thaleichthys</i>	warmouth	<i>Lepomis gulosus</i>
mosquitofish	<i>Gambusia affinis</i>	white catfish	<i>Ameiurus catus</i>
northern anchovy	<i>Engraulis mordax</i>	white crappie	<i>Pomoxis annularis</i>
Pacific herring	<i>Clupea pallasii</i>	white croaker	<i>Genyonemus lineatus</i>
Pacific lamprey	<i>Lampetra tridentata</i>	white sturgeon	<i>Acipenser transmontanus</i>
pink salmon	<i>Oncorhynchus gorbuscha</i>	yellow bullhead	<i>Ameiurus natalis</i>
plainfin midshipman	<i>Porichthys notatus</i>	yellow perch	<i>Perca flavescens</i>
prickly sculpin	<i>Cottus asper</i>	yellowfin goby	<i>Acanthogobius flavimanus</i>