

Fact Sheet **SPoT 2012**

LAND USE, POLLUTANTS, AND WATER QUALITY IN CALIFORNIA STREAMS

Stream Pollution Trends (SPoT) program begins with survey of watersheds statewide

Overview

The State Water Resources Control Board's Surface Water Ambient Monitoring Program (SWAMP) has released a report on results from the first year of a continuing statewide program that measures trends in pollution levels and toxicity in California streams. The report, *Statewide Perspective on Chemicals of Concern and Connections between Stream Water Quality and Land Use*, represents the first consistent assessment of large watersheds across California to determine how stream pollutant concentrations are affected by urban and agricultural development. The report describes surveys conducted in 2008, which established the baseline for determining long-term trends in the types and amounts of pollutants in stream sediments and their potential for impacting aquatic life. The study is valuable for improving our understanding of the effects of land development on stream water quality, for prioritizing areas in need of water quality management, and for evaluating the effectiveness of management programs designed to improve stream health. The study also provides a statewide perspective so that local and regional water quality monitoring efforts can evaluate how conditions in their streams compare to those in other California watersheds.

The State Water Resources Control Board's Surface Water Ambient Monitoring Program (SWAMP) has released a report on results from the first year of a continuing statewide program that measures trends in pollution levels and toxicity in California streams.



April 2012



www.waterboards.ca.gov/swamp

“The results indicate that, on a statewide basis, toxicity and concentrations of most measured pollutants were higher in streams that drain watersheds with higher proportions of urban land cover.”

“Stream sediment concentrations of heavy metals, such as cadmium, copper, lead and zinc, and industrial organic compounds such as PCBs, flame retardants, and hydrocarbons, were generally highest in stream sediments from Los Angeles and San Francisco Bay area watersheds.”

About the Survey

To most efficiently detect pollutant trends in California streams, the SPoT program measures contaminant concentrations and toxicity in stream sediments that accumulate in downstream reaches of large watersheds. In 2008, samples were collected from 92 of the nearly 200 major hydrologic units in California. In the SPoT program, sediment samples are collected once each year when streams return to base flow conditions after the high flows that carry pollutants washed from watershed surfaces during storms. Sediments are monitored because the majority of contaminants entering streams accumulate in sediments, and this environmental compartment integrates pollution signals over time. Each sample is analyzed for industrial compounds (PAHs, PBDEs, and PCBs), pesticides, and metals, and is tested for toxicity to aquatic organisms. Results are compared across watersheds throughout the state, and pollutant concentrations are compared to land use and other human activities.

Findings

A number of spatial patterns were evident in this survey of watersheds from throughout California. Stream sediment concentrations of heavy metals, such as cadmium, copper, lead and zinc, tended to be highest in urbanized Los Angeles and San Francisco Bay area watersheds. Metal concentrations in sediments from San Leandro Creek (SF Bay area), Ballona Creek and San Gabriel River (Los Angeles area) were among the highest in the state. These metals are released to the environment from brake pads, plumbing, industrial and commercial activities. Industrial organic compounds such as PCBs (banned since the 1970s), PBDEs (flame retardants) and PAHs (hydrocarbons) were also generally highest in stream sediments from the Los Angeles and San Francisco Bay areas, though high concentrations were also measured in some more remote locations. In contrast to this urban pattern, mercury concentrations were highest in sediments from watersheds where it is geologically abundant and historically mined, though

some urban streams also had relatively high mercury levels. The persistent legacy pesticide DDT was found in stream sediments from most urban and agricultural watersheds, where recent soil disturbance from development and tillage likely mobilizes DDT residues from applications more than 40 years ago. Pyrethroid pesticides are increasing in use in California and were detected in 55% of the samples statewide (51 of 92). The highest pyrethroid concentrations were measured in sediments collected from urban watersheds, plus two agricultural watersheds along the central coast. Approximately one quarter of the sediment samples collected were significantly toxic to amphipods, which are resident crustaceans representative of important aquatic food web links. Of these samples, 6.5% were identified as highly toxic. The highly toxic samples were collected from agricultural watersheds in the Tulare basin and central coast, in urban areas of southern California, and in the Tijuana River. Other toxic samples were collected from a wide range

of watershed types, including those along the north coast, the Tahoe area, and urban and agricultural areas across the state.

On a statewide basis, watersheds with greater than 10% urban land cover had consistently higher sediment pollutant concentrations and toxicity than did watersheds characterized by agricultural or other land cover types. Metals, industrial compounds, DDTs, and pyrethroid pesticides were all found at significantly higher concentrations in urban streams. This pattern is evident at the statewide level, but other local and regional studies have previously shown that pollution levels and toxicity can be strongly related to agricultural and other land uses in specific areas. This is because the “agricultural” land use category used in the SPoT analysis is comprised of many different types of crops and farming practices, and the amount of polluted runoff from these lands varies depending on the types of practices employed.



Pajaro River in central California. Photo courtesy of California Coastal Records Project. Used by permission. Copyright ©2002-2010 Kenneth & Gabrielle Adelman, California Coastal Records Project, www.californiacoastline.org

What's Next?

The SWAMP Team has continued to survey the SPoT program sites once each year to determine how stream pollution is changing over time. As land use changes and management programs work to improve water quality, the SPoT program will provide a valuable indication of the effectiveness of our efforts to protect California streams. The second SPoT report, covering data collected from 2008 to 2010, will be submitted for review later this year.




Urban stream near the southern California coast.

The SPoT program was developed to inform a number of management initiatives and provide a network linking state, regional, and local monitoring programs. These SPoT surveys provide statewide context and specific data for developing the California Integrated Report. SPoT data on stream pollution are useful for evaluating watersheds in the EPA Measure W program, as well as determining chemicals of concern for total maximum daily load allocations in impaired streams. Stormwater agencies and Regional Water Boards will use SPoT data to determine compliance with water quality regulations, and to evaluate the effectiveness of management practices to improve water quality. SWAMP will continue to integrate its statewide and regional programs so that SPoT chemistry and toxicity data will be assessed in conjunction with ecological data collected at the same sites by the SWAMP Perennial Streams Assessment, the Southern California Stormwater Monitoring Coalition, the Bay Area Stormwater Management Agency Association's Regional Monitoring Coalition and other regional programs. SPoT data on mercury and other bioaccumulating chemicals in streams will help identify sources and transport pathways for chemicals measured in edible fish tissues by SWAMP Bioaccumulation Program surveys of downstream lakes, rivers, and coastal waters.


Acknowledgments

"Pyrethroid pesticides were detected in stream sediments from more than half the watersheds surveyed, and were measured at concentrations high enough to cause toxicity in more than a quarter of the total samples."

The Stream Pollution Trend (SPoT) program surveys are funded by the State Water Board's Surface Water Ambient Monitoring Program (SWAMP) and the United States Environmental Protection Agency (USEPA). This survey was designed and is implemented in collaboration with the California Regional Water Quality Control Boards. The program is conducted by scientists from the University of California Davis' Marine Pollution Studies Laboratory at Granite Canyon, in cooperation with scientists from California State University's Moss Landing Marine Laboratories (MLML), California Department of Fish and Game's Water Pollution Control Laboratory, Rancho Cordova, and Trace Metal Laboratory at MLML, CSU Chico's Geographic Information Center, and the SWAMP data management and quality assurance teams.



Concentrations of pyrethroid pesticides measured in sediment from streams draining large watersheds throughout California. Each symbol represents concentrations of the 8 different pyrethroid pesticides measured at each site, with the concentrations converted to toxic units and then added together to give an estimate of their cumulative potential impact on aquatic organisms. One toxic unit is equivalent to a concentration lethal to half of the toxicity test organisms, which in this case are amphipods of a genus resident to California waters. Previous studies indicate that pyrethroid concentrations above 0.4 toxic units are associated with adverse impacts to similar organisms in the field.



For more information, please contact:

Brian Anderson

Department of Environmental Toxicology

University of California, Davis

c/o Marine Pollution Studies Laboratory

34500 Highway 1 Monterey, CA 93940

(ph) 831-624-0947 (fx) 831-626-1518

anderson@ucdavis.edu

<http://www.envtox.ucdavis.edu/GraniteCanyon/GraniteCanyon.htm>