## **Appendix B:**

Sediment Source Analysis for the Upper Eel River Sediment TMDL

Table B-1 identifies the terrain/geology types for each subwatershed and the entire basin. To determine the terrain types, the USFS Wilderness Area was removed and the USFS and State Geology maps were reviewed for the remaining basin. Similar rock types were then lumped into five basic categories: Schist, Melange, Alluvium, Coastal Belt, and Franciscan. A grid of 41.8 acre cells was then applied over the entire basin and the dominant terrain type for each cell was determined. Cells were randomly selected for inclusion in the stratified random sampling study of small sediment sources (plot features <3,000 yds³). Table B-1 also identifies the number of grid cells, their total area, and the number of plots sampled for each watershed, geology type, and ownership.

Table B-1. Number of grid cells, basin area, and sample plots by ownership, watershed, and terrain type for the Upper Eel River study area (USFS Wilderness Areas excluded)

Watershed	Stratum	Stratum Terrain/ Geology		Publ	ic domain			Priva	te domain		Total study area (public + private)			
		Geology Type	# grid cells	Area of grid cells (mi²)	Proposed grid cell sample	PWA # of grid cells sampled		Area of grid cells (mi <sup>2</sup> )	Proposed grid cell sample	PWA # of grid cells sampled	# grid cells	Area of grid cells (mi <sup>2</sup> )	Proposed grid cell sample	PWA # of grid cells sampled
Entire	1	Schist	1,955	127.6	11	11	268	17.5	4	4	2,223	145.1	15	15
Upper Eel watershed	2	Melange	609	39.8	7	7	376	24.6	7	7	985	64.4	14	14
study area	3	Alluvium	55	3.6	3	3	204	13.3	3	2	259	16.9	6	5
	4	Coastal Belt	164	10.7	5	5	830	54.2	10	8	994	64.9	15	13
	5	Franciscan	2,250	147.0	14	14	3,826	249.8	16	12	6,076	396.8	30	26
	Total		5,033	328.7	40	40	5,504	359.4	40	33	10,537	688.1	80	73
	Lake Pillst removed	oury area	46	3.0	Ī	-	ı		-	-	46	3.0	-	_
	Wilderness removed	s area	270	17.6	_	_	-		_	_	270	17.6	_	

Table B-1 (continued)

Watershed	Stratum	Terrain/		Publ	ic domain			Priva	ate domain		Total study area (public + private)			
		Geology Type	# grid cells	Area of grid cells (mi <sup>2</sup> )	Proposed grid cell sample	PWA # of grid cells sampled	# grid cells	Area of grid cells (mi²)	Proposed grid cell sample	PWA # of grid cells sampled	# grid cells	Area of grid cells (mi²)	Proposed grid cell sample	PWA # of grid cells sampled
Outlet	1	Schist	0	0.0		0	0	0.0		0	0	0.0		0
Creek	2	Melange	0	0.0		0	0	0.0		0	0	0.0		0
	3	Alluvium	0	0.0		0	189	12.3		2	189	12.3		2
	4	Coastal Belt	3	0.2		0	468	30.6		0	471	30.8		0
	5	Franciscan	62	4.0		0	1,725	112.7		4	1,787	116.7		4
	Subtotal		65	4.2		0	2,382	155.6		6	2,447	159.8		6
Rice Fork	1	Schist	95	6.2		2	28	1.8		1	123	8.0	-	3
	2	Melange	34	2.2		1	18	1.2		0	52	3.4		1
	3	Alluvium	0	0.0		0	0	0.0		0	0	0.0		0
	4	Coastal Belt	0	0.0		0	0	0.0		0	0	0.0	-	0
	5	Franciscan	1,059	69.2		6	131	8.6		1	1,190	77.8	-	6
	Subtotal		1,188	77.6		9	177	11.6		1	1,365	89.2	-	10
Soda Creek	1	Schist	482	31.5		1	116	7.6		1	598	39.1		2
	2	Melange	11	0.7		0	36	2.4		0	47	3.1		0
	3	Alluvium	0	0.0		0	0	0.0		0	0	0.0	-	0
	4	Coastal Belt	78	5.1		0	187	12.2		3	265	17.3		3
	5	Franciscan	0	0.0		0	10	0.6		0	10	0.6		0
	Subtotal		571	37.3		1	349	22.8		4	920	60.1		5

Table B-1 (continued)

Watershed	ĺ	Terrain/		Publ	ic domain			Priva	nte domain		Total study area (public + private)			
		Geology Type	# grid cells	Area of grid cells (mi <sup>2</sup> )	Proposed grid cell sample	PWA # of grid cells sampled	# grid cells	Area of grid cells (mi <sup>2</sup> )	Proposed grid cell sample	PWA # of grid cells sampled	# grid cells	Area of grid cells (mi²)	Proposed grid cell sample	PWA # of grid cells sampled
Tomki	1	Schist	229	15.0		1	16	1.0		0	245	16.0		1
Creek	2	Melange	30	2.0		0	321	21.0		7	351	23.0		7
	3	Alluvium	0	0.0		0	1	0.1		0	1	0.1		0
	4	Coastal Belt	83	5.4		5	175	11.3		5	258	16.7		10
	5	Franciscan	327	21.3		2	1,882	122.9		8	2,209	144.2		10
	Subtotal		669	43.7		8	2,395	156.3		20	3,064	200.0		28
Upper	1	Schist	1,149	75.0		7	108	7.0		2	1,257	82.0		9
Main Eel	2	Melange	534	34.9		6	1	0.1		0	535	35.0		6
	3	Alluvium	55	3.6		3	14	0.9		0	69	4.5		3
	4	Coastal Belt	0	0.0		0	0	0.0		0	0	0.0		0
	5	Franciscan	802	52.4		6	78	5.1		0	880	57.5		6
	Subtotal		2,540	165.9		22	201	13.1		2	2,741	179.0		24

Table B-2 illustrates the distribution of erosion features by ownership and terrain type for small sediment sources (<3,000 yds<sup>3</sup>) in addition to the total measured erosion and estimated sediment delivery. These numbers reflect actual plot data (before erosion and sediment delivery volumes were extrapolated to the entire basin).

Table B-2. Total measured erosion and sediment delivery measured in the 73 sample plots by ownership, terrain type and erosional feature type for the Upper Eel River watershed study area

	N	umber of Field I	Measured Featur	es by Terrain Type	(#)	Total measured	Total estimated
Erosional Feature Type	1. Schist	2. Melange	3. Alluvium	4. Coastal Belt	5. Franciscan	erosion (yds³)	sediment delivery (yds <sup>3</sup> )
Public Domain (plots/mi²)	11/127.6	7/39.8	3/3.6	5/10.7	14/147	40/328.7	40/328.7
Debris Slide (DL)	19	10	2	17	49	15,856	10,109
Debris Torrent Track (DT)	0	0	0	0	0	0	0
Bank Erosion (BE)	10	4	14	13	23	6,254	6,061
Road related gully (GU)	3	1	2	0	6	156	133
Non road related gully (GU)	11	0	0	1	2	414	238
Stream Crossing (XI)	5	0	0	0	2	183	132
Channel Incision (CI)	11	1	0	1	4	577	576
Surface Erosion (SE)	7	2	1	0	6	635	337
Subtotals	66	18	19	32	92	24,075	17,586

Table B- 2 (continued)

	N	umber of Field I	Measured Featur	es by Terrain Type (	(#)	Total measured	Total estimated
Erosional Feature Type	1. Schist	2. Melange	3. Alluvium	4. Coastal Belt	5. Franciscan	erosion (yds³)	sediment delivery (yds³)
Private Domain (plots/mi²)	4/17.5	7/24.6	2/13.3	8/54.2	12/249.8	33/359.4	33/359.4
Debris Slide (DL)	14	31	0	19	47	45,010	15,038
Debris Flow Source (DF)	1	0	0	0	0	444	400
Bank Erosion (BE)	5	16	21	16	60	5,212	4,934
Road related gully (GU)	3	0	1	1	12	658	578
Non road related gully (GU)	5	7	0	8	3	1,065	1,001
Stream Crossing (XI)	2	0	0	6	10	410	401
Surface Erosion (SE)	1	0	0	0	0	60	54
Active Earthflow (SDS)	0	3	0	0	4	1,461	175
Channel Incision (CI)	1	9	0	6	19	922	914
Subtotals	32	50	22	56	155	55,242	23,495
Total # of features, erosion and delivery volumes for all domains	98	68	41	88	247	79,317	41,081
Total # of plots/terrain type area in mi <sup>2</sup>	15/145.1	14/64.4	5/16.9	13/62.2	26/396.8	73/688.1	73/688.1

Table B-2a. Total measured erosion and sediment delivery measured in the 73 sample plots by ownership, terrain type and erosional feature type for the Outlet Creek subwatershed, Upper Eel River watershed study area.

	Nı	ımber of Field N	Aeasured Feature	es by Terrain Type (	#)	Total measured	Total estimated
Erosional Feature Type	1. Schist	2. Melange	3. Alluvium	4. Coastal Belt	5. Franciscan	erosion (yds³)	sediment delivery (yds³)
Public Domain (plots/mi²)	0/0	0/0	0/0	0/0.2	0/4	0/4.2	0/4.2
Debris Slide (DL)	0	0	0	0	0	0	0
Debris Flow Source (DF)	0	0	0	0	0	0	0
Debris Torrent Track (TT)	0	0	0	0	0	0	0
Bank Erosion (BE)	0	0	0	0	0	0	0
Road related gully (GU)	0	0	0	0	0	0	0
Non road related gully (GU)	0	0	0	0	0	0	0
Stream Crossing (XI)	0	0	0	0	0	0	0
Channel Incision (CI)	0	0	0	0	0	0	0
Surface Erosion (SE)	0	0	0	0	0	0	0
Subtotals	0	0	0	0	0	0	0

Table B- 2a (continued)

Tuote B 2a (commaca)	N	Number of Field I	Measured Featur	es by Terrain Type	(#)	Total measured	Total estimated
Erosional Feature Type	1. Schist	2. Melange	3. Alluvium	4. Coastal Belt	5. Franciscan	erosion (yds³)	sediment delivery (yds <sup>3</sup> )
Private Domain (plots/mi²)	0/0	0/0	2/12.3	0/30.6	4/112.7	6/155.6	6/155.6
Debris Slide (DL)	0	0	0	0	20	2,038	1,204
Debris Flow Source (DF)	0	0	0	0	0	0	0
Debris Torrent Track (TT)	0	0	0	0	0	0	0
Bank Erosion (BE)	0	0	21	0	12	1,899	1,748
Road related gully (GU)	0	0	1	0	7	354	291
Non road related gully (GU)	0	0	0	0	2	35	35
Stream Crossing (XI)	0	0	0	0	3	133	129
Surface Erosion (SE)	0	0	0	0	0	0	0
Active Earthflow (SDS)	0	0	0	0	0	0	0
Channel Incision (CI)	0	0	0	0	4	339	339
Subtotals	0	0	22	0	48	4,798	3,746
Total # of features, erosion and delivery volumes for all domains	0	0	22	0	48	4,798	3,746
Total # of plots/terrain type area in mi <sup>2</sup>	0/0	0/0	2/12.3	0/30.8	4/116.7	6/159.8	6/159.8

Table B-2b. Total measured erosion and sediment delivery measured in the 73 sample plots by ownership, terrain type and erosional feature type for the Rice Fork subwatershed, Upper Eel River watershed study area.

		Number of Field	Measured Featu (#)	res by Terrain Type		Total measured erosion	Total estimated sediment delivery	
Erosional Feature Type	1. Schist	2. Melange	3. Alluvium	4. Coastal Belt	5. Franciscan	(yds <sup>3</sup> )	(yds <sup>3</sup> )	
Public Domain (plots/mi²)	2/6.2	1/2.2	0/0	0/0	6/69.2	9/77.6	9/77.6	
Debris Slide (DL)	3	0	0	0	15	1,484	856	
Debris Flow Source (DF)	0	0	0	0	0	0	0	
Debris Torrent Track (TT)	0	0	0	0	0	0	0	
Bank Erosion (BE)	2	0	0	0	14	875	702	
Road related gully (GU)	1	0	0	0	4	46	39	
Non road related gully (GU)	2	0	0	0	0	22	22	
Stream Crossing (XI)	0	0	0	0	0	0	0	
Channel Incision (CI)	5	0	0	0	1	134	134	
Surface Erosion (SE)	0	0	0	0	3	82	57	
Subtotals	13	0	0	0	37	2,643	1,810	

Table B- 2b (continued)

	N	umber of Field I	Measured Featur	es by Terrain Type	(#)	Total measured	Total estimated
Erosional Feature Type	1. Schist	2. Melange	3. Alluvium	4. Coastal Belt	5. Franciscan	erosion (yds <sup>3</sup> )	sediment delivery (yds³)
Private Domain (plots/mi²)	1/1.8	0/1.2	0/0	0/0	0/8.6	1/11.6	1/11.6
Debris Slide (DL)	0	0	0	0	0	0	0
Debris Flow Source (DF)	0	0	0	0	0	0	0
Debris Torrent Track (TT)	0	0	0	0	0	0	0
Bank Erosion (BE)	0	0	0	0	0	0	0
Road related gully (GU)	1	0	0	0	0	12	6
Non road related gully (GU)	1	0	0	0	0	30	30
Stream Crossing (XI)	1	0	0	0	0	16	16
Surface Erosion (SE)	0	0	0	0	0	0	0
Active Earthflow (SDS)	0	0	0	0	0	0	0
Channel Incision (CI)	0	0	0	0	0	0	0
Subtotals	3	0	0	0	0	58	52
Total # of features, erosion and delivery volumes for all domains	16	0	0	0	37	2,701	1,862
Total # of plots/terrain type area in mi <sup>2</sup>	3/8	1/3.4	0/0	0/0	6/77.8	10/89.2	10/89.2

Table B-2c. Total measured erosion and sediment delivery measured in the 73 sample plots by ownership, terrain type and erosional feature type for the Soda Fork subwatershed, Upper Eel River watershed study area.

English LEngth of The		Number of Field	Measured Featu (#)	res by Terrain Type		Total measured erosion	Total estimated sediment delivery	
Erosional Feature Type	1. Schist	2. Melange	3. Alluvium	4. Coastal Belt	5. Franciscan	(yds <sup>3</sup> )	(yds <sup>3</sup> )	
Public Domain (plots/mi²)	1/31.5	0/0.7	0/0	0/5.1	0/0	1/37.3	1/37.3	
Debris Slide (DL)	5	0	0	0	0	222	189	
Debris Flow Source (DF)	0	0	0	0	0	0	0	
Debris Torrent Track (TT)	0	0	0	0	0	0	0	
Bank Erosion (BE)	5	0	0	0	0	49	49	
Road related gully (GU)	0	0	0	0	0	0	0	
Non road related gully (GU)	2	0	0	0	0	164	11	
Stream Crossing (XI)	0	0	0	0	0	0	0	
Channel Incision (CI)	2	0	0	0	0	22	22	
Surface Erosion (SE)	0	0	0	0	0	0	0	
Subtotals	14	0	0	0	0	457	271	

Table B- 2c (continued)

	N	umber of Field I	Measured Featur	es by Terrain Type	(#)	Total measured	Total estimated
Erosional Feature Type	1. Schist	2. Melange	3. Alluvium	4. Coastal Belt	5. Franciscan	erosion (yds <sup>3</sup> )	sediment delivery (yds³)
Private Domain (plots/mi²)	1/7.6	0/2.4	0/0	3/12.2	0/0.6	4/22.8	4/22.8
Debris Slide (DL)	4	0	0	7	0	10,669	2,005
Debris Flow Source (DF)	0	0	0	0	0	0	0
Debris Torrent Track (TT)	0	0	0	0	0	0	0
Bank Erosion (BE)	2	0	0	3	0	134	119
Road related gully (GU)	0	0	0	0	0	0	0
Non road related gully (GU)	1	0	0	2	0	120	100
Stream Crossing (XI)	0	0	0	0	0	0	0
Surface Erosion (SE)	0	0	0	0	0	0	0
Active Earthflow (SDS)	0	0	0	0	0	0	0
Channel Incision (CI)	0	0	0	4	0	40	40
Subtotals	7	0	0	16	0	10,963	2,264
Total # of features, erosion and delivery volumes for all domains	21	0	0	16	0	11,420	2,535
Total # of plots/terrain type area in mi <sup>2</sup>	2/39.1	0/3.1	0/0	3/17.3	0/0.6	5/60.1	5/60.1

Table B-2d. Total measured erosion and sediment delivery measured in the 73 sample plots by ownership, terrain type and erosional feature type for the Tomki Creek subwatershed, Upper Eel River watershed study area.

		Number of Field		Total measured erosion	Total estimated sediment delivery		
Erosional Feature Type	1. Schist	2. Melange	3. Alluvium	4. Coastal Belt	5. Franciscan	(yds <sup>3</sup> )	(yds <sup>3</sup> )
Public Domain (plots/mi²)	1/15	0/2	0/0	5/5.4	2/21.3	8/43.7	8/43.7
Debris Slide (DL)	6	0	0	17	8	6,620	5,048
Debris Flow Source (DF)	0	0	0	0	0	0	0
Debris Torrent Track (TT)	0	0	0	0	0	0	0
Bank Erosion (BE)	2	0	0	13	6	1,061	1,056
Road related gully (GU)	0	0	0	0	1	18	18
Non road related gully (GU)	0	0	0	1	2	48	48
Stream Crossing (XI)	0	0	0	0	1	11	11
Channel Incision (CI)	0	0	0	1	3	369	369
Surface Erosion (SE)	0	0	0	0	0	0	0
Subtotals	8	0	0	32	21	8,127	6,550

Table B- 2d (continued)

	N	umber of Field I	Measured Featur	es by Terrain Type	(#)	Total measured	Total estimated
Erosional Feature Type	1. Schist	2. Melange	3. Alluvium	4. Coastal Belt	5. Franciscan	erosion (yds <sup>3</sup> )	sediment delivery (yds <sup>3</sup> )
Private Domain (plots/mi²)	0/1	7/21	0/0.1	5/11.3	8/122.9	20/156.3	20/156.3
Debris Slide (DL)	0	31	0	12	27	30,798	10,822
Debris Flow Source (DF)	0	0	0	0	0	0	0
Debris Torrent Track (TT)	0	0	0	0	0	0	0
Bank Erosion (BE)	0	16	0	13	48	3,161	3,049
Road related gully (GU)	0	0	0	4	7	415	404
Non road related gully (GU)	0	7	0	3	6	707	672
Stream Crossing (XI)	0	0	0	6	7	253	248
Surface Erosion (SE)	0	0	0	0	0	0	0
Active Earthflow (SDS)	0	3	0	0	4	1,344	175
Channel Incision (CI)	0	9	0	2	15	538	530
Subtotals	0	66	0	40	114	37,216	15,900
Total # of features, erosion and delivery volumes for all domains	8	66	0	72	135	37,216	22,450
Total # of plots/terrain type area in mi <sup>2</sup>	1/16.1	7/23	0/0	10/16.7	10/144.2	28/200	28/200

Table B-2e. Total measured erosion and sediment delivery measured in the 73 sample plots by ownership, terrain type and erosional feature type for the Upper Main Eel subwatershed, Upper Eel River watershed study area.

		Number of Field	Measured Featu (#)	res by Terrain Type	:	Total measured erosion	Total estimated sediment delivery (yds³)	
Erosional Feature Type	1. Schist	2. Melange	3. Alluvium	4. Coastal Belt	5. Franciscan	(yds <sup>3</sup> )		
Public Domain (plots/mi²)	7/75	6/34.9	3/3.6	0/0	6/52.4	22/165.9	22/165.9	
Debris Slide (DL)	5	10	2	0	26	7,530	4,016	
Debris Flow Source (DF)	0	0	0	0	0	0	0	
Debris Torrent Track (TT)	0	0	0	0	0	0	0	
Bank Erosion (BE)	1	4	14	0	3	4,269	4,254	
Road related gully (GU)	2	1	2	0	1	92	76	
Non road related gully (GU)	7	0	0	0	0	180	157	
Stream Crossing (XI)	5	0	0	0	1	172	121	
Channel Incision (CI)	4	1	0	0	0	52	51	
Surface Erosion (SE)	7	2	1	0	3	553	280	
Subtotals	31	18	19	0	34	12,848	8,955	

Table B- 2e (continued)

	N	Number of Field I	Measured Featur	es by Terrain Type	(#)	Total measured	Total estimated
Erosional Feature Type	1. Schist	2. Melange	3. Alluvium	4. Coastal Belt	5. Franciscan	erosion (yds <sup>3</sup> )	sediment delivery (yds³)
Private Domain (plots/mi²)	2/7	0/0.1	0/0.9	0/0	0/5.1	2/13.1	2/13.1
Debris Slide (DL)	10	0	0	0	0	1,505	1,007
Debris Flow Source (DF)	1	0	0	0	0	444	400
Debris Torrent Track (TT)	0	0	0	0	0	0	0
Bank Erosion (BE)	3	0	0	0	0	18	18
Road related gully (GU)	2	0	0	0	0	17	17
Non road related gully (GU)	3	0	0	0	0	33	24
Stream Crossing (XI)	1	0	0	0	0	8	8
Surface Erosion (SE)	1	0	0	0	0	60	54
Active Earthflow (SDS)	0	0	0	0	0	0	0
Channel Incision (CI)	1	0	0	0	0	5	5
Subtotals	22	0	0	0	0	2,090	1,533
Total # of features, erosion and delivery volumes for all domains	53	18	19	0	34	14,938	10,488
Total # of plots/terrain type area in mi <sup>2</sup>	9/82	6/35	3/4.5	0/0	6/57.5	24/179	24/179

Table B-3 outlines the erosion and sediment delivery volumes by geology type by ownership for both small features <3,000 yds<sup>3</sup> and air photo-identified large features >3,000 yds<sup>3</sup>. The erosion and sediment delivery volumes for plot features were extrapolated to the entire Upper Eel River study area using the stratified random sampling method. Erosion and sediment delivery volumes for landslides are actual measurements from the air photo analyses.

Table B-3. Total past erosion and sediment delivery from small features  $<3,000 \text{ yds}^3$  and large features  $>3000 \text{ yds}^3$  by terrain type for each ownership in the Upper Eel River watershed study area

		Sr	nall Features (<3,000 y	vds <sup>3</sup> )	Lar	Large Features (>3,000 yds <sup>3</sup> )				
Ownership <sup>1</sup>	Terrain Type/ Geology	Non earthflow Erosion (yds <sup>3</sup> )	Non earthflow Sediment Delivery (yds <sup>3</sup> )	Earthflow Erosion (yds <sup>3</sup> )	Non earthflow erosion (yds <sup>3</sup> )	Non earthflow Sediment Delivery (yds <sup>3</sup> )	Earthflow erosion (yds³)			
Entire Upper	1	1,369,130	467,783	0	5,121,147	2,830,120	86,633			
Eel River	2	1,311,878	485,764	18,209	2,406,644	1,244,569	222,102			
study area	3	209,884	207,674	0	19,350	19,350	0			
	4	943,469	613,337	0	1,443,858	885,955	22,444			
	5	5,963,218	3,627,160	320,428	4,346,122	2,095,365	163,821			
	Totals	9,797,579	5,401,718	338,637	13,337,121	7,075,359	495,000			
Private	1	829,728	237,448	0	876,071	602,504	37,568			
	2	942,041	282,967	18,209	636,476	426,179	162,211			
	3	130,152	129,336	0	0	0	0			
	4	759,035	467,705	0	1,137,848	626,458	22,444			
	5	4,867,629	2,784,053	320,428	1,548,377	922,885	106,570			
	Totals	7,528,585	3,901,509	338,637	4,198,772	2,578,026	328,793			
Public	1	539,402	230,335	0	4,245,076	2,227,615	49,065			
	2	369,837	202,797	0	1,770,168	818,391	59,891			
	3	79,732	78,338	0	19,350	19,350	0			
	4	184,434	145,632	0	306,010	259,497	0			
Ī	5	1,095,589	843,107	0	2,797,765	1,172,480	57,251			
	Totals	2,268,994	1,500,209	0	9,138,349	4,497,333	166,207			

<sup>&</sup>lt;sup>1</sup> Total erosion and sediment yield for plot features (<3,000 yds<sup>2</sup>) for each ownership will not add up to the total yield for the entire Upper Eel River study area because each ownership is treated as a separate, smaller sample population, which is applied to each domain.

Table B-3a. Total past erosion and sediment delivery from small features <3,000 yds<sup>3</sup> and large features >3000 yds<sup>3</sup> by terrain type for each ownership in the Outlet Creek subwatershed, Upper Eel River watershed study area

		Sr	nall Features (<3,000 y	$vds^3$ )	Lar	ge Features (>3,000 yds	s <sup>3</sup> )
Ownership <sup>1</sup>	Terrain Type/ Geology	Non earthflow Erosion (yds³)	Non earthflow Sediment Delivery (yds³)	Earthflow Erosion (yds³)	Non earthflow erosion (yds³)	Non earthflow Sediment Delivery (yds³)	Earthflow erosion (yds³)
Private	1	0	0	0	0	0	0
	2	0	0	0	0	0	0
	3	120,582	119,826	0	0	0	0
	4	0	0	0	491,310	253,948	0
	5	1,518,862	1,068,638	0	387,636	230,538	84,114
	Sub -total	1,639,444	1,188,464	0	878,946	484,486	84,114
Public	1	0	0	0	0	0	0
	2	0	0	0	0	0	0
	3	0	0	0	0	0	0
	4	0	0	0	0	0	0
	5	0	0	0	0	0	0
	Sub -total	0	0	0	0	0	0
Totals		1,639,444	1,188,464	0	878,946	484,486	84,114

<sup>&</sup>lt;sup>1</sup> Total erosion and sediment yield for plot features <3,000 yds<sup>2</sup> for each subwatershed will not add up to the total yield for the entire Upper Eel River basin because each ownership is treated as a separate, smaller sample population which is applied to each domain.

Table B-3b. Total past erosion and sediment delivery from small features <3,000 yds<sup>3</sup> and large features >3000 yds<sup>3</sup> by terrain type for each ownership in the Rice Fork subwatershed, Upper Eel River watershed study area

		Sı	nall Features (<3,000 y	$(ds^3)$	Lar	ge Features (>3,000 yds	s <sup>3</sup> )
Ownership <sup>1</sup>	Terrain Type/ Geology	Non earthflow Erosion (yds³)	Non earthflow Sediment Delivery (yds³)	Earthflow Erosion (yds³)	Non earthflow erosion (yds <sup>3</sup> )	Non earthflow Sediment Delivery (yds³)	Earthflow erosion (yds³)
Private	1	1,624	1,456	0	3,968	1,190	0
	2	0	0	0	0	0	0
	3	0	0	0	0	0	8,505
	4	0	0	0	0	0	0
	5	0	0	0	72,019	25,101	5,478
	Sub -total	1,624	1,456	0	75,987	26,291	13,983
Public	1	22,658	13,062	0	59,624	15,121	830
	2	0	0	0	29,198	8,759	6,947
	3	0	0	0	0	0	0
	4	0	0	0	0	0	0
	5	382,299	270,928	0	1,088,179	438,533	42,808
	Sub -total	404,957	283,990	0	1,177,001	462,413	50,585
Totals		406,581	285,446	0	1,252,988	488,704	64,568

<sup>&</sup>lt;sup>1</sup> Total erosion and sediment yield for plot features <3,000 yds<sup>2</sup> for each subwatershed will not add up to the total yield for the entire Upper Eel River basin because each ownership is treated as a separate, smaller sample population which is applied to each domain.

Table B-3c. Total past erosion and sediment delivery from small features  $<3,000 \text{ yds}^3$  and large features  $>3000 \text{ yds}^3$  by terrain type for each ownership in the Soda Creek subwatershed, Upper Eel River watershed study area

		Sr	nall Features (<3,000 y	$vds^3$ )	Larg	ge Features (>3,000 yds	s <sup>3</sup> )
Ownership <sup>1</sup>	Terrain Type/ Geology	Non earthflow Erosion (yds³)	Non earthflow Sediment Delivery (yds³)	Earthflow Erosion (yds³)	Non earthflow erosion (yds³)	Non earthflow Sediment Delivery (yds³)	Earthflow erosion (yds³)
Private	1	1.187.376	227.244	0	674.610	428,341	27,410
	2	0	0	0	103,509	87,836	28,376
	3	0	0	0	0	0	0
	4	45,316	19,012	0	459,319	211,874	14,614
	5	0	0	0	0	0	0
	Sub -total	1,232,692	246,256	0	1,237,438	728,051	70,400
Public	1	220,274	130,622	0	915,639	539,462	18,553
	2	0	0	0	0	0	2,,082
	3	0	0	0	0	0	0
	4	0	0	0	65,516	39,213	0
	5	0	0	0	0	0	0
	Sub -total	220,274	130,622	0	981,155	578,675	20,635
Totals		1,452,966	376,878	0	2,218,593	1,306,726	91,035

<sup>&</sup>lt;sup>1</sup> Total erosion and sediment yield for plot features <3,000 yds<sup>2</sup> for each subwatershed will not add up to the total yield for the entire Upper Eel River basin because each ownership is treated as a separate, smaller sample population which is applied to each domain.

Table B-3d. Total past erosion and sediment delivery from small features <3,000 yds<sup>3</sup> and large features >3000 yds<sup>3</sup> by terrain type for each ownership in the Tomki Creek subwatershed, Upper Eel River watershed study area

		Sr	nall Features (<3,000 y	vds <sup>3</sup> )	Larg	ge Features (>3,000 yds	s <sup>3</sup> )
Ownership <sup>1</sup>	Terrain Type/ Geology	Non earthflow Erosion (yds³)	Non earthflow Sediment Delivery (yds³)	Earthflow Erosion (yds³)	Non earthflow erosion (yds³)	Non earthflow Sediment Delivery (yds³)	Earthflow erosion (yds³)
Private	1	0	0	0	2,218,593	1,306,726	91,035
	2	804,243	241,575	15,546	532,967	338,343	125,330
	3	0	0	0	0	0	0
	4	230,615	147,105	0	187,220	160,636	7,830
	5	2,763,011	1,471,254	236,426	999,388	618,871	16,978
	Sub -total	3,797,869	1,859,934	251,972	1,767,475	1,160,510	157,742
Public	1	78,776	68,242	0	1,034,037	757,718	27,815
	2	0	0	0	48,526	21,396	5,824
	3	0	0	0	0	0	0
	4	93,342	73,704	0	240,494	220,285	0
	5	353,160	296,262	0	625,629	213,031	0
	Sub -total	525,278	438,208	0	1,948,686	1,212,430	33,639
Totals		4,323,147	2,298,142	251,972	3,716,161	2,372,940	191,381

<sup>&</sup>lt;sup>1</sup> Total erosion and sediment yield for plot features <3,000 yds<sup>2</sup> for each subwatershed will not add up to the total yield for the entire Upper Eel River basin because each ownership is treated as a separate, smaller sample population which is applied to each domain.

Table B-3e. Total past erosion and sediment delivery from small features <3,000 yds<sup>3</sup> and large features >3000 yds<sup>3</sup> by terrain type for each ownership in the Upper Main Eel subwatershed, Upper Eel River watershed study area

		Sr	nall Features (<3,000 y	$rds^3$ )	Larg	ge Features (>3,000 yds	33)
Ownership <sup>1</sup>	Terrain Type/ Geology	Non earthflow Erosion (yds³)	Non earthflow Sediment Delivery (yds³)	Earthflow Erosion (yds³)	Non earthflow erosion (yds³)	Non earthflow Sediment Delivery (yds³)	Earthflow erosion (yds³)
Private	1	112,860	82,782	0	149,593	130,312	2,554
	2	0	0	0	0	0	0
	3	0	0	0	0	0	0
	4	0	0	0	0	0	0
	5	0	0	0	89,334	48,374	0
	Sub -total	112,860	82,782	0	238,927	178,686	2,554
Public	1	288,399	74,193	0	2,235,776	915,314	1,867
	2	378,339	207,459	0	1,692,444	788,235	45,038
	3	79,732	78,338	0	19,350	19,350	0
	4	0	0	0	0	0	0
	5	332,964	253,833	0	1,083,937	520,916	14,442
	Sub -total	1,079,433	613,823	0	5,031,507	2,243,815	61,347
Totals		1,192,293	696,605	251,972	5,270,434	2,422,501	63,901

<sup>&</sup>lt;sup>1</sup> Total erosion and sediment yield for plot features <3,000 yds<sup>2</sup> for each subwatershed will not add up to the total yield for the entire Upper Eel River basin because each ownership is treated as a separate, smaller sample population which is applied to each domain.

Table B-4 identifies sediment yield rates by time period for <3,000 yds<sup>3</sup> small features and >3,000 yds<sup>3</sup> large features for the Upper Eel River basin. The results are divided into earthflow and non-earthflow contributions for the small and large sediment sources. The sediment delivery rates for plot features were extrapolated to the entire Upper Eel River study area using the stratified random sampling method. The separation in the time period corresponds with the approximate changes in the Forest Practice Rules.

Table B-4. Sediment yield (in yds³/mi²/year and tons/mi²/year) by ownership and time frame for small features <3,000 yds³ and large features >3,000 yds³ in the Upper Eel River watershed study area

Watershed	Ownersh	nip domain			Pre -197	0		Post-1970					
				features 0 yds³)		Features 00 yds³)		Small f (<3,00		Large Fo (>3,000			
			non Ef	EF	non Ef	EF	Total	non Ef	EF	non Ef	EF	Total	
Entire Upper Eel	Private	yds <sup>3</sup> /mi <sup>2</sup> /yr	243	20	188	24	475	105	10	45	6	166	
River watershed study area		tons/mi²/yr	374	30	289	37	730	162	16	70	9	257	
	Public	yds <sup>3</sup> /mi <sup>2</sup> /yr	70	0	311	9	390	73	0	128	7	207	
		tons/mi <sup>2</sup> /yr	107	0	479	14	601	112	0	197	10	319	
	Totals	yds <sup>3</sup> /mi <sup>2</sup> /yr	160	10	247	17	434	90	5	85	6	186	
		tons/mi²/yr	247	16	380	26	669	138	8	130	10	286	
	Del este	yds <sup>3</sup> /mi <sup>2</sup> /yr	107	0	94	18	219	130	0	9	0	139	
	Private	tons/mi²/yr	165	0	144	28	337	200	0	14	0	214	
0.4.6.1	D 1.1	yds <sup>3</sup> /mi <sup>2</sup> /yr	0	0	0	0	0	0	0	0	0	0	
Outlet Creek P	Public	tons/mi²/yr	0	0	0	0	0	0	0	0	0	0	
	Sub-	yds <sup>3</sup> /mi <sup>2</sup> /yr	104	0	91	18	213	127	0	9	0	135	
	totals	tons/mi²/yr	161	0	140	27	328	195	0	13	0	208	

Table B-4 (continued)

Watershed	Ownersh	nip domain			Pre -197	0				Post-1970		
				features 0 yds³)		Features 00 yds <sup>3</sup> )		Small f (<3,00		Large Fo (>3,000		
			non Ef	EF	non Ef	EF	Total	non Ef	EF	non Ef	EF	Total
	Duissata	yds <sup>3</sup> /mi <sup>2</sup> /yr	4	0	30	0	34	0.4	0	40	35	76
	Private	tons/mi <sup>2</sup> /yr	6	0	47	0	52	0.7	0	61	54	117
D: E 1	D 1.11	yds <sup>3</sup> /mi <sup>2</sup> /yr	43	0	120	10	173	70	0	69	10	149
Rice Fork	Rice Fork Public	tons/mi²/yr	66	0	185	15	267	107	0	106	16	230
	Sub-	yds³/mi²/yr	38	0	109	9	155	61	0	65	14	140
	totals	tons/mi²/yr	58	0	167	13	239	93	0	101	21	215
	Deissata	yds <sup>3</sup> /mi <sup>2</sup> /yr	132	0	861	49	1042	201	0	179	48	428
	Private	tons/mi²/yr	203	0	0	75	1605	310	0	276	73	659
0.1.0.1	D 11	yds <sup>3</sup> /mi <sup>2</sup> /yr	0	0	391	2	393	103	0	111	15	229
Soda Creek	Public	tons/mi²/yr	0	0	602	3	605	159	0	171	23	353
	Sub-	yds <sup>3</sup> /mi <sup>2</sup> /yr	50	0	569	1	621	140	0	137	27	305
	totals	tons/mi²/yr	77	0	877	2	955	216	0	211	42	469

Table B-4 (continued)

Watershed	Ownersh	nip domain			Pre -197	0				Post-1970		
				features 0 yds³)		Features 00 yds <sup>3</sup> )		Small f (<3,00		Large Fo (>3,000		
			non Ef	EF	non Ef	EF	Total	non Ef	EF	non Ef	EF	Total
	Deissata	yds <sup>3</sup> /mi <sup>2</sup> /yr	304	33	180	29	546	82	18	60	4	163
	Private	tons/mi <sup>2</sup> /yr	468	51	277	45	841	126	28	92	6	252
T 1: C 1	D 1.11	yds <sup>3</sup> /mi <sup>2</sup> /yr	260	0	703	26	989	65	0	196	0	261
Tomki Creek	Public	tons/mi²/yr	401	0	1083	40	1524	100	0	301	0	401
	Sub-	yds <sup>3</sup> /mi <sup>2</sup> /yr	294	0	294	29	617	78	0	90	3	171
	totals	tons/mi²/yr	453	0	453	44	950	120	0	138	4	263
	Dei 111	yds <sup>3</sup> /mi <sup>2</sup> /yr	112	0	371	0	483	87	0	73	6	167
	Private	tons/mi²/yr	172	0	571	0	743	134	0	113	9	257
Hansa Mais Est	D 1.1	yds <sup>3</sup> /mi <sup>2</sup> /yr	40	0	287	6	333	74	0	144	5	223
Upper Main Eel	Public	tons/mi²/yr	61	0	442	10	514	114	0	222	8	344
	Sub-	yds <sup>3</sup> /mi <sup>2</sup> /yr	45	0	293	6	344	75	0	139	5	219
	totals	tons/mi²/yr	69	0	452	9	530	115	0	214	8	337

 $<sup>^{1}</sup>$  Total erosion and sediment yield for plot features <3,000 yds $^{2}$  for each ownership will not add up to the total yield for the entire Upper Eel River basin study area because each ownership is treated as a separate, smaller sample population which is applied to each domain.

Table B-5 compares pre-1970 and post-1970 sediment volumes for management and non-management land use association on private and public lands.

Ownership		Total Yield by Time Per Sediment Yield		Total Yield by Time Perio Sediment Yiel	
		Non Earthflow	Earthflow	Non Earthflow	Earthflow
		Basir	- n-wide		
Private	Pre-1970 (30 years)	2,299,049 (32%)	27,961 (<1%)	2,342,797 (33%)	439,219 (6%)
	Post-1970 (34 years)	698,328 (10%)	3,703 (<1%)	1,139,359 (16%)	202,832 (3%)
	Subtotals	2,997,377 (42%)	31,664 (<1%)	3,482,156 (49%)	642,051 (9%)
Public	Pre-1970 (30 years)	711,708 (12%)	10,516 (<1%)	3,044,859 (49%)	80,159 (1%)
	Post-1970 (34 years)	612,852 (10%)	8,748 (<1%)	1,628,123 (26%)	66,785 (1%)
	Subtotals	1,324,560 (22%)	120,242 (<1%)	4,672,982 (75%)	146,949 (2%)
Basin-wide To	tal	4,321,937 (32%)	50,928 (<1%)	8,155,138 (61%)	789,000 (6%)
		Outlet Creek	subwatershed		
Private	Pre-1970 (30 years)	304,216 (17%)	11,330 (<1%)	633,198 (36%)	72,784 (4%)
	Post-1970 (34 years)	279,910 (16%)	0	455,625 (26%)	0
	Subtotals	584,126 (33%)	11,330 (<1%)	1,088,823 (62%)	72,784 (4%)
Public	Pre-1970 (30 years)	0	0	0	0
	Post-1970 (34 years)	0	0	0	0
	Subtotals	0	0	0	0
Totals for the	Outlet Creek subwatershed	584,126 (33%)	11,330 (<1%)	1,088,823 (62%)	72,784 (4%)

Table B-5 (continued)

Ownership	,	Total Yield by Time Per Sediment Yield		Total Yield by Time Perio Sediment Yiel	
		Non Earthflow	Earthflow	Non Earthflow	Earthflow
		Rice Fork s	ubwatershed		
Private	Pre-1970 (30 years)	3,556 (9%)	0	8,279 (20%)	0
	Post-1970 (34 years)	8,980 (21%)	0	6,933 (16%)	13,984 (34%)
	Subtotals	12,536 (30%)	0	15,212 (36%)	13,984 (34%)
Public	Pre-1970 (30 years)	38,903 (5%)	2,682 (<1%)	341,247 (43%)	20,295 (3%)
	Post-1970 (34 years)	67,388 (8%)	830 (<1%)	298,866 (37%)	26,779 (3%)
	Subtotals	106,291 (13%)	3,512 (<1%)	640,113 (80%)	47,074 (6%)
Totals for the I	Rice Fork subwatershed	118,827 (14%)	3,512 (<1%)	655,325 (78%)	61,058 (7%)
		Soda Creek s	subwatershed		
Private	Pre-1970 (30 years)	206,403 (20%)	10,526 (1%)	472,939 (45%)	22,972 (2%)
	Post-1970 (34 years)	220,572 (21%)	0	74,392 (7%)	36,901 (4%)
	Subtotals	426,975 (41%)	10,526 (1%)	547,331 (52%)	59,873 (6%)
Public	Pre-1970 (30 years)	73,674 (10%)	0	363,916 (50%)	1,974 (<1%)
	Post-1970 (34 years)	160,228 (22%)	639 (<1%)	111,478 (15%)	18,022 (2%)
	Subtotals	233,902 (32%)	639 (<1%)	475,394 (65%)	19,996 (3%)
Totals for the S	Soda Creek subwatershed	660,877 (37%)	11,165 (<1)	1,022,725 (58%)	79,869 (5%)

Table B-5 (continued)

Ownership	mucu)	Total Yield by Time Per Sediment Yield		Total Yield by Time Perio Sediment Yiel	
		Non Earthflow	Earthflow	Non Earthflow	Earthflow
		Tomki Creek	subwatershed		
Private	Pre-1970 (30 years)	1,270,622 (37%)	6,105 (<1%)	997,098 (29%)	288,046 (8%)
	Post-1970 (34 years)	206,090 (6%)	3,703 (<1%)	546,634 (16%)	111,859 (3%)
	Subtotals	1,476,712 (43%)	9,808 (<1%)	1,543,732 (45%)	399,905 (12%)
Public	Pre-1970 (30 years)	256,799 (15%)	971 (<1%)	1,006,542 (60%)	32,668 (2%)
	Post-1970 (34 years)	105,203 (6%)	0	282,093 (17%)	0
	Subtotals	362,002 (21%)	971 (<1%)	1,288,635 (77%)	32,668 (2%)
Totals for the T	Tomki Creek subwatershed	1,838,714 (36%)	10,779 (<1%)	2,832,367 (55%)	432,573 (9%)
		Upper Main E	el subwatershed		
Private	Pre-1970 (30 years)	154,450 (59%)	0	35,261 (13%)	0
	Post-1970 (34 years)	41,103 (15%)	0	30,654 (12%)	2,554 (1%)
	Subtotals	195,553 (74%)	0	65,915 (25%)	2,554 (1%)
Public	Pre-1970 (30 years)	290,888 (10%)	6,863 (<1%)	1,336,805 (46%)	25,221 (1%)
	Post-1970 (34 years)	294,600 (10%)	7,279 (<1%)	935,344 (32%)	21,984 (1%)
	Subtotals	585,488 (20%)	14,142 (<1%)	2,272,149 (78%)	47,205 (2%)
Totals for the U	Jpper Main Eel subwatershed	781,041 (25%)	14,142 (<1%)	2,338,064 (73%)	49,759 (2%)

Table B-6 identifies sediment delivery rates by management practice and ownership for small features ( $<3,000 \text{ yds}^3$ ) and large features ( $>3,000 \text{ yds}^3$ ). These results are also separated by earthflow and non-earthflow contributions.

Table B-6. Sediment yield (in yds³/mi²/year, tons/mi²/year and %) by ownership and primary land use association for small and large features in the Upper Eel River watershed study area

					Non E	arthflow					Earthf	low			Total
	main (Private a nership) (mi²)	nd Public	No land use	Road Related	Timber harvest	Ag/ Grazin g	Fire	Total non EF sediment yield	No land use	Road Related	Timber harvest	Ag/ Grazin g	Fire	Total EF sediment yield	sediment yield (non EF+ EF)
	Plot <3,000	yds <sup>3</sup> /mi <sup>2</sup> /yr	94	26	30	19	0	170	15	0	0	0	0	15	184
	yds <sup>3</sup> sediment	tons/mi <sup>2</sup> /yr	145	40	46	30	0	261	23	0	0	0	0	23	284
ui <sup>2</sup> )	sources	%	56	15	18	11	0	100	100	0	0	0	0	100	100
4	<b>3.</b> 3,000yds <sup>3</sup>	yds <sup>3</sup> /mi <sup>2</sup> /yr	56	22	30	2	2	112	12	1	0	0	0	13	125
(359.	sediment	tons/mi <sup>2</sup> /yr	86	35	47	3	3	173	18	2	0	0	0	20	192
Private	sources	%	50	20	27	2	1	100	91	9	0	0	0	100	100
Pri		yds <sup>3</sup> /mi <sup>2</sup> /yr	150	48	61	21	2	282	26	1	0	0	0	27	309
	Sub-total/ %	tons/mi²/yr	230	74	93	33	3	434	41	2	0	0	0	43	476
		%	53	17	22	8	<1	100	96	4	0	0	0	100	100

Table B-6 (continued)

	ne D-0 (commue	<u>u,                                      </u>			Non E	arthflow					Earthf	low			Total
	main (Private ar nership) (mi²)	nd Public	No land use	Road Related	Timber harvest	Ag/ Grazin g	Fire	Total non EF sediment yield	No land use	Road Related	Timber harvest	Ag/ Grazin g	Fire	Total EF sediment yield	sediment yield (non EF+ EF)
	Plot <3,000	yds <sup>3</sup> /mi <sup>2</sup> /yr	44	9	18	0	0	71	0	0	0	0	0	0	71
	yds <sup>3</sup> sediment	tons/mi <sup>2</sup> /yr	68	14	28	0	0	110	0	0	0	0	0	0	110
mi²)	sources	%	62	13	25	0	0	100	0	0	0	0	0	0	100
	>3000yds <sup>3</sup>	yds <sup>3</sup> /mi <sup>2</sup> /yr	177	23	12	<1	1	214	5	<1	<1	0	<1	5	219
Public (328.7	sediment	tons/mi <sup>2</sup> /yr	272	36	18	<1	2	330	7	<1	<1	0	<1	7	337
ıblic	sources	%	82	11	6	<1	<1	100	94	<1	<1	0	5	100	100
Pu		yds <sup>3</sup> /mi <sup>2</sup> /yr	221	33	30	<1	1	285	5	<1	<1	0	<1	5	290
	Sub-total/ %	tons/mi <sup>2</sup> /yr	340	50	46	<1	2	439	7	<1	<1	0	<1	7	446
		%	77	11	10	<1	<1	100	94	<1	<1	0	5	100	100
	Plot <3,000	yds <sup>3</sup> /mi <sup>2</sup> /yr	70	18	24	10	0	123	8	0	0	0	0	8	130
	yds <sup>3</sup> sediment	tons/mi <sup>2</sup> /yr	108	28	38	16	0	189	12	0	0	0	0	12	201
$mi^2$ )	sources	%	57	15	20	8	0	100	100	0	0	0	0	100	100
88.1	PWA	yds <sup>3</sup> /mi <sup>2</sup> /yr	114	23	22	1	1	160	8	0.7	<1	0	<1	9	170
ge (6	>3,000yds <sup>3</sup> sediment	tons/mi <sup>2</sup> /yr	175	35	33	2	2	247	13	1	<1	0	<1	14	261
n-wi	sources	%	71	14	13	1	1	100	93	7	<1	0	<1	100	100
Basin-wide (688.1		yds <sup>3</sup> /mi <sup>2</sup> /yr	184	41	46	11	1.5	283	16	0.7	<1	0	<1	17	300
	Sub-total/ %	tons/mi²/yr	283	63	71	17	2	436	25	1	<1	0	<1	26	462
		%	65	14	16	4	<1	100	95	4	<1	0	<1	100	100

Table B-6a. Sediment yield (in yds³/mi²/year, tons/mi²/year and %) by ownership and primary land use association for small and large features in the Outlet Creek subwatershed, Upper Eel River watershed study area

Do	main (Private	and Public			Non	Earthflow					Earthfl	ow		Total
	nership) (mi <sup>2</sup> )		No land use	Road Related	Timber harvest	Ag/ Grazing	Fire	Total non EF sediment yield	No land use	Road Related	Timber harvest	Ag/ Grazing	Total EF sediment yield	sediment yield (non EF+ EF)
	Plot	yds <sup>3</sup> /mi <sup>2</sup> /yr	80	20	10	10	0	119	0	0	0	0	0	119
	<3,000 yds <sup>3</sup>	tons/mi <sup>2</sup> /yr	122	31	16	15	0	184	0	0	0	0	0	184
	sediment sources	%	67	17	8	8	0	100	0	0	0	0	0	100
ها		yds <sup>3</sup> /mi <sup>2</sup> /yr	30	11	5	2	0	49	73	1	0	0	74	123
Private	>3,000yds <sup>3</sup> sediment	tons/mi <sup>2</sup> /yr	46	17	8	4	0	75	113	2	0	0	114	189
H	sources <sup>5</sup>	%	61	23	11	5	0	100	98	2	0	0	100	100
		yds <sup>3</sup> /mi <sup>2</sup> /yr	109	31	15	12	0	168	73	1	0	0	74	242
	Sub-total/ %	tons/mi²/yr	168	48	24	18	0	259	113	2	0	0	114	373
		%	65	19	9	7	0	100	98	2	0	0	100	100

Table B-6a (continued)

	<i>ole B-6a (contii</i> main (Private	,			Non	Earthflow					Earthfl	ow		Total
	nership) (mi²)		No land use	Road Related	Timber harvest	Ag/ Grazing	Fire	Total non EF sediment yield	No land use	Road Related	Timber harvest	Ag/ Grazing	Total EF sediment yield	sediment yield (non EF+ EF)
	Plot	yds <sup>3</sup> /mi <sup>2</sup> /yr	0	0	0	0	0	0	0	0	0	0	0	0
	<3,000 yds <sup>3</sup>	tons/mi <sup>2</sup> /yr	0	0	0	0	0	0	0	0	0	0	0	0
	sediment sources	%	0	0	0	0	0	0	0	0	0	0	0	0
၁	>3000yds <sup>3</sup>	yds <sup>3</sup> /mi <sup>2</sup> /yr	0	0	0	0	0	0	0	0	0	0	0	0
Public	sediment	tons/mi <sup>2</sup> /yr	0	0	0	0	0	0	0	0	0	0	0	0
	Sub-total/	%	0	0	0	0	0	0	0	0	0	0	0	0
		yds³/mi²/yr	0	0	0	0	0	0	0	0	0	0	0	0
	Sub-total/ %	tons/mi²/yr	0	0	0	0	0	0	0	0	0	0	0	0
		%	0	0	0	0	0	0	0	0	0	0	0	0
mi²)	Plot	yds <sup>3</sup> /mi <sup>2</sup> /yr	77	20	10	9	0	116	0	0	0	0	0	116
59.8	Plot <3,000 yds <sup>3</sup>	tons/mi <sup>2</sup> /yr	119	30	15	14	0	179	0	0	0	0	0	179
Totals (159.8 mi²)	sediment sources	%	67	17	8	8	0	100	0	0	0	0	0	100
\ Tot	PWA	yds <sup>3</sup> /mi <sup>2</sup> /yr	29	11	5	2	0	47	71	1	0	0	72	119
WA.	>3,000yds <sup>3</sup> sediment	tons/mi²/yr	45	17	8	4	0	73	110	2	0	0	111	184
CAL	sources <sup>5</sup>	%	61	23	11	5	0	100	98	2	0	0	100	100
Outlet Creek CALWA.		yds <sup>3</sup> /mi <sup>2</sup> /yr	106	31	14	12	0	164	71	1	0	0	72	236
tlet C	Sub-total/ %	tons/mi²/yr	164	47	23	18	0	252	110	2	0	0	111	363
Ou		%	65	19	9	7	0	100	98	2	0	0	100	100

Table B-6b. Sediment yield (in yds³/mi²/year, tons/mi²/year and %) by ownership and primary land use association for small and large features in the Rice Fork subwatershed, Upper Eel River watershed study area

					Non I	Earthflow				_	Earthflo	w		Total
	main (Private nership) (mi <sup>2</sup> )		No land use	Road Related	Timber harvest	Ag/ Grazing	Fire	Total non EF sediment yield	No land use	Road Related	Timber harvest	Ag/ Grazing	Total EF sediment yield	Total sediment yield (non EF+ EF)
	Plot	yds <sup>3</sup> /mi <sup>2</sup> /yr	0	<1	2	0	0	2	0	0	0	0	0	2
	<3,000 yds <sup>3</sup>	tons/mi²/yr	0	<1	3	0	0	3	0	0	0	0	0	3
	sediment sources	%	0	12	88	0	0	100	0	0	0	0	0	100
le te		yds <sup>3</sup> /mi <sup>2</sup> /yr	20	3	12	0	0	35	0	0	0	0	0	35
Private	>3,000yds <sup>3</sup> sediment	tons/mi <sup>2</sup> /yr	32	5	18	0	0	55	0	0	0	0	0	55
	sources <sup>5</sup>	%	58	9	33	0	0	100	0	0	0	0	0	100
	Sub-total/	yds <sup>3</sup> /mi <sup>2</sup> /yr	20	3	13	0	0	37	0	0	0	0	0	37
		tons/mi²/yr	32	5	21	0	0	58	0	0	0	0	0	58
		%	55	9	36	0	0	100	0	0	0	0	0	100

Table B-6b (continued)

140	le B-6b (conti	nucu)			Non	Earthflow					Earthfl	ow		Total
	main (Private nership) (mi <sup>2</sup>		No land use	Road Related	Timber harvest	Ag/ Grazing	Fire	Total non EF sediment yield	No land use	Road Related	Timber harvest	Ag/ Grazing	Total EF sediment yield	sediment yield (non EF+ EF)
	Plot	yds <sup>3</sup> /mi <sup>2</sup> /yr	45	10	3	0	0	57	0	0	0	0	0	57
	<3,000 yds <sup>3</sup>	tons/mi²/yr	69	15	5	0	0	88	0	0	0	0	0	88
	sediment sources	%	78	17	5	0	0	100	0	0	0	0	0	100
၁		yds <sup>3</sup> /mi <sup>2</sup> /yr	84	2	7	0	0	93	4	0	0	0	4	97
Public	>3000yds <sup>3</sup> sediment	tons/mi²/yr	130	3	11	0	0	143	7	0	0	0	7	150
	sources <sup>5</sup>	%	90	2	8	0	0	100	100	0	0	0	100	100
		yds <sup>3</sup> /mi <sup>2</sup> /yr	129	11	10	0	0	150	4	0	0	0	4	155
	Sub-total/ %	tons/mi²/yr	198	18	15	0	0	231	7	0	0	0	7	238
		%	86	8	6	0	0	100	100	0	0	0	100	100
	Plot	yds <sup>3</sup> /mi <sup>2</sup> /yr	39	8	3	0	0	50	0	0	0	0	0	50
$2 \mathrm{mi}^2$	<3,000 yds <sup>3</sup>	tons/mi²/yr	60	13	4	0	0	77	0	0	0	0	0	77
Total Rice Fork CALWAA (89.2 mi²)	sediment sources	%	78	17	5	0	0	100	0	0	0	0	0	100
WA	PWA	yds <sup>3</sup> /mi <sup>2</sup> /yr	76	2	8	0	0	86	4	0	0	0	4	90
CAI	>3,000yds <sup>3</sup> sediment	tons/mi²/yr	117	3	12	0	0	132	6	0	0	0	6	138
Fork	sources <sup>5</sup>	%	89	2	9	0	0	100	100	0	0	0	100	100
Rice		yds <sup>3</sup> /mi <sup>2</sup> /yr	115	10	10	0	0	136	4	0	0	0	4	140
<b>Total</b>	Sub-total/ %	tons/mi²/yr	177	16	16	0	0	209	6	0	0	0	6	215
		%	85	8	7	0	0	100	100	0	0	0	100	100

Table B-6c. Sediment yield (in yds³/mi²/year, tons/mi²/year and %) by ownership and primary land use association for small and large features in the Soda Creek subwatershed, Upper Eel River watershed study area

					Non E	arthflow					Eart	thflow			Total
	nain (Private a ership) (mi <sup>2</sup> )	nd Public	No land use	Road Related	Timber harvest	Ag/ Grazing	Fire	Total non EF sedimen t yield	No land use	Road Related	Timber harvest	Ag/ Grazing	Fire	Total EF sediment yield	sediment yield (non EF+ EF)
	Plot	yds <sup>3</sup> /mi <sup>2</sup> /yr	11	0	157	0	0	169	0	0	0	0	0	0	169
	<3,000 yds <sup>3</sup>	tons/mi²/yr	17	0	242	0	0	260	0	0	0	0	0	0	260
	sediment sources	%	7	0	93	0	0	100	0	0	0	0	0	0	100
e je		yds <sup>3</sup> /mi <sup>2</sup> /yr	336	115	21	0	28	499	35	7	0	0	0	42	541
Private	>3,000yds <sup>3</sup> sediment	tons/mi²/yr	518	176	32	0	42	768	54	11	0	0	0	65	834
	sources <sup>5</sup>	%	67	23	4	0	6	100	83	17	0	0	0	100	100
		yds <sup>3</sup> /mi <sup>2</sup> /yr	348	114	178	0	28	668	35	7	0	0	0	42	710
	Sub-total/ %	tons/mi²/yr	535	176	274	0	42	1028	54	11	0	0	0	65	1093
		%	52	17	27	0	4	100	83	17	0	0	0	100	100

Table B-6c (continued)

	te B-oc (comin	,			Non E	arthflow					Eart	hflow			Total
	nain (Private a ership) (mi²)	and Public	No land use	Road Related	Timber harvest	Ag/ Grazing	Fire	Total non EF sedimen t yield	No land use	Road Related	Timber harvest	Ag/ Grazing	Fire	Total EF sediment yield	sediment yield (non EF+ EF)
	Plot	yds <sup>3</sup> /mi <sup>2</sup> /yr	0	0	55	0	0	55	0	0	0	0	0	0	55
	<3,000 yds <sup>3</sup>	tons/mi²/yr	0	0	84	0	0	84	0	0	0	0	0	0	84
	sediment sources	%	0	0	100	0	0	100	0	0	0	0	0	0	100
ږ	>3000yds <sup>3</sup>	yds <sup>3</sup> /mi <sup>2</sup> /yr	199	33	11	0	0	242	6	0	0	0	2	8	250
Public	sediment	tons/mi²/yr	307	50	17	0	0	373	10	0	0	0	3	13	386
	sources <sup>5</sup>	%	82	13	5	0	0	100	74	0	0	0	26	100	100
		yds³/mi²/yr	199	33	65	0	0	297	6	0	0	0	2	8	305
	Sub-total/ %	tons/mi²/yr	307	50	101	0	0	458	10	0	0	0	3	13	470
		%	63	10	21	0	0	100	74	0	0	0	26	100	100
$\mathbf{i}^2$ )	Plot	yds <sup>3</sup> /mi <sup>2</sup> /yr	4	0	94	0	0	98	0	0	0	0	0	0	98
0.1 m	<3,000 yds <sup>3</sup>	tons/mi²/yr	7	0	144	0	0	151	0	0	0	0	0	0	151
Soda Creek CALWAA Total (60.1 mi²)	sediment sources	%	4	0	96	0	0	100	0	0	0	0	0	0	100
A To	PWA	yds <sup>3</sup> /mi <sup>2</sup> /yr	251	64	14	0	10	340	17	3	0	0	1	21	361
LWA	>3,000yds <sup>3</sup> sediment	tons/mi²/yr	387	98	22	0	16	523	27	4	0	0	2	33	556
k CA	sources <sup>5</sup>	%	74	19	4	0	3	100	81	13	0	0	6	100	100
Cree		yds³/mi²/yr	255	64	108	0	10	437	17	3	0	0	1	21	459
Soda	Sub-total/ %	tons/mi²/yr	393	98	167	0	16	674	27	4	0	0	2	33	707
رو	% <u>t</u>	%	58	15	25	0	2	100	81	13	0	0	6	100	100

Table B-6d. Sediment yield (in yds³/mi²/year, tons/mi²/year and %) by ownership and primary land use association for small and large features in the Tomki Creek subwatershed, Upper Eel River watershed study area

					Non I	Earthflow					Ear	thflow			Total
	ain (Private ar ership) (mi <sup>2</sup> )	nd Public	No land use	Road Related	Timber harvest	Ag/ Grazing	Fire	Total non EF sediment yield	No land use	Road Related	Timber harvest	Ag/ Grazing	Fire	Total EF sediment yield	Total sediment yield (non EF+ EF)
	Plot	yds <sup>3</sup> /mi <sup>2</sup> /yr	113	30	16	23	0	182	25	0	0	0	0	25	207
	<3,000 yds <sup>3</sup>	tons/mi <sup>2</sup> /yr	175	46	24	35	0	280	39	0	0	0	0	39	319
	sediment sources	%	62	17	9	12	0	100	100	0	0	0	0	100	100
e.		yds <sup>3</sup> /mi <sup>2</sup> /yr	41	16	57	2	<1	116	15	<1	0	0	0	15	131
Private	>3,000yds <sup>3</sup> sediment	tons/mi²/yr	63	25	87	4	<1	179	22	1	0	0	0	23	202
	sources <sup>5</sup>	%	35	14	49	2	<1	100	96	4	0	0	0	100	100
		yds <sup>3</sup> /mi <sup>2</sup> /yr	154	46	72	25	<1	298	40	<1	0	0	0	40	338
	Sub-total/ %	tons/mi²/yr	238	71	111	39	<1	459	61	1	0	0	0	62	521
		%	52	16	24	8	<1	100	98	2	0	0	0	100	100

Table B-6d (continued)

			Non Earthflow							Earthflow						
	main (Private ai nership) (mi <sup>2</sup> )	nd Public	No land use	Road Related	Timber harvest	Ag/ Grazing	Fire	Total non EF sediment yield	No land use	Road Related	Timber harvest	Ag/ Grazing	Fire	Total EF sediment yield	Total sediment yield (non EF+ EF)	
	Plot <3,000 yds <sup>3</sup> sediment sources	yds <sup>3</sup> /mi <sup>2</sup> /yr	83	46	28	0	0	157	0	0	0	0	0	0	157	
		tons/mi <sup>2</sup> /yr	128	70	43	0	0	241	0	0	0	0	0	0	241	
		%	53	29	18	0	0	100	0	0	0	0	0	0	100	
. <u>2</u>	>3000yds <sup>3</sup> sediment sources <sup>5</sup>	yds <sup>3</sup> /mi <sup>2</sup> /yr	369	21	34	1	8	434	12	<1	0	0	0	12	446	
Public		tons/mi²/yr	569	32	52	2	13	668	18	1	0	0	0	19	686	
		%	85	5	8	<1	2	100	97	3	0	0	0	100	100	
	Sub-total/ %	yds <sup>3</sup> /mi <sup>2</sup> /yr	452	66	62	1	8	590	12	<1	0	0	0	12	602	
		tons/mi²/yr	697	102	95	2	13	909	18	<1	0	0	0	19	927	
		%	77	11	10	<1	1	100	97	3	0	0	0	100	100	
mi²)	Plot <3,000 yds <sup>3</sup> sediment sources	yds <sup>3</sup> /mi <sup>2</sup> /yr	107	34	18	18	0	176	20	0	0	0	0	20	196	
(200 mi <sup>2</sup> )		tons/mi²/yr	164	52	28	27	0	272	30	0	0	0	0	30	302	
Total		%	61	19	10	10	0	100	100	0	0	0	0	100	100	
<	PWA >3,000yds <sup>3</sup> sediment sources <sup>5</sup>	yds <sup>3</sup> /mi <sup>2</sup> /yr	113	17	52	2	2	185	14	<1	0	0	0	14	200	
LWA		tons/mi²/yr	173	26	80	3	3	285	21	1	0	0	0	22	308	
CAI		%	61	9	28	1	1	100	96	4	0	0	0	100	100	
reek	Sub-total/ %	yds <sup>3</sup> /mi <sup>2</sup> /yr	219	51	70	20	20	362	33	<1	0	0	0	34	395	
Tomki Creek CALWA		tons/mi²/yr	338	78	108	31	31	557	52	1	0	0	0	53	610	
Tor		%	61	14	19	5	5	100	98	2	0	0	0	100	100	

Table B-6e. Sediment yield (in yds³/mi²/year, tons/mi²/year and %) by ownership and primary land use association for small and large features in the Upper Main Eel subwatershed, Upper Eel River watershed study area

				Non Earthflow							Earthflow						
Domain (Private and Public ownership) (mi <sup>2</sup> )			No land use	Road Related	Timber harvest	Ag/ Grazing	Fire	Total non EF sediment yield	No land use	Road Related	Timber harvest	Ag/ Grazing	Fire	Total EF sediment yield	Total sediment yield (non EF+ EF)		
	Plot <3,000 yds <sup>3</sup> sediment sources	yds <sup>3</sup> /mi <sup>2</sup> /yr	<1	1	97	0	0	99	0	0	0	0	0	0	99		
Private		tons/mi²/yr	<1	2	150	0	0	152	0	0	0	0	0	0	152		
		%	<1	1	99	0	0	100	0	0	0	0	0	0	100		
	>3,000yds <sup>3</sup> sediment sources <sup>5</sup>	yds <sup>3</sup> /mi <sup>2</sup> /yr	78	88	47	0	0	213	0	0	0	0	0	0	213		
		tons/mi²/yr	121	136	72	0	0	328	0	0	0	0	0	0	328		
Ь		%	37	41	22	0	0	100	0	0	0	0	0	0	100		
	Sub-total/	yds <sup>3</sup> /mi <sup>2</sup> /yr	79	89	144	0	0	312	0	0	0	0	0	0	312		
		tons/mi²/yr	121	137	222	0	0	480	0	0	0	0	0	0	480		
		%	25	29	46	0	0	100	0	0	0	0	0	0	100		

Table B-6e (continued)

Table B-be (continued)			Non Earthflow							Earthflow						
	main (Private a nership) (mi <sup>2</sup> )	nd Public	No land use	Road Related	Timber harvest	Ag/ Grazing	Fire	Total non EF sediment yield	No land use	Road Related	Timber harvest	Ag/ Grazing	Fire	Total EF sediment yield	Total sediment yield (non EF+ EF)	
	Plot <3,000	yds <sup>3</sup> /mi <sup>2</sup> /yr	45	2	11	0	0	58	0	0	0	0	0	0	58	
	yds <sup>3</sup> sediment	tons/mi²/yr	69	3	17	0	0	89	0	0	0	0	0	0	89	
	sources	%	77	4	19	0	0	100	0	0	0	0	0	0	100	
၂	>3000yds <sup>3</sup> sediment sources <sup>5</sup>	yds <sup>3</sup> /mi <sup>2</sup> /yr	169	33	9	<1	<1	211	2	0	<1	0	0	2	213	
Public		tons/mi <sup>2</sup> /yr	261	50	14	<1	<1	325	4	0	<1	0	0	4	329	
-		%	80	15	4	<1	<1	100	98	0	2	0	0	100	100	
	Sub-total/%	yds <sup>3</sup> /mi <sup>2</sup> /yr	214	35	20	<1	<1	269	2	0	<1	0	0	2	271	
		tons/mi²/yr	329	54	31	<1	<1	414	4	0	<1	0	0	4	418	
		%	79	13	7	<1	<1	100	98	0	2	0	0	100	100	
	Plot <3,000 yds <sup>3</sup> sediment sources	yds <sup>3</sup> /mi <sup>2</sup> /yr	41	2	17	0	0	61	0	0	0	0	0	0	61	
(179 mi²)		tons/mi <sup>2</sup> /yr	64	3	27	0	0	94	0	0	0	0	0	0	94	
(179		%	68	3	29	0	0	100	0	0	0	0	0	0	100	
otal	PWA >3,000yds <sup>3</sup> sediment sources <sup>5</sup>	yds <sup>3</sup> /mi <sup>2</sup> /yr	163	37	12	<1	<1	211	2	0	<1	0	0	2	213	
Eel T		tons/mi²/yr	250	56	18	<1	<1	326	4	0	<1	0	0	4	329	
ain		%	77	17	6	<1	<1	100	98	0	2	0	0	100	100	
Upper Main Eel Total	Sub-total/ %	yds <sup>3</sup> /mi <sup>2</sup> /yr	204	39	29	<1	<1	272	2	0	<1	0	0	2	274	
Upp		tons/mi²/yr	314	60	45	<1	<1	419	4	0	<1	0	0	4	423	
		%	75	14	11	<1	<1	100	98	0	2	0	0	100	100	