

Appendix E

Sensitivity Plots for the EFDC Toxics Model at the Impaired Shoreline Areas

Figure E-1. Comparison of TPAH concentrations at the Paleta Creek mouth for the baseline and sensitivity runs.	1
Figure E-2. Comparison of TPAH concentrations at the Chollas Creek mouth for the baseline and sensitivity runs.	1
Figure E-3. Comparison of TPAH concentrations at the Switzer Creek mouth for the baseline and sensitivity runs.	2
Figure E-4. Comparison of TPAH concentrations at the B-Street mouth for baseline and sensitivity runs.	2
Figure E-5. Comparison of TPAH concentrations at the Downtown Anchorage mouth for the baseline and sensitivity runs.	3




Figure E-1. Comparison of TPAH concentrations at the Paleta Creek mouth for the baseline and sensitivity runs.




Figure E-2. Comparison of TPAH concentrations at the Chollas Creek mouth for the baseline and sensitivity runs.




Figure E-3. Comparison of TPAH concentrations at the Switzer Creek mouth for the baseline and sensitivity runs.




Figure E-4. Comparison of TPAH concentrations at the B-Street mouth for baseline and sensitivity runs.




Figure E-5. Comparison of TPAH concentrations at the Downtown Anchorage mouth for the baseline and sensitivity runs.