



ELSEVIER

Available online at [www.sciencedirect.com](http://www.sciencedirect.com)

SciVerse ScienceDirect

Current Opinion in  
Environmental  
Sustainability

# Ecological stoichiometry and its implications for aquatic ecosystem sustainability

Patricia M Glibert

Aquatic ecosystems are increasingly stressed not only by increased nutrient loads (eutrophication) but also by changing forms and proportions of nutrients. Nutrient enrichment, composition and stoichiometry interact with aquatic food web dynamics in complex ways. Both algal species composition and emergent properties within species change with changing nutrient composition, in turn affecting food webs at all levels. Consumers further regulate – and may even accelerate – discrepancies in nutrient stoichiometry by various feedbacks, release, and recycling pathways. Stoichiometric regulation of aquatic ecosystem structure also occurs at the sediment interface via altered biogeochemical processes and benthic food webs when nutrient composition changes. Thus, multiple feedbacks serve to alter food web structure when nutrient loads are altered. Such feedbacks may also lead to conditions conducive to invasive species and altered stable states as illustrated for the San Francisco Bay Delta and the Rhine River.

## Address

University of Maryland Center for Environmental Science, Horn Point Laboratory, PO Box 775, Cambridge, MD 21613, United States

Corresponding author: Glibert, Patricia M ([glibert@umces.edu](mailto:glibert@umces.edu))

Current Opinion in Environmental Sustainability 2012, 4:1–6

This review comes from a themed issue on  
Aquatic and marine systems  
Edited by R Ian Perry, Alida Bundy, Eileen E Hofmann

Received 2 February 2012; Accepted 14 May 2012

1877-3435/\$ – see front matter  
© 2012 Elsevier B.V. All rights reserved.

<http://dx.doi.org/10.1016/j.cosust.2012.05.009>

## Introduction

Much has been written about the eutrophication of lakes, estuaries and coasts ([1<sup>••</sup>,2,3] and references therein). Anthropogenic activities have significantly altered nutrient loads and their composition, in turn profoundly affecting ecosystem health [4]. For example, well-documented effects of increased nutrient loads and eutrophication include loss of biodiversity, increased harmful algal blooms, and development of dead zones [5,6]. Beyond eutrophication, changes in nutrient loads are impacting ecosystem dynamics in complex ways. Changes in nutrient form (chemical state, oxidized vs reduced, organic vs inorganic, dissolved vs particulate) and the proportion of different elements (C, N, P, Si, etc.) have both proximal

and ultimate effects on ecosystems (*sensu* [7]). The stoichiometry (proportions) of nutrients, especially nitrogen (N) and phosphorus (P), in many aquatic systems is changing as a result of both increasing nutrient loads (often dominated by N) on the one hand, and management efforts focused on single nutrient control (often P) on the other. At the primary producer level, many of these effects are well understood, framed largely in the concepts of Liebig's law of the minimum and Redfield stoichiometry (reviewed by [8<sup>•</sup>,9,10<sup>•</sup>]). At the proximal scale, here defined as the scale of growth of primary producers, nutrient form and concentration affect the cell physiological processes of uptake, assimilation, and those, in turn, regulate the internal concentrations that regulate growth (e.g. [11]). At the ultimate scale, the total load and balance of nutrient elements have effects that propagate through the food web, with the potential of transforming ecosystems to new stable states. The regulation of food web structure by nutrient stoichiometry is further controlled by the various and complex feedbacks, release, and recycling pathways that that are all fundamentally constrained by nutrient load, form or stoichiometry. Just as different elemental ratios may affect the composition of the primary producers, different nutrient requirements of higher trophic levels will have an impact on their ability to thrive as community composition changes at the base. The latter is the emphasis of this brief communication.

## Stoichiometry, biotic feedbacks, and biogeochemistry

An ecological stoichiometric approach is based on the transfer of elements through the food web, not just the flow of carbon (C) [12<sup>••</sup>]. Generally, homeostatic mechanisms keep the acquisition of materials and energy in balance with the cellular growth demands of primary producers and the general consistency of Redfield stoichiometry in phytoplankton is corroborative (e.g. [10<sup>•</sup>]). Ecological stoichiometry recognizes that at the base of the food web, the elemental composition of the primary producers is affected by nutrient composition whether nutrients are limiting or not. While the total nutrient load of the nutrient considered to be limiting for phytoplankton production may control overall biomass, the composition, both in form of nutrients and the proportion of different nutrient elements affects the composition of the primary producers. When nutrient loads change in amount, form or proportion, cellular adjustments in acquisition efficiency lead to altered emergent properties such as the proportions of ribosomes, enzyme activities, gene regulation, cellular pigmentation complement, and

## 2 Aquatic and marine systems

cellular elemental composition [12<sup>••</sup>]. These properties have implications for competition and species success, leading, eventually, to changes in species dominants and biodiversity. In addition, properties such as lipid composition, toxin production, cell membrane thickness, and other chemical constituents, which are also, at least partly, functions of cellular elemental availability, can alter the quality of food for consumers, in some cases turning 'good' food to 'bad' [13,14]. Production of toxins in algae, for example, often occurs when nutrient stoichiometry is not in classic ('Redfield') stoichiometric proportions [14,15]. Toxic species can be harmful to higher trophic levels, disrupting normal ecosystem function. The dominance of such algae can result in a failure of normal predator-prey interactions, which in turn enhances the transfer of nutrients that sustain such species at the expense of competing algal species [16–18].

While primary producers may be flexible to some degree in their elemental composition [19<sup>••</sup>,20], heterotrophs are typically far less flexible [10<sup>•</sup>,12<sup>••</sup>,21<sup>••</sup>]. The nutritional quality of the food at the base of the food web affects all aspects of the abilities of consumers to meet their nutritional and reproductive demands. Nutrient stoichiometry and food quality can have differential effects on consumers depending on their life stage, for example, larval or adult [22–25]. Larvae would be expected to have higher P demands, for example, than adults, owing to their higher growth rates [22]. Heterotrophs typically maintain a more strict stoichiometry than phytoplankton through their various excretion and release mechanisms [12<sup>••</sup>,26], and the relationship between their biomass elemental requirements and the proportion of those elements in their food determines the extent to which they are a sink or a source of N, P or C [12<sup>••</sup>,27]. In general, as one moves up the food web, stoichiometric constraints on grazers increase, as does the organismal requirement for P, owing to fixed stoichiometry in body tissue, especially bone [12<sup>••</sup>]. Heterotrophs thus can exert significant control on the N and P available for primary producers [12<sup>••</sup>,28].

The biota thus modify the environment through nutrient uptake and regeneration which differs from species to species based on their elemental requirements and the nutrient substrates available to them as dissolved nutrients or as food. In addition to the biotic feedbacks and regulatory controls whereby individual organisms maintain their stoichiometry under mass balance constraints, at the ecosystem level, growth of different species also modifies the physical and chemical environment through alterations in physical habitat (e.g. turbidity, flow), light, oxygen, pH, as well as substrate availability. These factors can alter the pathways by which nutrients are released from the sediment, or the rates by which they are transformed in the sediment or water column by such processes as nitrification, denitrification, anammox, or dissimilatory nitrate reduction to ammonium (DNRA).


In eutrophic systems, increased algal productivity may lead to depressed water column oxygen which may result in increased recycling of N and P by changes in redox potential, or pH [29<sup>••</sup>,30,31]. These fluxes can positively reinforce an ecosystems' degradation trajectory, as suggested to be the case for the Chesapeake Bay [31]. By contrast, in systems with a reduced nutrient load, lower algal abundance in the water column, higher light, and higher redox potential may help to reinforce higher rates of nitrification and denitrification, leading to nutrient removal and potentially eutrophication reversal [31]. Biogeochemical pathways together with homeostatic control serve to provide the mechanism(s) whereby nutrient dynamics support trophodynamic structure. Positive reinforcing feedbacks of biogeochemistry and homeostasis shift ecosystems to new stable states; such shifts can be gradual or abrupt and communities may not return to their original state once the disturbance (in this case, altered nutrient loads) is removed.

Altered ecological stoichiometry and associated ecological feedback mechanisms may also help to explain the potential for a system to support the growth of organisms previously not found in an area, that is, biological invasions. While changes in climate, land-use and other factors have long been recognized to be related to a system's susceptibility for invasive species to thrive, so too is the possibility that altered nutrient loads may make an environment more suitable for species with different nutritional requirements [29<sup>••</sup>]. Such a suggestion does not negate the importance of external vectors bringing invaders to a new area, but nutritional loads, and the associated ecological feedbacks that may alter habitat, adds a mechanism to explain why some shifts occur when they do, and when and why particular species are successful in new environments [29<sup>••</sup>].

### Examples from comparative systems

Two estuarine examples illustrate the important feedbacks between changes in nutrient loads and their stoichiometry, biogeochemistry, food web structure and potential for biological invasions. The San Francisco Bay Delta, a heavily modified and anthropogenically impacted estuarine system, has had major changes in nutrient loads over the past several decades ([29<sup>••</sup>,32] and references therein). In particular, increases over time in the N:P ratio of the major nutrient loads have been attributed, at least partly, to increases in wastewater N and removal of P from laundry detergents [29<sup>••</sup>]. Coincident with, and related to, these changes are declines in water column algal biomass (as chlorophyll *a*), a shift in the dominant zooplankton species from the calanoid *Eurytemora affinis*, to the invasive cyclopoid *Limnithona tetraspina*, as well as invasions of the Asiatic clam, *Corbula amurensis*, the macrophyte, *Egeria dense*, and various species of centrarchid fish [29<sup>••</sup>]. *Egeria dense* is thought to do particularly well under increasing water column N:P

Figure 1



Conceptual depiction of the change over time in major nutrients, flow, dominant biogeochemical processes, and the food web of the San Francisco Bay Delta illustrating how multiple feedbacks serve to alter food web structure when nutrient loads are altered. The first panel represents the period from 1975 to ~1982, when flow was low, and diatoms were the dominant phytoplankton, and the calanoid copepod *Eurytemora* was the dominant copepod, and pelagic planktivorous fish were common. The second panel represents the period from ~1982 to 1986 when flow was high, and  $\text{NH}_4^+$  was increasing. During this period the food web began to change. Under very low flow conditions, depicted by the third panel, and representing ~1987–1995, the  $\text{NH}_4^+$  load was high but  $\text{PO}_4^{3-}$  began to decrease. The food web also began to change significantly, with changes in the dominant phytoplankton and zooplankton, increasing abundance of macrophytes, increased importance of sediment nutrient processes, and increase in piscivores. Finally, post 1995,  $\text{NH}_4^+$  loads remain high, while  $\text{PO}_4^{3-}$  loads have decreased. Sediment biogeochemical processes are of increasing importance in nutrient processing, macrophyte production is important and piscivorous fish have increased. Reproduced from Glibert *et al.* [29\*\*] with permission of the publisher.

#### 4 Aquatic and marine systems

loads because it can access P through the sediments, and can tolerate high N, especially in the form of  $\text{NH}_4^+$ , in the water column. It also serves as an 'ecological engineer', decreasing nutrients through uptake, reducing turbidity by trapping sediments, and providing refuge for zooplankton and habitat for fish such as largemouth bass [33]. More importantly from a chemical perspective, it is able to use bicarbonate effectively through a well-developed carbon-concentrating mechanism (e.g. [34]). The consequence is that it does not become C-limited even under periods of high productivity, in turn elevating pH on a diel basis in the surrounding water as it grows, in some cases substantially. Once pH is elevated, the fundamental physical-chemical relationships related to P adsorption-desorption in the sediment change, as does N biogeochemistry [35,36]. Moreover, under increased pH conditions, the biogeochemistry of calcification is altered, increasing the potential for calcification and the growth of calcifying organisms. The change in the abundance of the clam *C. amurensis* from the time of its introduction in the mid-1980s to 2005 has been shown to be highly and positively correlated to the increase in total N:total P ( $r^2 = 0.46$ ;  $n = 20$ ;  $p < 0.01$ ; all data log transformed), and the average annual abundance of this species has also been found to be highly and positively correlated with mean annual average pH in the estuary ( $r^2 = 0.64$ ;  $n = 19$ ;  $p < 0.01$ ; species abundance data log transformed) [29\*\*]. Thus, over time nutrient loading changes, biogeochemical changes and foodweb changes were intertwined (Figure 1).

The River Rhine presents a parallel example [37,38]. From 1977 to 2005, total P loads declined over 6-fold owing to reductions in wastewater loading from municipal and industrial facilities and the total N:total P ratio increased 3-fold, in turn resulting in an 8-fold decline in water column chlorophyll *a* [38]. Cumulative non-indigenous species increased from ~20 to >50 during these years, a rate of accumulation higher than seen in any year of record, dating back to 1825 [37]. Most of these species were molluscs (22% of total) and crustacea (51% of total); among the molluscs, the most abundant were the clam *Corbicula fluminea* and the mussel *Dreissena polymorpha* (zebra mussel) [37]. The annual rate of change in number of animal non-indigenous species and the rate of change in the ratio of total N:total P in the water column over this ~30-year time period are highly and positively correlated ( $r^2 > 0.40$ ;  $n = 28$ ;  $p < 0.05$ ). In addition, invasive macrophytes, including *E. dense*, are increasingly common in the Rhine [39], as they are in the San Francisco Bay Delta.

While there are many vectors that may deliver non-indigenous species to new areas, there is no question that altered biogeochemical composition of the water column, and the associated shifts in primary production can set in motion a cascade of changes that facilitate the

success of new species. Thus, both the Rhine and the Bay Delta had similar types of ecosystem changes following changes in N:P stoichiometry: significant declines in chlorophyll *a*, invasions of *E. dense*, bivalve clams, and changes in crustacea. These examples illustrate that when P is reduced relative to N, and when the system production becomes more benthic in nature, new dominants may emerge. In the case of filter-feeding molluscs, they, in turn, may help to maintain the new water quality condition by enhanced filtration of the water column.

#### Conclusions and implications

Nutrient stoichiometric control of ecosystem structure and sustainability must be viewed within the context of other factors in the environment, including the multiple stressors that now impact systems, as well as the scale (spatial, temporal and organismal; proximate or ultimate) on which the system is being examined. Trophodynamics and biogeochemical processes are fundamentally coupled, but are also fundamentally constrained by the availability and composition of elements relative to the needs of the organisms. Ecological stoichiometry affects ecosystems by setting elemental constraints on the growth of organisms, with several important implications for the health and sustainability of aquatic systems. Stoichiometric imbalances may accelerate transformations of nutrients or may alter the processes by which nutrients are cycled in the ecosystem, further altering nutrient availability or form for primary producers [28,40]. Food *quality* is linked to food web outcome via feedback effects and nutrient biogeochemistry, key processes determining biodiversity. Models of ecosystem function based on classic nutrient or growth kinetic relationships, or flow of single elements (i.e. C), are ill-suited to capture the complexities of stoichiometric effects on ecosystems [41]. Although far more complex than single currency models, multiple currency models and ecological stoichiometric concepts should be among the considerations that should go into a new generation of ecosystem models [11,29\*\*,42\*,43].

Ecological stoichiometry should be incorporated into our thinking, management, and modeling of the dynamics of nutrient-impacted systems. Imbalances in stoichiometry may have impacts on ecosystems even at nutrient loads normally taken to be saturating or supersaturating. The effect of stoichiometric constraints has implications, for example, on bioenergetics and all aspects of fish nutrition, both from a basic ecological perspective as well as applied, including aquaculture [21\*\*,44]. Moreover, ecological stoichiometry bears significantly on the debate of whether aquatic ecosystem restoration efforts should focus on P removal, N removal, or both (e.g. [45,46,47\*\*] and references therein). Single nutrient removal strategies drive ecosystems into states of stoichiometric imbalance [29\*\*]. Imbalances in stoichiometry may destabilize the dynamics of consumers, shifting systems to new conditions. Single

nutrient removal strategies may have unintended consequences for aquatic ecosystems.

## Acknowledgements

The author thanks I. Perry, A. Bundy, E. Hofmann, and M. Barange for the invitation to submit this paper, and D. Fullerton, J. Burkholder, J. Cornwell, and T. Kana for numerous discussions and insights regarding these relationships. The author received support from the State and Federal Contractors Water Agency and the Delta Stewardship Council during the preparation of this review; the funding agencies had no involvement in its preparation, the decision to submit, nor in the interpretation of any data reported herein. This is contribution number 4653 from the University of Maryland Center for Environmental Science.

## References and recommended reading

Papers of particular interest, published within the period of review, have been highlighted as:

- of special interest
- of outstanding interest

1. Doney SC: **The growing human footprint on coastal and open ocean biogeochemistry.** *Science* 2010, **328**:1512-1516.  
This paper reviews the effects of human impacts on ocean biogeochemistry and the projections for further global scale changes in future decades.
2. Howarth RW: **Coastal nitrogen pollution: a review of sources and trends globally and regionally.** *Harmful Algae* 2008, **8**:14-20.
3. Van Drecht G, Bouwman AF, Harrison J, Knoop JM: **Global nitrogen and phosphate in urban waste water for the period 1970-2050.** *Global Biogeochem Cycles* 2009, **23**:GB0A03 <http://dx.doi.org/10.1029/2009GB003458>.
4. Johnson P, Townsend A, Cleveland CC, Glibert PM, Howarth R, Mackenzie V, Rejmankova E, Ward M: **Linking environmental nutrient enrichment and disease emergence in humans and wildlife.** *Ecol Appl* 2010, **20**:16-29.
5. Anderson DA, Glibert PM, Burkholder JM: **Harmful algal blooms and eutrophication: nutrient sources, composition, and consequences.** *Estuaries* 2002, **25**:562-584.
6. Cloern JE: **Our evolving conceptual model of the coastal eutrophication problem.** *Mar Ecol Prog Ser* 2001, **210**:223-253.
7. Vitousek P, Porder S, Houlton BZ, Chadwick OA: **Terrestrial phosphorus limitation: mechanisms, implications, and nitrogen-phosphorus interactions.** *Ecol Appl* 2012, **20**:5-15.
8. Allen AP, Gillooly J: **Towards an integration of ecological stoichiometry and the metabolic theory of ecology to better understand nutrient cycling.** *Ecol Lett* 2009, **12**:369-384.  
This paper discusses the integration of the metabolic theory of ecology and ecological stoichiometry and its guiding principles.
9. Falkowski PG: **Rationalizing elemental ratios in unicellular algae.** *J Phycol* 2000, **36**:3-6.
10. Klausmeier CA, Litchman E, Daufresne T, Levin SA: **Phytoplankton stoichiometry.** *Ecol Res* 2008, **23**:479-485.  
This paper reviews phytoplankton stoichiometry in the context of classic kinetic models for nutrient uptake and provides an overview of how multiple nutrient models can advance our understanding of oceanic stoichiometry and its homeostasis.
11. Flynn KJ: **Do external resource ratios matter? Implications for modelling eutrophication events and controlling harmful algal blooms.** *J Mar Syst* 2010, **83**:170-180.
12. Sterner RW, Elser JJ: *Ecological Stoichiometry: The Biology of Elements from Molecules to the Biosphere.* Princeton, NJ: Princeton University Press; 2002.  
This book considers the principles of ecological stoichiometry from the fundamental building blocks of elements to macroscale perspectives. The foundational ideas are presented.
13. Granéli E, Edvardsen B, Roelke DL, Hagström JA: **The ecophysiology and bloom dynamics of *Prymnesium* spp.** *Harmful Algae* 2012, **14**:260-270.
14. Mitra A, Flynn KJ: **Predator-prey interactions: is "ecological stoichiometry" sufficient when good food goes bad?** *J Plankt Res* 2005, **27**:393-399.
15. Glibert PM, Burkholder JM: **Harmful algal blooms and eutrophication: strategies for nutrient uptake and growth outside the Redfield comfort zone.** *Chin J Limnol Oceanogr* 2011, **29**:724-738.
16. Irigoien X, Flynn KJ, Harris RP: **Phytoplankton blooms: a 'loophole' in microzooplankton grazing impact?** *J Plankt Res* 2005, **27**:313-321.
17. Mitra A, Flynn KJ: **Promotion of harmful algal blooms by zooplankton predatory activity.** *Biol Lett* 2006, **2**:194-197.
18. Sunda WG, Granéli E, Gobler CJ: **Positive feedback and the development and persistence of ecosystem disruptive algal blooms.** *J Phycol* 2006, **42**:963-974.
19. Finkel ZV, Beardall J, Flynn KJ, Quiqq A, Rees TA, Raven JA: **Phytoplankton in a changing world: cells size and elemental stoichiometry.** *J Plankt Res* 2010, **32**:119-137.  
This paper takes the perspective that phytoplankton cell size and stoichiometry have important implications for growth rates, food web structure and biogeochemical cycling especially in world where CO<sub>2</sub> is changing and acidification of the oceans is occurring.
20. Geider RJ, La Roche J: **Redfield revisited: variability of C:N:P in marine microalgae and its biochemical basis.** *Eur J Phycol* 2002, **37**:1-17.
21. McIntyre PB, Flecker AS: **Ecological stoichiometry as an integrative framework in stream fish ecology.** *Am Fish Soc Symp* 2010, **73**:539-558.  
This paper presents an overview of the principles of ecological stoichiometry and how they apply across individual, population, community and ecosystem levels in fish ecology.
22. Boersma M, Aberle N, Hantzsche FM, Shoo KL, Wiltshire KH, Malzahn AM: **Nutritional limitation travels up the food chain.** *Int Rev Hydrobiol* 2008, **93**:479-488.
23. Hessen DO, Andersen TR: **Excess carbon in aquatic organisms and ecosystems: physiological, ecological, and evolutionary implications.** *Limnol Oceanogr* 2008, **53**:1685-1696.
24. Malzahn AM, Hantzsche F, Shoo KL, Boersma M, Aberle N: **Differential effects of nutrient-limited primary production on primary, secondary or tertiary consumers.** *Oecologia* 2010, **162**:35-48.
25. Shoo KL, Aberle N, Malzahn AM, Boersma M: **Does the nutrient stoichiometry of primary producers affect the secondary consumer *Pleurobrachia pilleus*?** *Aquat Ecol* 2010, **44**:233-242.
26. Vanni MJ: **Nutrient cycling by animals in freshwater ecosystems.** *Annu Rev Ecol Syst* 2002, **33**:341-370.
27. Sterner RW, Hessen DO: **Algal nutrient limitation and the nutrition of aquatic herbivores.** *Annu Rev Ecol Syst* 1994, **25**:1-29.
28. Nugraha A, Pondaven P, Tréguer P: **Influence of consumer-driven nutrient recycling on primary production and the distribution of N and P in the ocean.** *Biogeosciences* 2010, **12**:1285-1305.
29. Glibert PM, Fullerton D, Burkholder JM, Cornwell JC, Kana TM: **Ecological stoichiometry, biogeochemical cycling, invasive species and aquatic food webs: San Francisco Estuary and comparative systems.** *Rev Fish Sci* 2011, **19**:358-417.  
Using principles of eutrophication, ecological stoichiometry and stable state theory, this paper presents an overall framework for considering how long-term changes in the food web of the San Francisco Bay Delta and other comparative ecosystems have resulted from changes in nutrients loads and forms.
30. Gao Y, Cornwell JC, Stoecker DK, Owens MS: **Effects of cyanobacterial-driven pH increases on sediment nutrient fluxes and coupled nitrification-denitrification in an shallow fresh water estuary.** *Biogeosci Discuss* 2012, **9**:1-38.
31. Kemp WM, Boynton WR, Adolf JE, Boesch DF, Boicourt WC, Brush G, Cornwell JC, Fisher TR, Glibert PM, Hagy JD *et al.*: **Eutrophication in Chesapeake Bay: historical trends and ecological interactions.** *Mar Ecol Prog Ser* 2005, **303**:1-29.

## 6 Aquatic and marine systems

32. Kimmerer WJ: **Open water processes of the San Francisco Estuary: from physical forcing to biological responses.** *San Francisco Estuary Watershed Sci* 2004, **2** scholarship.org/uc/item/9bp499mv.
33. Yarrow M, Marin VH, Finlayson M, Tironi A, Delgado LE, Fischer F: **The ecology of *Egeria densa* Planchón (Liliopsida: Alismatales): a wetland ecosystem engineer?** *Rev Chil Hist Nat* 2009, **82**:299-313.
34. Bowes G: **Aquatic plant photosynthesis: strategies that enhance carbon gain.** In *Plant Life in Aquatic and Amphibian Habitats*. Edited by Crawford RMM. Oxford: Blackwell Scientific; 1987:79-89. Special Pub No 5 of the British Ecol Soc.
35. Bange HW: **Gaseous nitrogen compounds (NO, N<sub>2</sub>O, N<sub>2</sub>, NH<sub>3</sub>) in the ocean.** In *Nitrogen in the Marine Environment*. Edited by Capone DG, Bronk DA, Mulholland MR, Carpenter EJ. London: Elsevier; 2008:51-94.
36. Jordan TE, Cornwell JC, Boynton WR, Anderson JT: **Changes in phosphorus biogeochemistry along an estuarine salinity gradient: the iron conveyer belt.** *Limnol Oceanogr* 2008, **53**:172-184.
37. Leuven RSEW, van der Velde G, Bajjens I, Snijders J, van der Zwart C, Lenders HJR, bij de Vaate A: **The river Rhine: a global highway for dispersal of aquatic invasive species.** *Biol Invasions* 2009, **11**:1989-2008.
38. Van Nieuwenhuysse E: **Response of summer chlorophyll concentration to reduced total phosphorus concentration in the Rhine River (Netherlands) and the Sacramento-San Joaquin Delta (California, USA).** *Can J Fish Aq Sci* 2007, **64**:1529-1542.
39. Nienhuis PH: *Environmental History of the Rhine-Meuse Delta*. Springer Science and Business Media, B.V.; 2008.
40. Elser JJ, Hamilton A: **Stoichiometry and the new biology: the future is now.** *PLoS Biol* 2007, **5**:e181 <http://dx.doi.org/10.1371/journal.pbio.0050181>.
41. Flynn KJ: **Ecological modelling in a sea of variable stoichiometry: dysfunctionality and the legacy of Redfield and Monod.** *Prog Oceanogr* 2009, **84**:52-65.
42. Allen JI, Polimene L: **Linking physiology to ecology: towards a new generation of plankton models.** *J Plankt Res* 2011 <http://dx.doi.org/10.1093/plankt/fbr032>.  
This paper outlines the need for incorporation of underlying cell physiology and food web interactions in ecosystem models in order to understand and quantify emergent properties – a departure from numerical parameter fitting.
43. Glibert PM, Allen JI, Bouwman L, Brown C, Flynn KJ, Lewitus A, Madden C: **Modeling of HABs and eutrophication: status, advances, challenges.** *J Mar Syst* 2010, **83**:262-275.
44. Hendrixson HA, Sterner RW, Kay AD: **Elemental stoichiometry of freshwater fishes in relation to phylogeny, allometry and ecology.** *J Fish Biol* 2007, **70**:121-140.
45. Carpenter SR: **Phosphorus control is critical to mitigating eutrophication.** *Proc Natl Acad Sci USA* 2008, **105**:11039-11040.
46. Conley DJ, Paerl HW, Howarth RW, Boesch DF, Seitzinger SP, Havens KE, Lancelot C, Likens GE: **Controlling eutrophication: nitrogen and phosphorus.** *Science* 2009, **323**:1014-1015.
47. Duarte CM, Conley DJ, Carstensen J, Sánchez-Camacho M: **Return to Neverland: shifting baselines affect eutrophication restoration targets.** *Estuaries Coasts* 2008, **32**:29-36.  
This paper presents a synthesis of systems that have undergone nutrient loading increases and decreases and the challenges of returning systems to their pre-enrichment conditions.